

This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

#### Usage guidelines


Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:


- + *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.
- + Refrain from automated querying Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.
- + *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.
- + *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

#### **About Google Book Search**

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at http://books.google.com/









## THE

# INFLUENCE OF TROPICAL CLIMATES

ON

EUROPEAN CONSTITUTIONS.



# INFLUENCE OF TROPICAL CLIMATES

ON

# EUROPEAN CONSTITUTIONS,

INCLUDING

PRACTICAL OBSERVATIONS ON THE NATURE AND TREATMENT
OF THE DISEASES OF EUROPEANS ON THEIR RETURN
FROM TROPICAL CLIMATES.

BY

## JAMES RANALD MARTIN, F.R.S.,

SURGEON BENGAL ARMT, RETIRED; LATE PRESIDENCY-SURGEON, AND SURGEON TO THE NATIVE HOSPITAL, CALCUTTA; MEMBER OF THE BOYAL ASIATIC SOCIETY OF GREAT BEITAIN AND IRELAND, AND OF THE ASIATIC SOCIETY OF BENGAL, ETC. ETC.

A NEW EDITION.




LONDON:

JOHN CHURCHILL, NEW BURLINGTON STREET.

MDCCCLVI.

45



## PREFACE.

When, little more than twelve months ago, I was called upon to prepare anew the work composed by the late Dr. James Johnson and myself, in 1841, on the "Influence of Tropical Climates on European Constitutions," I felt that a mere reprint could not in justice be proposed as a Seventh Edition. Many years had elapsed since Dr. Johnson's original and admirable work on the subject first appeared. The additions in subsequent re-publications consisted chiefly of his able reviews of the labours and researches of other authors on tropical diseases, which appeared in the "Medico-Chirurgical Review." The work thus became in great part a collection of reviews, which, excellent as they were at the time, could not now be made to represent the advancing knowledge of the day.

I have therefore ventured to re-cast and re-write the entire work. Even the introductory chapter is entirely re-written—so much only of Dr. Johnson's observations being retained as appeared to me proper to the subject, and all beyond is exclusively my own composition.

I am well aware of the responsibility I have thus reluctantly assumed. Throughout the present Edition I have adhered to, and preserved the medico-topographic and statistical character of my original Official Reports, which were printed and circulated at the public cost in Bengal, by order of the Supreme Government of India, while I was serving under its orders. In doing so, I am

actuated by the conviction that, next to the importance of physical investigations, stand those which relate to the discovery the EXTERNAL CAUSES OF DISEASE. By such a course of inqui we shall, I think, most readily appreciate both the prevention at cure of those formidable diseases to which Europeans are subject in hot climates.

I have likewise continued, so far as the scope and intent of the work admits, an examination of the physiological influences climate, both Asiatic and European, while every suggestion offered which seemed to me likely to enhance the value a importance of PREVENTIVE MEDICINE.

For the discussion of the nature, causes, and treatment tropical diseases, my opportunities, whether rightly or wrong applied, have been both varied and extended. I served in varies of India—in peace and war—amongst natives and Europ—in hospital and in private practice—for two-and-twenty and during this time I did my best to collect materials in the field which came under my observation; I wish much that I command more leisure for the arrangement and completion subjects.

A treatise purporting to describe "The Influence of Climates on European Constitutions," which should not some description of the sequelæ to tropical diseases, a counter-influences which, under an European climate, health of persons who have resided long in warm reg be obviously incomplete. I therefore here reprint Lancet" certain articles published by me during years, on "The Nature, Causes, and Treatment of th Europeans on their Return from Tropical Climates of such a supplementary treatise was felt by my fri Johnson, who urged me, on my return from India taking; and having now been several years in extion of this class of invalids, I venture on the duty which my friend proposed to me. In corefrain from offering a tribute of respect to the

author of the "Influence of Tropical Climates on European Constitutions," originally published in 1813, from materials collected and arranged by him when he was not twenty-four years of age. Having served not quite three years in the Indian seas, as surgeon to a ship-of-war, he had at this time returned home with broken health, several abscesses having been discharged from the liver, rendering his life one of extreme suffering for years thereafter. By great native elasticity of mind and body, he overcame his distresses and difficulties, and ended his career as an esteemed and eminent physician of this metropolis.


Referring to the subject of the sixth edition of the work, composed jointly by Dr. Johnson and myself, in 1841, he observes, with a just pride, that "Although successive editions have received additions and improvements, yet, very few of the Author's original doctrines have been subverted, or practices exploded, during a period of more than a quarter of a century."

Few men in our profession could say so much; and I speak but the general sentiment when I say, that fewer still have departed from us, leaving a higher reputation for kindliness of feeling, and high sense of honour, than did Dr. James Johnson.

To his talented son, Mr. Athol Johnson, I am under much obligation for the aid he has given me in revising the work while being printed.

J. R. MARTIN.


Grosvenor Street, Grosvenor Square, December, 1855.



# CONTENTS.

# PART I.

		P	AGE.
SKETCH OF THE PHYSICAL CLIMATE OF CALC	UTTA,	IN	-
BENGAL, WITH GENERAL REMARKS ON THE CAUSES	OF	PHY-	
SICAL CLIMATE			1
Action of the Sun upon the Atmosphere	16.1		5
Internal Heat			7
Elevation of the Ground			9
General and Local Aspects			10
Position of Mountains			11
Effects of the Neighbourhood of the Sea	-		11
Geological Nature of the Soil			13
Influence of the Labour of Man-General Population			20
The Predominant Winds			22
The Rains—Sources of Aqueous Exhalations—Humidity		-	25
The River			27
The Salt Lake			29
Woods and Marshes of the Sunderbunds			30
MEDICAL CLIMATE AND THE INFLUENCE OF SEASON	100		33
Physical Effects of Climate			36
The Hot Season and its Effects			38
The Rainy Season			42
The Cold Season	-	8	
STATISTICS OF THE INFLUENCE OF CLIMATE AND SEASON	IN PE	LODUC-	
ING SICKNESS AND MORTALITY IN CALCUTTA.	-		52
Mortality of the various Races, European and Native .			59
Mortality of British Soldiers and Seamen		1	74



						AGE.
Catalogue—raisonnè of the			ttent Feve	ers by diff	erent	
Authors, in different ag		ntries				178
General Remarks in Conclus	ion .					180
INTERMITTENT FEVER						
Historical and General Notic	ces .	-				185
Causes			-		6	187
Symptoms and Progress			4			187
Pathology						189
Treatment						192
THE CONGESTIVE CONTINU	UPD PRVI	P OF TI	IF COLD	SEASON	JIN	
BENGAL .	DED PEAL	or II	IE COLD	SEASOI	, 114	200
BENGAL						200
THE ARDENT CONTINUED	FEVER	OF THE	нот	SEASON	IN	
BENGAL						204
THE FEVERS OF THE NATIVI	S OF INT	OTA				
General Topographical Obse			dia .			210
Physical Character of its In		Direish In	uia .			212
State of Agriculture and the			- 1			213
Native Armies—Character		or of Ron	en]			214
Official and Social Position						215
British Officers of the Indian		car Omcer			100	215
Native Army of Bengal, its s	The state of the s					700
Average Sickness and Morta		. V.		1.0		216
		enty lears	1	-		216
Native Army of Madras, its			*			217
Its average Mortality and In	validing				1.6	217
Native Army of Bombay						217
Its average Mortality and I						217
Liability of the Native Troc						217
Influence of change of Clin				. 18		217
Seasons of Unhealthiness co			European	ns .	1.8	220
Fevers constitute the first in		000000000000000000000000000000000000000	*			220
Treatment of the Fevers of	the Native	8 .				221
ACUTE DYSENTERY OF BENC	AL					
General and Historical Obse						222
Predisposing Causes .				-		223
Symptoms and Progress						224
Complications .		-			1	226
Hepatic complication .			10			227
Splenic			- 3			229
m at at		-	10			229
Pathology .	-		- 2	2		230
Treatment of acute Tropica	Dysantern		1			233
reatment of acute fropica	T PASCHICE A	140	*		- 4	400

	xii		CONTEN	TS
STATE OF THE PERSON NAMED IN	Treatmen	of Hæmorrhagic	Dysentery	
	,,	Hepatic Dyser		4
	n	Scorbutic "		,
		Dysentery, co		with
	mitte			
	***	Enlargement of		n .
	"	Convalescence	7	
		-raisonne of the		-
		countries, by differ		
	General C	bservations in con	clusion	1
	CHRONIC DYS	ENTERY .		
		in the Natives of	India	
	DISEASES OF	THE LIVER		
		Secretion of Bile		1
	Treatmen		2	
		n of the Liver	-	
	Treatment			
	Hepatitis			
		and Progress		
	Causes			
	Pathology			
	Treatmen			
	Hepatic A	bscess .		
	Treatment			
	Jaundice	(6) (6)		
	Causes		*	
	Pathology			
	Treatmen			
2		nlargement and Ir	iduration of	the
	Treatmen			
1913		EMIC CONGESTI	ON OF TH	HE
100	Symptom			
100				
	Pathology			
100	Treatmen			
	EPIDEMIC CH			
12		the Earlier Indian		
11/1		the Medical Boar		
		cal Progress of Ep	idemic Chol	lera
		and Progress	2	
	Predisposi	ng Causes .		

#### CONTENTS.

								AGE.
	Comparative duration	n of fatal	Diarrhœa	, Summer	Cholera, a	nd Epidem	ic	
	Cholera			*		Acres .		302
	Relation of elevation							303
	Diagram representing					elevations		305
	The six theories as t				Cholera			306
	Dr Baly's Report to	the Colleg	e of Physi	icians				307
	Preventive Measures		,					314
	17 11	In the Ar				0 1		320
	Morbid appearance a	fter death	in Collap	se		4		325
	Pathology of Cholera						-	330
	Treatment of Cholers	-Dr. Gu	ll's Repor	t to the Co	llege of P	hysicians		335
	Epidemic Cholera in	Calcutta						343
	Cholera Fever of 183	4		4				344
	Influence of high ten	perature	4					345
	,, Season							346
	Predisposing Causes							348
	Premonitory Diarrho	ea						348
	Pathology .			2				349
	Treatment .		41					350
	Tables of average S	ickness ar	d Mortal	ity from C	holera in	our foreig	gn	
	possessions	*						354
HE	YELLOW, OR HÆM General and topograp			R		4		356
	Symptoms and Prog				4			360
	Sequelæ .							362
	The Black Vomit							362
	Diagnosis .							363
	Prognosis .							365
	Mortality .					1	-	366
	Post mortem appeara	nces			0			368
	Predisposing Causes					-		370
	Endemic Causes					1		371
	Infection .							375
	Prevention .							382
	Cure					i i	1	386
	outo .	-			-	*		000
EL	IRIUM TREMENS.							
40.84	Symptoms and Progr	ress			14		V.	392
	Causes .		1	1	2	5		394
	Complications							395
	Pathology .		16	154		-	1.	395
	Prevention .		-		10.	16		397
	Curative Treatment							398

	xiv CONTENTS.
	PAGE.
	HABITS OF THE BRITISH SOLDIERY AND THEIR INFLUENCE ON
	HEALTH
	Barrack Life
	Necessity of Exercise
	Intemperance
	ON THE SELECTION AND IMPROVEMENT OF LOCALITIES FOR THE EUROPEAN TROOPS IN OUR INTERTROPICAL POSSESSIONS. LOCALITIES TO CHOOSE, LOCALITIES TO AVOID 413
March 1997	Memorandum relating to the construction and arrangement of barracks
	and hospitals
	Table of Mortality in town and country throughout the different seasons
	of the year
	Neglect of Army Medical Officers by the Authorities
	,, Avoid
	, Atold
	ON THE TERM OF EFFICIENT SERVICE IN INDIA
	General Observations
	CHILDREN IN BENGAL
	PART II.
	PRACTICAL OBSERVATIONS ON THE NATURE AND TREATMENT OF THE DISEASES OF EUROPEANS ON THEIR RETURN FROM TROPICAL
S111	CLIMATES
	Introductory Chapter .
	Anæmia .
	Cases—Treatment
	ON THE TREATMENT OF INVALIDS SUFFERING FROM THE SEQUITOR TO THE FEVERS OF TROPICAL CLIMATES
	TO THE FEVERS OF TROPICAL CAMARIE
	Cases
	CHRONIC DIABRHEA Cases—Treatment
	Cases—Treatment

					P/	GE.
CHRONIC DYSENTERY		•	•	•		<b>525</b>
Table of Statistics .			•	•	•	527
Symptoms		· .				530
Cases—Treatment .	•			•		531
Congestion of the Live	R.	•				548
Cases—Treatment .		•				550
TORPOR OF THE LIVER	•		•	•		558
Symptoms			•	•	•	559
Treatment						561
Nitro-muriatic acid bath		•	•			564
Mode of using .	•	•	•	•		564
Cases		•		•	•	568
CHRONIC ENLARGEMENT O	F THE I	IVER.	•			575
Treatment		•	•			578
Cases				•	•	579
CONCLUSION .			_	_		597

CONTENTS.

xv

## ON CLIMATE.

# SKETCH OF THE PHYSICAL CLIMATE OF CALCUTTA IN BENGAL,

WITH GENERAL REMARKS

## ON THE CAUSES OF PHYSICAL CLIMATE.

"L'ensemble de toutes les circonstances naturelles et physiques, au milieu desquelles nous vivons dans chaque lieu."—Cabanis.

"The best observations upon climate, often lose half their value from want of an exact description of the surface of the country."—Malte Brun,

Whoever considers climate, with reference to its vast importance to human welfare, must feel some degree of disappointment at the meagreness in which the advanced state of knowledge in the nineteenth century has yet left this most interesting branch of

inquiry.

On this all-important subject it is but too evident that, in the language of Sir John Herschel, we have not yet determined "what ought to be observed," and thus it has happened, for want of "a right direction," that much of the labour of observation and minute investigation has been "but labour lost." In medical pursuits, we find continually that the apparent cause of sickness, or of death, is not the real cause; and so it is with the causes of physical climate. "When a thing is accounted for in two ways," says Fontenelle, "the truth is usually on the side most opposed to appearances."

their being comparable, and this in an especial manner applies to inquiries relating to medical topography and medical statistics. I am satisfied that, in a professional sense, it is impossible to take too extended a view of the subject of climate, and that he who succeeds best must follow the indication of Cabanis; "L'ensemble de toutes les circonstances naturelles et physiques, au milieu desquelles nous vivons dans chaque lieu;" for this much is certain, that the framers of elaborate tables of the winds and of the degrees of the thermometer, have as yet done little to inform our minds, or guide our inquiry. "Le climat n'est donc point resserré dans les circonstances particulieres des latitudes, ou du froid et du chand; il embrasse, d'une maniere absolument générale, l'ensemble des circonstances physiques attachées a chaque local, il est cet ensemble lui-même; et tous les traits caractéristiques par lesquels la nature a distingué les differents pays entrent dans l'idée que nous devons former du climat."

Medical men, to whom a right understanding of the subject is paramount, whether we consider the good or the evil to be derived from climate, are happily beginning to be aware of this fact; and it is with much satisfaction, I find one of the latest and best writers on medical topography in England, declaring, in furtherance of the just and comprehensive views above quoted, that the character of a climate is much more faithfully indicated by such a natural test as its influence on vegetable products, "than by any

instrumental, or artificial means whatever."

Dr. Mason Good, asserts, with truth, that it is no exclusive reproach to medicine that meteorology has hitherto been turned to so little practical advantage; "for of all the subdivisions of general philosophy, there is none so little entitled to the name of science as meteorology itself. And, till the naturalist has explained the variations of the barometer, the physician need not blush at being incapable of turning to account the supposed influence of the planets, or unfolding the origin, or tracing the capricious courses of epidemics and pestilence." Difficult as is the subject, however, it ought to have its great attractions to the physician; for, assuredly, no other affords so free a scope for original observation and inquiry; and this applies in an especial manner to tropical climates. Dr. Watson observes justly, that "to ascertain the causes of any disease, and to display them before the public mind, are large steps towards the ultimate removal of such as human endeavours are competent to remove." . . . . "To know the cause of a disease is sometimes to be able to cure, often to be able to prevent it."

Malte Brun's division of physical climate shall here be adopted, because it marks the geographical points to which our inquiries may most profitably be directed, inasmuch as a consideratio the details, as given by that admirable writer, brings us at once the discovery of such errors in the physical condition of any giv place, as may lead to its improvement, and the consequent prevention of disease—the great end of medical topography.

When again, we look back to our native country, and boast its pure and bracing air, let us not forget the important fact, "tl it is man himself who has in a great measure created these sal brious climates. France, Germany, and England, not more th twenty ages ago, resembled Canada and Chinese Tartary, countr situated, as well as our Europe, at a mean distance between equator and the pole."

"Whenever nature is more powerful than industry, whether good or for bad, man receives from the climate an invariable irresistible impulse;" and so it is in Bengal, 'for bad,' in quence of the total absence of that 'industry' which elemonds nature to the purposes of man, for his 'good.'

Physical climate, says Malte-Brun, comprehends the heat and cold, the drought, the humidity, and the salul occur in any given region of the earth.

The causes of physical climate he states to be ni ber:-

1st. The action of the sun upon the atmosphere.

2nd. The interior temperature of the globe.

3rd. Elevation of the earth above the level of the

4th. The general inclination of the surface a posure.

5th. The position of its mountains relative)

6th. The neighbourhood of great seas and tion.

7th. Geological nature of the soil.

8th. Degree of cultivation and of populati has arrived.

9th. The prevalent winds.

To these perhaps may be added:-

10th. Position in respect to the equate

11th. Position in respect to large rive 12th. Position in respect to forests.

By tracing these causes, and by ur general points of view, the results of p we shall, according to Malte-Brun, climatology, in some measure corresp the other sciences.

#### ACTION OF THE SUN UPON THE ATMOSPHERE.

At all places between  $20^{\circ}$  and  $23\frac{1}{2}^{\circ}$  of latitude, the solar rays during two months fall at noon either perpendicularly, or at an angle which deviates from a right angle only by  $3\frac{1}{2}^{\circ}$  at most. The degree of temperature is also increased by the greater length of the longest days, which near the tropic, are  $13\frac{1}{2}$  hours. The application of these facts to the geographical position of Calcutta during portions of May, June, July, and August, are sufficiently obvious, and would not seem to require the lengthened details offered by some writers.

The difference between sensible heat and that indicated by instruments, is nowhere more remarkable than within this city during the hot months, where, from the length of time the houses, walls, and roads are getting heated by the sun, the first hours of the night are rendered even more oppressive than the day, from the copious radiation going on for some time after the sunset. "The true indication of the force of the solar rays," says Herschel, "would seem to be, not the statical effect upon the thermometer, but their momentary intensity measured by the velocity with which they communicate heat to an absorbent body."

Professor Daniel's observations also go to show that the power of solar radiation in the atmosphere increases from the equator to the poles, and from below upwards.

Chowringhee and Garden Reach have a perceptible advantage over Calcutta, from their openness and the great extent of evaporation from trees, which tends so much to reduce temperature.

This delightful property in trees should always be turned to our advantage; and where the grounds are kept in proper order, a moderate number of trees will prove grateful and beneficial in every sense, but especially in the hot dry season.

In the rains, the evaporation from trees can add but little to the humidity already existing in the atmosphere. It has been computed that a country covered with trees emits more vapour by one-third, than one even covered with water, and it is this property that gives to the shade of vegetables a coolness so much more effectual and agreeable than that of rocks or walls.

In the sweeping condemnation by Dr. M'Culloch, of trees, &c.—down to the very flower-pots, he goes a length unwarranted by any known facts. It is surely unphilosophical to view everything in nature as made only for man's destruction.

The following table will exhibit the state of the atmospheric temperature in Calcutta, its weight and humidity, during the years specified. The thermometer in the open air is 40° to 110° in this city.

		MAXIMUM.	MINIMUM.	
	Therm	90° 4	53° 3	Me
1830	Barom	30 131	29 512	Wet
1	Hygrom	16 9	1 5	
i	Therm	95° 8	57° 8	
1831	Barom	30 122	29 456	160
1	Hygrom	14 5	1 5	Me
i	Therm	96° 9	54° 6	2
1832 }	Barom	30 071	29 467	
1	Hygrom,	15 2	1 6	
1	Therm	97° 2	61° 1	Me
1833	Barom	30 095	29 485	
1	Hygrom	17 3	2 3	
(	Therm	99° 4	58° 9	
1834 }	Barom	30 022	29 470	
(	Hygrom	13 8	1 4	1
		In the Sun.	On the ground.	
ſ	Therm	110° 2	42° 0	
1835	Barom	30 038	29 480	
l	Hygrom	13: 9	3 2	
		May.	January	
(	Therm	97°.9	53° 5	
)	Barom	Jan. 30.087	Aug at 9 h. 50	
1836	1		29.569	1
1	Hygrom	March 14.8	August 5.6	,
	Therm	June 98.2	not absd.	
1837	Barom	Jan. 31.101	July 29.547	
	Hygrom	April 20.4	July, Aug. 42	

The mean temperature of each month by the annexed table:—

January.	February	March.	April.	May.	June.	Iv.
66°.2	69°.8	80°.0	85°.4	85°.7	83°.7	8

The greatest range of temperat January, when it amounts to about when it is about 6°. The me for the year is about sun ris 2.40 P.M.

#### INTERNAL HEAT.

Some philosophers have sought the cause of climate in the internal heat of the globe. The subject has engaged the attention of scientific men in various countries, but it is only of late that anything approaching to precise information has been obtained. From experiments recently made, it would appear certain that in proportion to the depression below the earth's surface, so does the temperature steadily increase; and even where the depths were very small the elevation was quite perceptible. At the depth of 100 feet from the surface, the temperature of the earth is the average temperature of the climate, and differs with the latitude; thus at Walso, in Lapland, it is 36°; at St. Petersburgh, 40°; in England, 52°; at Paris, 54°; at Rome, 61°; and at Cairo, 70°.—T. Griffiths.

It is stated, on the other hand, by Fourier, that at 100 feet below the surface the influence of the solar rays is extinct. The position where this takes place he calls the invariable stratum; all variations above this place are imputed to the influence of radiation, all below to the nature or primeval heat of the globe.

"It is obvious," says Professor Alfred Taylor, "that this increase of our temperature cannot be due to solar influence. 1st. Because the existence of a zone of invariable temperature shows that the power of the sun must be limited to that portion of the earth's crust which is above it. 2nd. Because, if the interior heat of the mass of the earth were due to external or solar influence alone, the thermometer should fall just in proportion as we descend below the invariable stratum. But the reverse of this is the fact, as we may presently be able to establish from the results of accurate observations. Hence we are irresistibly led to this conclusion-that in going below the invariable stratum, we are approaching some great source of heat, which must be situated within the interior of our globe." . . . " The experiments hitherto made lead to the following interesting results. Diurnal variations of temperature are not perceived beyond two or three feet. Variations depending on the months or seasons extend somewhat lower; and annual variations are entirely lost at a depth of from 60 to 100 feet, varying slightly in different localities. The maximum depth at which the changes in the thermometer are perceptible, amounts to only 1-400,000th part of the earth's diameter! Upon the alternate heating and cooling of this film, which does not exceed the nine-millionths of an inch in a globe of three feet in diameter, depend all the vicissitudes of temperature in climates, seasons, and cycles of years." . . "The extent to which the strata become warmed by the sun



elevation, we shall have a field beformany parts of our empire, and of the Tropics—in the East and West India

The importance, in a sanatory poin subject of elevation, is exemplified Egypt, yellow fever in the west, and and west, have never been known to height. The limit of yellow fever, in long ago observed by Robert Jackso the plague of Egypt confines itself t districts—never ascending so far ever and, consequently, the atmosphere ar

With the elevation of the land co progression, being caused by the enl

quires by rarefaction.

The rate of increase of cold is or yards of altitude; and this beneficent siderably increases the number of Torrid Zone. "In ascending from gine ourselves in a few days transpor Pole."

The site of Calcutta is said to be, at all, above the level of the tides a language of Asia, "what more need our position!

GENERAL AND LO

The general aspect should be aspect. The general declivity of not exclude the most opposite looks admitted as a general principle the local aspects is in the same

This principle can only be appeared for example, the entire tract of a Every one knows of what effect of a soil relatively to the sun, when the sun is elevated 45°, r whilst upon a plain the same of 45°, that is, with one quarte 45° to the north will be structed direction, which makes them

Malte-Brun, speaking of the south-south-west and so all; whilst, on the contrary, those of the north-east are the coldest. The general aspect of the valley of the Ganges is about east and west, with an inclination of the delta to the south.

#### POSITION OF MOUNTAINS.

Mountains act on climates in two ways. They attract the vapours suspended in the air; these vapours, by their condensation, produce clouds and fogs; often, also, these assemblages of watery substances, which the winds waft in every direction, are stopped in their devious course by chains of mountains, in the elevated valleys of which they continue to accumulate. These effects are still more sensibly felt, when a chain of mountains is crowned with extensive forests.

The elevated lands in Bengal-Proper hardly merit the name of mountains, and are mostly situated in the districts of Bheerboom, Sylhet, Chittagong, and along the eastern boundaries of Tipperah. They are limited in extent, and exert little or no influence on the climate of Calcutta.

#### EFFECTS OF THE NEIGHBOURHOOD OF THE SEA.

The neighbourhood of the sea moderates the excess of temperature, besides contributing, according to Pouillet, through evaporation, one of the most important sources of atmospheric electricity.

In hot climates the maritime regions are not so warm as the centre of the plains;—of this we have an annual example in the marine current of the south-west monsoon.

Calcutta is situated about 100 miles from the sea, but so level is the country that the tides ascend in the dry season as far as Sooksagur, being 140 miles from Sagor Point.

In the beginning of March, when the south-west monsoon sets in, the currents set up the Bay of Bengal, and gradually raise the sea at its head several feet, raising the river Hooghly with it, and that long before the freshes are felt. This continues till October, when the pouring of the rivers into the bay, during the rains in July, August, and September, and the changes of the monsoon to north-east in the end of October, give the current a set in the contrary direction, and gradually restore the sea and river to their condition in the previous March. The effect of the two monsoons upon the currents, and the height of the sea in the Bay of Bengal, may therefore be considered, as that of two long unequal tides

during the year—eight months of floo the point of lowest low water in the highest high water in the freshes is 20

The greatest mean rise of the tide f in the freshes is ten feet. The smallest dry season, neap tides, is four feet. To occur only in the highest, or alternate ance may with certainty be predicted and the parallax of the moon. They fluence of the south-west monsoon, an tions of the river where the peculiar direction and set of the tides in any p rise.\*

Persons who speak confidently, and on the tides and levels, assert that w irruptions of the sea, by the counterthe friction of the tides against their b in the Salt Lake to be but two fee Sagor, and consequently several feet Sagor. Of the truth of some of these mitous illustration on the 20th and 21 sea, backed by a storm from the eas overran the low countries along the Hidgelee, and up to Tumlook and D breach over the Sunderbund as far nor ing its devastations as far east as Dace irruptions of the sea within fifty years, in violence and extent. In some d 28 inches and 8 decimals, being lower for years, the thermometer averaged 84

The tide, which ran for seven hours, feet above its ordinary level, destroying ings of every kind, while the storm stood within the influence of tide and life and destruction of cattle must have

Sagor Island was covered seven feet lation of 7,000 persons few escaped month later than usual in setting in, was arrested from the saturation of thing survivors from these united calami suburbs of Calcutta, and were at last supporting the parents' lives by the win their children, the most emaciated rupee.

<sup>\*</sup> Vide the interesting paper by Mr. Jar

and four of ebb. From y season, to that of the t 10 inches.

low to high water mark an rise of the tide in the "Bores" in the Hoogly ing tides; their appearthe season of the year greatest under the ine only felt in those porof the sands, and the ular reach actuate their

er into minute details e only protected from nts of the rivers, and They state the tides ove the mean level at ood-tide, below that at rtions we had a cala-May, 1833, when the th-east and south-east from Balasore, along nd Harbour, making a Calcutta, and extend-There have been many none to equal this last he barometer stood at n it had been known

to a height of twelve bankments and builded the trees; nothing and the loss of human a enormous.

e rains were nearly a ne ordinary cultivation with salt. The starvcowded about the outer ed to the necessity of d and unnatural traffic m sold at last for one

, Trans. A. Society.

Government did much to relieve the more of these poor people; but there was much relieve. Towards the end of August there epidemic fever, which, by the end of Septe drying process, ripened into terrible violen cording to the reports of respectable natives of the remaining population to the south and did Calcutta escape; thousands of the nativin the suburbs, nearest the land inundated Europeans suffered severely, though there we ties in proportion to the numbers attacked wi

#### GEOLOGICAL NATURE OF THE

Whenever the natives of India speak of cl it by the terms "air and water." They ha definition of it.

The circumstance is here mentioned to simpression, on this important head, through least as well-founded in fact as some others entertained by philosophers.

The nature of the soil, which must of nequalities of both air and water, in an essent unnoticed in most definitions, European a geological nature of the soil would seem portant of the causes of physical climate. Of fied very early in my service in India.

Clay-lands, while they retain water and medo not readily admit either heat or air, own and adhesive nature. They are proverbially chalky soils. Soils containing much carbons matter, on the other hand, acquire a very high the solar rays, while they likewise cool mosoils, especially when dry.

Dark soils, too, absorb heat more powerful more energetically than light-coloured soils, portion of heat. In short, a very slight attervarious qualities of soils will be sufficient to affect very powerfully both the temperature the atmosphere; and hence the vast important study.

The internal nature or composition of the fluence on climate in a variety of ways. heated equally soon. One soil quickly pa heat, while another retains it for a long time.

hilsl

VEL OF SEA

300 MILES

Keunjhur on the Byturnee.

> Bonaie on the Brahmini.

Sumbhulpoor on the Mahanuddi. trees, and the climate deadly—water very scarce, impregnated with vegetable and mineral matter—iron ore in great abundance, the soil red with its oxide.

This country is very deadly; the soil chiefly red from iron oxides.

Keunjhur and Mahagirri Hills, sienite, gneiss, talcite, hornblende, kidney-iron ore and iron in every shape, it affects the needle—deadly climate except on the heights, which vary from 1,700 to 4,000—gneiss—gold found in the rivers.

Beaumurra Hills—gneiss, quartzose, iron ore in abundance—some very fine falls of water strongly impregnated with calcareous matter, leaving stalactite—considered wholesome—height 1,200 feet?

Height of this range about 300 feet above Sumbhulpoor, or 800 feet above the sea?

Gneiss formation, talcite—much iron, though less than in the other ranges—diamonds and gold worked for—matrix red clay and sand—height supposed to be about 500 feet—climate very unhealthy in the hot season—to strangers deadly.

In the "Madras Quarterly Medical Journal," for 1841, is published a very interesting report "On the Hill Fevers of the Southern Peninsula of India," by Dr. Heyne, of the Madras army. This officer, after describing the hill ranges, "which have rendered themselves known to Europeans for the malignity of the fever," and likewise those that are "as constantly free of the hill fever," goes on to state that here, the received opinions, as to the vegetable or marshy origin of fever, will not hold, for that the hills in question " are not more woody than in other healthy places:some, indeed, where the epidemic of 1808 and 1810, as well as the endemic, were most destructive, are quite naked of trees, as Dindigal, Madura, and the rocks west of Seringapatam." . . " Now, if it should be found that the fever exists constantly and invariably among certain description of hills, when others of a different composition are as constantly free from the same, would it not become reasonable to suppose that the nature or composition of the rock itself must furnish the cause of the calamity? The hills where it is found to prevail appear, at first view, to be who bestowed great attention upon the subject, attributed the fatal fever of that country, in 1844, "to exhalations from limestone rocks on which the barracks were built." Others again attributed the fever, so fatal to the 78th Highland Regiment, to neglect of the tanks; but the sickness was more generally ascribed to "an unusually high and anomalous inundation, and an equally anomalous fall of the Indus, which brought an extraordinary fertile but premature vegetation. The early and entire subsidence of the waters left this vegetation to be withered up by the sun, which produced, as it always does in Sindh, malaria."

Sir William Napier, referring to these several opinions, justly observes, that probably all the causes were here condensed. He adds, that "this and other epidemics which prevail at irregular periods in Sindh, arise from exhalations produced by volcanic action; for the country, though alluvial, is so subject to sudden and extensive changes from earthquakes, that in 1819 nearly the

whole surface of Cutch was changed."

But soils of various colours may be quite as ferruginous as the red soils—the difference being that, when dark, they retain organic matters, while, in the purely red earths, all organic substances have been, more or less, decomposed, or dissipated. May not this decomposition by the iron, together with the magnetic phenomena, elicited by heat and other agents, be productive of disease, and of fever especially, in certain climates and localities?

I am aware of the numerous objections that may be urged to this, or to any other hypothesis; but, in the absence of conclusive aids from chemistry and meteorology, we can do no better than have recourse to rational conjectures. "There is nothing so clear and obvious," says Biot, "as the discovery of yesterday:

nothing so obscure as that of to-morrow."

"Whatever may be the cause," says Hennen, "it is certain that, in many countries, the malaria does not arise until all the surfacewater has totally disappeared, and leaves the whole surface of the country, including the many courses of the winter streams, an arid desert."

The difference in the composition of the soil on the sea-coast and in the interior of the United States, is declared by Dr. Forey to cause the vast difference observable in the prevalence of inter-

mittent fever.

Where the soil is primitive, with a sandy sterile soil, the rate of intermittent fever is but 36 per 1,000 annually, while in the interior, where the soil is alluvial, composed of a rich vegetable mould from three to six feet deep, the fever is nearly six times as prevalent. In the suburbs of London it is said that in places drained within the last few years the temperature has been raised, fogs have disappeared, and influenza, marsh fevers, rheumatism

as the exhalations from the earth, the force of epidemics, &c. The varying states of the earth's magnetism cannot fail of exercising powerful influences on human health, however little may be the present amount of our knowledge respecting the nature and

operations of these influences.

In 1845 I obtained, through Sir Charles Trevelyan, samples of three kinds of earth from Sierra Leone;—viz. that of the Coast, of the Isles de Loss, and of Grand Cape Mount, as also of the Rivers Sierra Leone and Pongos. The following replies to questions submitted by me were likewise received from the colony of Sierra Leone:—

"1. What is the geological character of the soil in and around

the settlement?

- "Above high water mark, red earth—specimen sent. Below high water mark, black mud—specimen sent from Congo Creek. Flats and valleys, black earth—specimen sent from King Tom's Point.
- "2. Does ferruginous or red sand-stone prevail, or red earth?
  "Red earth. Red sand-stone in the strata (used for building)
- underneath—specimen sent from Freetown. Occasionally large blocks and some strata of blue granite (now used in building)—specimen sent from Fort Thornton.

"3. Is the soil of the most unhealthy stations of a ferruginous

nature?

" I have never heard any one station in this colony called less

healthy than another.

"4. Does the soil of the valleys or flats partake of the ferruginous character?

" Yes.

"5. Do the low lands emit any offensive gases or smells, and, if so, at what period of the twenty-four hours?

"The mud by the river banks smells when exposed to the sun

at ebb tide.

"6. Does magnetic iron exist, and, if so, is the variation of the compass affected by it?

"Yes-specimen sent from Tower Hill Barrack Yard.

"7. Has the nature of the soil in general, or of the most unhealthy localities in particular, been chemically analysed?

"Answer by Dr. Fergusson—So far as I know there has not been any minute analysis of the soil of this colony beyond the general one, that the soil is ferruginous; gravelly in some parts; loamy in others; in others sandy; but in all ferruginous."

For the following report, on the qualities of the specimens of earth from Sierra Leone, I am indebted to Professor Bowman, of King's College, London. The numbers, as given below, do not Parisot remarks justly, that animalised vegetable matters in a state of decomposition are condensed and not decomposed by porous bodies, amongst which fine sand may be classed. "In the mild and moist winter putrid decomposition commences. The subtile ferments it produces are condensed in the sandy soil of which Egypt is formed, and in which the dead repose. When the elevation of the temperature makes the ferments evaporate, man receives them by the superficial absorbent vessels, or by those of

the lungs in the act of respiration."

Let us contemplate a desert country, the rivers, abandoned to themselves, become choked and overflow, and their waters serve only to form pestilential marshes. A labyrinth of thickets and of brambles overspread the most fertile hills. In the meadows the unsightly wild mushroom, and the useless moss choke the nutritious herbs; forests become impenetrable to the rays of the sun; no wind disperses the putrid exhalations of the trees which have fallen under the pressure of age; the soil, excluded from genial and purifying warmth of the air, exhales nothing but poison; and an atmosphere of death gathers over all the country. But what do not industry and perseverance accomplish? The marshes are drained; the rivers flow in their disencumbered channels; the axe and the fire clear away the forests; the earth furrowed by the plough is opened to the rays of the sun, and the influence of the wind; the air, the soil, and the waters acquire by degrees a character of salubrity; and vanquished nature yields its empire to man, who thus creates a country for himself.

Agriculture must be much improved in Bengal before the European, in the language of Malte-Brun, can be said to have created a country for himself. A Hindu field is described by Mill to be in the highest state of cultivation, where only so far changed by the plough, as to afford a scanty supply of mould for covering the seed, while the useless and hurtful vegetation is so far from being eradicated, that where burning precedes not, the grasses and steriles which have bid defiance to the plough, cover a large portion of the surface. The same author concludes, that "everything which savours of ingenuity, even the most natural results of common observation and good sense, are foreign to the agriculture of the Hindus. Their ideas of improvement are very limited; they scarcely extend beyond the introduction of irrigation into land which was formerly cultivated dry. Each small proprietor is content to follow the customs of his forefathers; the same rude implements of husbandry, the same inferior race of cattle, and the same practices are still in operation, which have existed unchanged for centuries. As to any new experiments of general measuring, draining, differences in the rotation of crops, introducing new grain or vegetables, or new sorts of those already

known, any attention to their better and more combined syst people could raise the same or all these are out of the question

I cannot find that the examp much influence on the state of of India. It is certain that is supplied with excellent vegetal I believe matters are much the nock, the founder of Calcutt In the Appendix to the Parlia the population of the 24-Per at 1,225,000, which I have rea

THE PREDO

The united influence of all sical climate is variously mod all their variations depend on the heat of one climate and the tinual influence on each other, accumulation in it of the not emanating from the earth's sur

Mr. Atkinson, referring to the winds on plants and insects—observes, that "there is some change of wind, causes a sin stances and structures. The plass,' and the chickweed," he barometers, opening their deliclosing them against rain. So north and east winds, which 'shelter themselves in holes.'

The northern parts of a growth their cold air towards to they will receive warm air changes some time after the course that hemisphere in which this luminary on the atmosphe causes, the cold air from the course for half the year, and to the other.

The south-west rainy mon periodical winds, begins on the Delhi by the end of June, ext Affghanistan, but greatly modified. It prevails more in the mountains than the flats of the Punjaub; the hills and valleys of Cashmere have their share of it, and it gradually loses itself westward in the valley of Peshawur, where it appears only in clouds and showers. On the Coromondel Coast it is retarded, the clouds brought by the south-west winds being detained by the Ghauts. It reaches Bengal by the 15th of June.

Owing to the arrest of the south-west monsoon by the mountains, and consequent accumulation of vapour, an extraordinary deposition of rain takes place on the Malabar Coast, being not less than 123-5 inches in the year in the latitude of 11½ north.

When not influenced by elevated lands, this monsoon generally prevails north of the Equator from April to October, accompanied by tempests, storms, and rain, while a north-east wind blows during the other six months. The periodical winds that prevail in the Bay of Bengal extend their influence over the flat country, until they are diverted by chains of mountains into another direction, nearly correspondent, however, with the course of the Ganges. When the sun has passed into the southern hemisphere, the monsoon alters its direction; the mass of air which had been accumulated during the hot season and rains on the central platform of Asia, now bestirs itself, and moves towards the regions south of the Equator, where the atmosphere has been dilated and dissipated by the solar heat. Over most parts of the Indian Ocean this monsoon proceeds from the north-east, because the central platform lies to the north-east. On the other hand, as the seas of China, of Borneo, of New Guinea, of Java, have the centre of Asia to the north and north-west, the monsoon comes to them from these points.

In the south of Bengal the prevalent winds are north and south; in Behar east and west, and the same takes place in Assam, follow-

ing the course of the Berhampooter.

That the monsoons exercise a beneficial influence on health cannot be doubted, but especially the south-west, from its prevalence during the greatest heats, and from its greater power of thoroughly ventilating the country—"Qualis aer, talis spiritus." Stagnation would prove immediately destructive to health in a climate where there are so many various and abundant sources of noxious effluvia, to ripen into activity by such a cessation of wind as should admit of their accumulation, and that of heat, in any one place, or in such streets, for instance, as those of the native portion of Calcutta. Of the north-east monsoon I must limit my praises solely to its ventilating properties; for in every other respect it exercises an unfavourable influence on health to a degree not generally known; indeed, so far from it, that it is common to hear the accession of the cold season hailed by in-

valids who are ignorant of the many dangers it carries along. It is the true "Sirocco of the North."

The following table gives the direction of the winds at noon during the years specified:—

			N.	NW.	w.	s.w.	S.	S.E.	E.	N.E.	Calm			
			Days											
In 1832	he wir	nd was	44	60	26	44	67	49	4	71	9	No	regist.	1 day.
1833	**	12	56	39	14	30	115	24	25	32	29	,,	1)	2 days
1834	**	"	53	52	12	25	99	22	26	40	36	-		
1835	"	- 11	41	65	31	84	41	45	12	31	7	29	99	8 days
1836	25	- 311	67	58	88	38	46	7	23	20	12	"	- 19	6 day
1837	**	25	83	26	77	43	76	6	21	13	7	**	at	3 da
1838	39	**										-		7
T	otal .		344	300	248	264	444	153	111	207	100			7

The following table,\* given by Mr. Smeaton (Philos. T and confirmed by Mr. Hutton (Mathematical Dictionar hibits in pounds avoirdupois the pressure which differe will exert upon a square foot of surface exposed direct) them. The first column is a rough representation second:—

Velocity	of Winds.	Force on one Square Foot	Chamatana		
Miles per Hour.	Feet per Second.	in pounds Avoirdupois.	Character o		
1					
2	1.47	•005	Hardly '		
2 3 4 5	2.93	-020	Just pe		
4	4:40	*044			
	5.87	.079	Gent!		
10	7.33	123			
15	14.67	492	Ple		
20	22.00	1-107			
25	29.34	1.968	V		
30	36.67	3.075			
35	44·01 51·34	4.429	7		
40	20.00	6.027			
45 50	58.68 66.01	7·873 { 9·963 }			
55	73:35	12.300			
60	88.02	17:1715			
80	117.36	31.490			
100	146.70	49.200			

\* A. B. Maddock, M.D., on the Influ

THE RAINS-SOURCES OF AQUEOUS EXHALATIONS-HUMIDITY.

The endiometric processes having failed of discovering in the atmosphere of places the most opposite, such as the narrow lanes of London, and the summits of lofty mountains, any difference in the constituent properties of their permanent gases, it becomes a question of the highest interest and importance to ascertain the varying quantity of aqueous vapour in our atmosphere; for it is the only ascertainable fluctuating ingredient in its composition. To the medical topographer, a minute inquiry of the kind here suggested may lead to a knowledge interesting and important respecting the causes and cure of certain diseases, in so far as these may be connected with climate, and this is no small benefit from the proper application of instruments.

"The rain in falling," says Mulder, "carries with it everything that floats in the atmosphere, and which is not essential to its constitution; which brings back to the earth what came from the earth; and which, while it thus purifies the atmosphere from these hurtful adulterations, restores to the soil these numberless volatile substances, where they have abundant opportunities of forming

not only harmless but useful combinations."

Without taking into view the expanse of the Bay, the coup-d'wil of a good map of Bengal will at once show how bountiful nature has been to that country, by means of her majestic rivers with their innumerable tributaries, in yielding the sources of aqueous exhalations, and it were gratifying to the medical topographer, could his description be limited to these. There are not any lakes in Bengal resembling those of Scotland or Canada, but there is a profusion of extensive jheels, which may be either denominated shallow lakes or morasses. A large proportion of these in the dry season contain little or no water, but during the rains present immense sheets, over which boats of the greatest magnitude may be navigated, and some are navigable to a certain extent throughout the year. There is reason to believe that nearly all these stagnant sheets of water rest in what were, at a remote period, the channels of large rivers, which have since altered their courses and now flow in other directions. The area of Bengal and Behar is 149,217 square miles, and with Benares not less than 162,000 square miles. The following proportions of the surface are grounded upon many surveys after making allowance for large rivers :-

Rivers and lakes (one-eig Deemed irreclaimable and Sites of towns and villa	barr	en (one-six	th)	cc. (one to	wenty-	Parts.
fourth)	urths	**	**		**	1 3 9
Waste (one sixth)				2.		4
			Total		**	24*

According to another calculation, Bengal contains 97,244 square miles; if from this, that portion of Tipperah which is independent, the tract of the Sunderbunds and other wastes, equal to 13,244 miles, be deducted, the remaining inhabited country will be equal to 84,000 square miles; but the extent of waste and surface, occupied by rivers, marshes, &c.. seems here greatly under-rated.

When all this is considered, along with the complete saturation during five months in the year, of every inch of soil, even that which may not be actually inundated, the extent and sources of aqueous exhalation—the commerce of land and water—may be imagined. It is ascertained that the capacity of the atmosphere for moisture varies with its temperature, so that at 113° it holds a twentieth part of its weight of moisture; at 80° a fortieth, and so on.

To a scientific officer well acquainted with the localities, I put the two following questions: 1st. Taking the area of the 24 pergunnahs to be 882 square miles, what proportion should you say the water surface bears to the land, on the 30th May and the 10th of October, the first being just before the rains, and the latter just after? 2nd. What proportion does the cultivated land bear to the waste and jungle within the said area?

Answer to 1st Question.—I should say that on the 10th of June, you might assume one-twentieth part as the proportion of the water to the land; of course, I mean by water, ground from which exhalations could arise. Answer to Question 2nd.—The cultivated land may be about 14-20ths; water, 1-20th; roads and villages, 2-20th; uncultivated, 3-20ths.

The subjoined table will show the annual fall of rain during the years specified. The average annual fall is about 60 inches.

			RAIN.			
1830						inches 63.28
1831				**		57.50
1832	**		**		**	49.26
1833	2.0	**	**		**	60.56
1834		**	**	**	44	68.73

\* W. Hamilton.

1835*						inches. 85.50
1836	**	**				45.39
1837						43.06
1838		**	**		· ce	44 16
				Mean		59.228

The following are the average rates of evaporation for the dry months: January, three inches; February, five; March, seven;

April and May, nine.

It is during the periods of the year when the drying process is in greatest activity that unhealthiness prevails with greatest severity, namely, the commencement and termination of the rainy season. In the former, or that called by the natives the lesser rain (Chota bursat), common remittent fevers arise, and at the termination, or that from 15th September to the end of October, the severest forms of the same fever prevail, but chiefly amongst the indigent natives, and Europeans newly arrived.

The connexion of the rainy season with disease, would, in such a climate as this, form a highly interesting branch of inquiry; but the absence of everything like statistical object in our hospital reports, prevents even an approach to accuracy on this, or indeed

on any subject connected with climatorial influence.

The following statement is taken from the "Bengal Hurkaru"

of September 6th, 1845.

"The Englishman is informed that seventy-four inches of rain fell in Arakan during the last month; and that fifty inches had fallen during the first twelve days of the present month. The whole fall during the monsoon, to the 15th August, was two hundred inches."

The quantity of rain that occasionally falls in a brief space of time within the tropics, and especially near the equator, is enormous. On the 10th May, 1835, sixteen inches fell in Calcutta, in twelve hours. But, owing perhaps to the direction and elevation of the mountain ranges, and to the nearer approach to the equator, it is in the Burmese territories that we find the most surprising examples of great falls of rain. Those who served at Rangoon, and in Upper Ava, as the first European observers, will never forget the rainy seasons of those countries.

THE HOOGHLY RIVER—THE WESTERN BRANCH OF THE GANGES— THE BHAGIRATHI, OR TRUE GANGES OF THE HINDUS.

By some persons who have spoken of our climate, a large portion of the evils under which we suffer, has been ascribed to the river—its supposed overflowings—its sluggish tides and foul waters—its muddy and slimy banks, and the action of a vertical

<sup>\*</sup> Sixteen inches fell on the 10th May, in less than twelve hours.

sun upon them, &c., but I shall view the river in a different and

more friendly light, as the purifier of our city.

Certain am I that without this great scavenger, to whose tides we owe more than Captain Hamilton ever dreamt of, we should now be in a worse condition even than when he left us an hundred years ago. The truth is that, under moderate supervision on the part of the police, the river banks are inoffensive; and along their whole extent, although crowded with buildings for a space of nine miles, disease will be found less prevalent by far, than in the interior quarters towards the east; in short, the causes of fever are to be traced to other and more palpable sources than the riverbank, which is the most elevated of all our grounds, being from three to four feet above the surrounding levels. The causes of w healthiness in Garden Reach, after the salt water inundation 1833, could be readily traced to the state of the back-groundsone ever thought of looking for them in the river-bank. The avrise of the Ganges and its branches, is in

						Inches.
May	7.5	**	**	**	6	0
June	12		**.		9	6
July		**			12	6
Half of A	lugust	***	**		4	0
			Total		32	0

From above 350 observations of temperature made Prinsep, of which the details are given in the JAsiatic Society, it would appear that "the mean the surface water exceeds 81° Fahr., everywhere b and the sea." In the dry season the mean rate of than three miles per hour; in the rainy season, a undations are draining off, the current runs from seven eight miles in particular situations.

The river is at its lowest in the beginning of freshes are at their height in September, v scarcely visible off Calcutta; and the river sweet, far beyond Saugor in the open sea."

The following table exhibits the depths of the different seasons.

Minimum depth

Maximum average depth of eleven of the shallow
the Hooghly spring ebb

Maximum average depth over the same places sp

Minimum depth
Minimum average depth of eleven of the shalthe Hooghly spring ebb
Maximum average depth over the same place
Highest rise, ditto, ditto
Difference between higher

The situation and extent of the salt-water lake have already been described; and it only now remains to state the probable advantages of causing its area to be drained, filled up, and brought under cultivation. Owing to the shallowness of the lake generally, it is easy to drain it for the purposes of native agriculture; but to deprive the ground so drained of the sources of noxious exhala-

tion is not so easy.

It is not sufficient to convert the ground into a state of soft low meadow land; for the most dangerous exhalations are those which are retained, and occasionally emitted from under a crust of earth during the drying process, whereby they would appear to acquire unusual concentration, and prove the origin of the worst fevers. It is necessary that the grounds be thoroughly drained, leaving none of the characters of the marsh, otherwise it had better be left as it is; its present condition being one of far greater safety than such half drained soil as that obtained from the marsh of Chartreuse, for instance, near Bordeaux, which caused in the year 1805 alone, 12,000 persons to be affected with fever, within the

city, of whom 3,000 died within five months!

Two modes of effecting the drainage suggest themselves; the one by letting in the river during the rains, and thereby gaining a succession of deposits of the river silt, so as gradually to fill the lake, and thereby bring it in time to a level with the surrounding land; this would seem the easiest; it imitates the simple operations of nature, and would be the cheapest; but perhaps not the most conducive to health. Another mode is by a deep and wellconstructed canal, so as to effect the drainage; but, as even this must, to a certain degree, prove a receptacle for noxious matter, and offer a considerable surface for evaporation, a close line of umbrageous trees should, under the proposed plan, be planted along each side of the canal, as being powerfully attractive of marsh exhalation. This property in trees was practically known to the ancients,\* and is now beneficially exemplified in Demerara, and other parts of Guiana, "where the humid heat constantly cherishes the seeds of disease."

The ground cleared from water, should be well ploughed and cultivated, the ploughing to be done during the heaviest rain, so as to prevent exhalation; for it is during a certain stage of the drying process, that marsh exhalation is most concentrated, and it has been observed in many countries that the drying up of brackish water is more injurious than that of either salt or fresh alone. A succession of crops purifies and evaporates the soil, and thereby obviates exhalation; but they should not be of rice, or such crops

<sup>\*</sup> Regaud de L'Isle says of the malaria of Italy, that various obstacles form barriers which they cannot pass, and against which they deposit themselves.

as require profuse irrigation. The want of attention to some of the precautionary measures above hinted at, has neutralized the advantages that would otherwise have resulted from extensive draining executed in some parts of France and Italy; and I have only thus long dwelt on that of the Salt-Water-Lake, because I believe its proper performance to be a matter of great importance to this city, so far as regards the prevention of disease; and I need not here insist on the superior efficacy of preventive measures, such as have advanced in our own country apace with our civilization, and altogether banished from us some of the severest calamities that have ever afflicted the human race.

#### THE WOODS AND MARSHES OF THE SUNDERBUNDS,

The enormous extent of the Sunderbunds-covering a superfic of more than 20,000 square miles, and extending 180 miles to South and East of Calcutta—is composed of marshy land, cover with forest and underwood, together with the numerous embe chures of the Ganges-a region from whence the solar light heat, and the air, are excluded by the forest trees, to the ey tion of all minor vegetation. When we consider all thes cumstances, and their common influences on all the condition the atmosphere necessary to health, we shall be at no loss ! ceive how great must be the deteriorating influences of s region on the climate of Calcutta. That "forest on the bo the sea," must necessarily exercise a very powerful influ the temperature, humidity, electricity, and freedom of cir of the atmosphere in and around it, not to speak of the exhalations generated within, and disseminated without evident that the electric condition of the air must be affected by the combination of oxygen with the material plants over so vast a surface, as also by the extent of tion from it; and they are subjects deserving our most ca sideration.

How far the geological nature of the soil—the amouration—the presence of forests—the vicinity of seas rivers; how, in fact, all the local circumstances, noted graphic charts, may affect the electrical states of the which "we live, move, and have our being;" these a of the last importance to be observed and noted.

That the clearing of the extensive surface of the S or of any considerable portion of it, leaving belts a forest trees, would tend greatly to improve the local c around Calcutta, there can be no doubt. The histo and that of the effects of clearing marshes and forests in the neighbourhood of other cities in other countries, offers a demonstration of this fact.

Such a measure would open out the city to the more free influence of the sea-breezes, diminish the moisture of its atmo-

sphere, and purify it. These are no speculative results.

It is found, contrary to the vulgar opinion, that in cleared and cultivated tracts, the air is rendered drier and warmer in summer, and colder in winter, than in such as, from want of cultivation, remain, like the Sunderbunds, covered with wood and marsh.

"Earth-clouds" are commonly observed in the early morning to the east of Calcutta, over the borders of the Salt-Water Lake, and the cleared portions of the Sunderbunds. They are caused by the immense radiation of heat in the calm nights, and by the cooling of the earth's surface far below that of the air above—thus forming vast clouds or mists, close to the earth's surface. If then, to counterbalance the only disadvantage attendant on clearingsome little increase of temperature-we obtain purity and dryness of our atmosphere, we shall still be very greatly the gainers; for it is not so much from the high rate of temperature we Europeans suffer, as from the excessive humidity that is conjoined to it for so many months in the year, and both which, commingled with the terrestrial exhalations, tend gradually through their united influence, by inducing what may be termed a Cachexia Loci, to undermine the best and most robust of constitutions.

The clearing and draining immediately around the city, partial and imperfect as these are, have yet removed only some of the concentrated evil; but that emanation which was death within a few yards, cannot be other than insalubrious even at the distance of

a few miles.

## POSTSCRIPT.

Referring to the influences of all the physical and geographical circumstances detailed in the foregoing notes as affecting the electricity of the atmosphere, I would call attention to the conclusions arrived at by Dr. Pallas of the French army in Algeria, offering them here only as subject-matter for examination and test in other hot climates. Dr. Pallas considers:-

1st. That, just as light and air are the essential agents of vision and respiration, so electricity is the functional agent of innerva-

2nd. That the greater number of diseases, and especially those which belong to the class of neuroses, are occasioned by the exaggerated influence of general electricity, of which clouds, storms,

and marshy regions are the most fruitful sources.

3. Marshes, in their geographical constitution, and in the effects which they produce upon the economy, present the greatest analogy to the galvanic pile. Thus their action is much the more baneful as they contain certain proportions of water, and their activity is considerably increased when the water contains organic or saline matters in a state of solution. This explains why salt marshes and such as are near maritime rivers are the most insalubrious. The drying up or submersion of marshes produces analogous conditions to those of a galvanic pile deprived of humidity, or which is under water, and the effects of which are there insignificant.

4. The researches of philosophers and physiologists have shown that the electricity produced by our machines exerts a special action upon the nervous system. Experience and rigorous observation of facts proves that the diseases which are produced by a marshy atmosphere are primarily nervous, and become inflammatory only by the re-action of the nerves upon the vascular system,

inducing consecutive, local, or general irritation.

5. The neuroses are occasioned generally by the effects of eletricity, and intermittent fevers have a similar origin; that is to see they are due to the electrical emanations of the marshy pile, we are very active in hot countries, and not to miasmata, which

never been met with.

Professor Schönbein considers, on the other hand, the physiological importance of electricity has, upon the whole much exaggerated, that agent, in comparison to heat an acting but an inferior part in the economy of organized "Electricity," he says, "would affect neither the sense o smell, if atmospheric air did not contain oxygen and and the phenomena of sound and light, perceived during discharges, are due to the vibrations into which the part are thrown by the electrical discharges, electricity have nothing to do with them."

# MEDICAL CLIMATE AND THE INFLUENCE OF SEASON.

"Si l'histoire naturelle a besoin d'une bonne géographie physique, la science de l'homme a besoin d'une bonne géographie médicale."—Cabanis.

Under this head I shall offer some cursory observations on what is usually called by physicians "medical climate," and also some brief notices of the geography of disease, as connected with our

Bengal climate and seasons.

Temperature and humidity being the elements that give activity to terrestrial emanations, and all the external causes affecting health, derived from locality, I shall now consider them apart from the subjects above classed; remarking generally that the heat of Hindustan as compared to that of other regions, is more characterized by its duration than by its intensity. On the other hand, where the cold is excessive, the rise of temperature in spring and summer is sudden and intense, while again in those countries where the range is more limited, the seasons glide gradually and

imperceptibly into each other.

Speaking of the average of men, the terms hot, warm, cool, cold, as applied to the surrounding air, are regulated by the sensations produced; and if the heat be carried off as fast as it is generated, and no faster, no particular sensation is felt—the bodily powers being neither stimulated nor exhausted. Supposing then that no extraordinary exertions are made, the equilibrium is maintained when the thermometer stands at 62°, or thereabouts; and this point in the scale is therefore called temperate. All degrees above that point, up to 70°, are reckoned warm; all above 70°, hot. Descending again in the scale, we speak of the temperature denoted by any degree between the 60th and the 50th as being cool, and any lower degree of temperature is cold. Changes of temperature seem to be as readily felt at one part of the thermometric scale as at the other, and in whichever direction they take place. Captain Parry found that a rise in the thermometer from 13° below zero

23° caused discomfort to his men. "I may possibly incur the charge of affectation in stating that this temperature was much too high to be agreeable to us; but it was nevertheless a fact that everybody felt and complained of the change." On the other hand, the fall to from 40° or 50° at night in Bengal, during the cold season, might seem to an European as both agreeable and salutary; but a very brief experience teaches him the contrary. Much has been said and written on the superior capabilities of adaptation to climate in man over the lower animals; but if the power and just application of the arts of civilization be deducted, I am disposed to think with Dr. James Johnson, that the difference would be but small; for, even with these aids we find that in this climate "many die suddenly, others droop, and all degenerate," very much as with the lower animals of more temperate regions; and all we car hope to learn is how best to conquer by obeying nature. thus that man, the weakest of animals, is, in reality, the strong and it is under circumstances apparently calculated to overy and destroy his vigour that he finds the means of developir faculties and resources, which excite even his own astonis'

Gibbon, after stating that the Roman soldiers, from the cellent discipline," maintained "health and vigour in all adds, that "man is the only animal which can live and mevery country from the equator to the poles. The hog approach the nearest to our species in that privilege." as stated by the historian, that men do "live" in othe natural climates, but their existence is very unlike and vigour of the Roman soldier, whose habit and not allow his manly character and physical energy

in indolence.

"The truth is," says Dr. Johnson, "the tender incapable of sustaining the degree of exposure to the causes and effects incident to, or arising from vicis which so speedily operate a change on the struct exterior of unprotected animals. The object of that first sight might seem irrelevant, will now sevident that nature does not operate more pacting the ill effects of climate on man, that follows that we should not implicitly confithe spontaneous efforts of the constitution call into its aid those artificial means of pation, which reason may dictate and experit that we should study well the climate, and frames to the nature of the skies under v

Although the physical effects of cliencing the differences by which the var species are characterized, are foreign one cannot help remarking, that, if the native of Bengal Proper is to be classed among the Caucasians—the standard of the human race—the effects of climate and locality must indeed be great and remarkable.

No climates exist that are uniformly hot and dry, hot and moist, cold and dry, or cold and moist; yet certain countries have such a preponderance of one or other of these qualities, as to give a very marked character to the physical and moral nature of man; and physicians would do well to observe these results of climate more closely than has yet been done:—"Si l'histoire naturelle a besoin d'une bonne géographie physique, la science de l'homme a besoin d'une bonne géographie médicale."

The moral as well as the physical influences of climate have been considered so powerful by some philosophers, as to make some persons doubt whether a people, situated as our Asiatic subjects, are capable of receiving the impress of European knowledge and institutions. There is in hot climates, it has been well observed, a vis inertiæ which indisposes men to change their cus-

toms, or to cope with abuses; and the indolence which the climate occasions conduces to the stability of their barbarous institutions.

"The astonishing rapidity of political revolutions in Asia," says Montesquieu, "arises, however, out of one fact which is really dependent on its physical geography. In that part of the world, weak nations are opposed to strong; people warlike, brave, and active, border upon those who are effeminate, idle, and timid; the one must necessarily be conquerors, the others conquered. Here we have the principal reason of the liberty of Europe and the slavery of Asia." "It is necessary," says Malte-Brun, "to combine this just remark with another truth proved by physical geography, namely, that Asia has no temperate zone, no intermediate region between very cold and very hot climates. The slaves inhabit the hot, and the conquerors the elevated and cold regions."

In European countries, a certain amount of injury is caused to public health by the agitations of ceaseless competition, commercial speculation, religious controversy, and party politics; and though these influences are not apparent in India generally, we cannot altogether overlook them, or the many causes arising from long ages of civil and sacerdotal dominations, and other abuses of government which, with difference of race and of religion, produce so remarkable a difference between the moral and physical constitution of the Asiatic and European, so as to render the latter so much more liable to be affected by tropical climates.

The hot and dry season in Bengal extends from the beginning of March to the middle of June, during which the winds are steady and strong from south and south-west. The temperature rises gradually from 80° to about 90°—95° in the shade, and reache

to 100°—120° and upwards in the open air. Notwithstanding the high temperature, this season is rendered far less oppressive to the feelings than might be supposed, by means of the moisture carried along with the monsoon in its passage over the bay, and likewise by the frequency of refreshing storms, accompanied by rain lightning, and thunder. Of Calcutta at mid-day, in April, May, and part June, it may be said, however, with truth, that it is "a city o stone, in a land of iron, with a sky of brass." The local news papers of May, 1851, speak of the heat as more intense than it had been for years. "The thermometer in the coolest rooms stand at 92° to 94°, and the breeze which should bring refreshment at the close of the sultry day, has been as the breath of a furnace."

The most ordinary and simple effects of the season just de scribed are—determination of the fluids generally to the surface the body, the blood being venalized in proportion to the elevation of temperature; and respiration being less perfectly carried o owing to the rarefaction of the air, and the consequent diminuties of its oxygen in a given bulk, a vicarious decarbonization of blood is established in the great increase of the biliary secretio while, at the same time, the urine is surcharged with saline pregnation, and much diminished in quantity. Recent ex ments render it probable that, as temperature rises, the quant oxygen consumed in respiration is less, while the amount of bonic acid exhaled is less also. At this season too, it is be that both the relative and absolute quantity of carbonic ac' haled is diminished. There seems at length some reason that the vicarious nature of the relations existing between purative functions of the liver, kidneys, mucous surfaces, lungs may receive demonstration from chemistry.

"Vierordt has made a very extensive series of ob with the view of ascertaining the connection between the rure and the pulse, the respiratory movements, and the air expired, and of carbonic acid, in one minute. I ments were made at every degree of temperature between

76° Fahr.

"From a table of observations, we see that elevation rature is accompanied by a diminution of the number depth of the inspirations; its effect on the excretion acid is, to a certain degree, of an indirect nature, sin inshed number and depth of the expirations must siderable influence; but inasmuch as the diminumerely show itself in the total quantity excreted, I per centage of carbonic acid in the expired air, it that the elevation of temperature has a more direct the excretion of carbonic acid than could be accounted in the expiratory action.

"The degree of atmospheric moisture, to a certain degree, influences the respiratory functions and the excretion of carbonic acid. Lekmann, some years ago, experimented on this subject with pigeons, greenfinches, and rabbits. The quantity of carbonic acid exhaled in a moist air was much greater than in a drier atmosphere." . . . . "The researches of Vierordt on the influence of atmospheric pressure, show that this also is by no means an unimportant element." . . . "Vierordt and Baral agree in the opinion that more carbonic acid is excreted in winter than in the summer.

"The excretion of carbonic acid is very considerably diminished during sleep. This is most decisively proved by the experiments of Scharling, who found that a man, during one hour of the night, exhaled only 22.77 grammes, who, during an hour of the next day, immediately after a meal, exhaled 33.69 grammes; and who, in another case, found that the horary excretions of carbonic acid during this night and during the day were as 31.39: 40.74."—British and Foreign M. C. Review, July, 1854. The alternations here mentioned cannot fail of being important in relation to European health within the tropics.

It appears to me that the physiological action of heat is exhibited, first, in its temporary stimulation of both the ganglionic and cerebral systems of nerves, and in the increased manifestation of all those functions governed by them; and secondly, in its sedative and more lasting influence on the same systems of nerves, and in diminished manifestation, consequently, of all those functions governed by them. This last result is strikingly exemplified in the diminished muscular powers of both the heart and uterus.

Bearing in mind these important and interesting facts, as they appear to me, we shall at once perceive how it is, that for a certain time, the respiration, circulation, the hepatic function, and that of innervation itself, are unnaturally excited, and how the newly arrived European is exhilarated in mind, and so prone also to acute inflammatory and congestive abdominal diseases. But, as this state of over-excited or exalted function cannot be maintained beyond a certain limited term of years, without danger to the power and equable functional action of the source—the ganglionic nerves-we see how, along with a feeble respiration and circulation, diminished power of generating heat, and torpor of the hepatic function eventually succeed ;-all tending to congestion and other dangerous forms of more passive disease. Again, we perceive how the long-continued operation of alternate stimulation, and of subsequent depressive and sedative influences on the functions of the organic and animal systems of nerves, produces and confirms that eventual bodily and mental inertness and indisposition to healthy action, which not even the most powerful of Europeans can he always to resist, beyond certain defined limits as to time.

The hot season swells the exterior and produces that general chubbiness of appearance which is so remarkable in the torrid zone, even where the weight of the body is sensibly diminished: it increases the animal heat, and accelerates the pulse, accompanied by a prodigious increase of the pulmonary and cutaneous transudations: it produces nervous excitability varying in degree in most persons—an exaltation of the general sensibility: eruptive diseases, latent during the cold season, become actively developed; and the cutaneous vessels, even in healthy persons, are excited to the extent of producing the distressing eczema solare, known in the tropics by the name of prickly-heat. In ordinary seasons, we find that here, as in the West Indies, the most healthy months are from February to May, inclusive; while in the seasons of epidemics (cholera in particular) these are the months in which the disease is most fatal and long continued; indeed, it seldom vanishes till the setting in of the rains.

Under exposure, excessive fatigue, mental depression, or neglect of temperance in diet, results ardent fever, with some serious loc determination, and that very frequently to the cerebral organs—casionally to the liver; but though this is admitted, under the r sures of precaution dictated by common sense and experience very hottest are yet the healthiest of our seasons, and of our sta also,\* which goes far to prove that it is not heat alone that do the mischief, but something else in the climate and in the tution and habit of the stranger European, not common native of the country. From the results of personal obtain active field service both in India and in Ava, I am lee clude that mere heat, unless long continued, and combintemperance, with chills, and other untoward circum very rarely the direct cause of disease.

Édwards states it as a fact that the quantity of cuts lation from the human body is sometimes ten times than in moist air, and that it is doubled in the mer 32° to 64° F.

I have seen men of the Bengal Pilot service, pound laboured in the sun, during their youth and must the functions of the skin appeared at length to hausted. They exhibited a permanent drynes' scaliness of the whole surface of the body, from rienced distress in the cold season of Bengal English winter and spring. One old pilot furaceous that he would often say: "I could

<sup>\*</sup> At Agra, the hottest of our stations, the percentage of one in fifty per annum, out of a garrison of one thousand than shown in any table hitherto prepared in India.—As Henderson.

my nail on any part of my body and limbs." With these men the functions of the kidneys were always disturbed also.

The troops from Bengal and Madras were exposed, in the expedition to Egypt, to an excessive heat in crossing the desert from Kosseir to the Nile; yet they enjoyed excellent health, because they were not exposed to excesses, and their minds as well as

their bodies were kept in activity.

The superior power of enduring heat, under a sudden effort, or while the mind is keenly occupied, possessed by Europeans in India, even beyond the natives, has often been exhibited in other climates. It would seem to arise from the same causes that enabled the natives of the southern countries of Europe, according to Larrey, to bear the cold of the Russian winter during the French retreat from Moscow, "better than the natives of the Northern and moister climates—such as the Hanoverians, the Dutch, the Prussians, and the other German people: the Russians themselves, from what I learned at Wilna, suffered more from the cold than the French." They also suffered in Holland, in 1799, much more from cold than the British with whom they co-operated, as observed on the spot by Robert Jackson.

The inhabitants of the temperate regions of the globe, it is supposed, having their constitutions matured by genial climates, are able to bear the extremes of both heat and cold better than those whose frames have been weakened either by the severities of the

arctic, or the relaxations of tropical climates.

These remarks must be understood to apply only to sudden efforts or brief exposures; for it is now well known that to the duration of heat more than to its intensity—to a long continued exposure to a high range of temperature, more than to the temperature itself—must be referred much of the injurious influence of tropical climate on European constitutions. Robert Jackson, speaking of the first American campaigns, says, "Bad effects from the greatest exertions in the hottest weather of summer were extremely rare in that country, after the campaign had been continued for a few days."

The equable determination to the surface consequent on the progressive increase of temperature, seems to exercise an agreeable as well as a favourable influence on general health, especially in the old Indian; and even a new-comer seems to bear without complaint or apparent injury the great augmentation of the sensible perspiration. It is only the opposite condition—the total suppression of it in our cold season—that is felt by all so unnatural and unhealthy:—indeed, it is then we become aware of the effect of long continued exposure to a high range of temperature, through our extreme predisposition to be injuriously influenced by cold.

Miscarriages, frequent at all seasons in India, occur yet more

serious complication. It renders the cure difficult and tedious, the system being cachectic, the blood dissolved, there being a

general anæmia, in short.

To give a more concentrated view of the physical circumstances above related, it may be said that in hot climates, the air being expanded, less oxygen is taken in at each inspiration. The necessity for hydrocarbonaceous food is therefore lessened. Less of the "elements of respiration" ought to be taken in the food than would be taken in colder climates. In consequence of the internal increase of temperature, less internal heat is required.

Exercise increases the heat of the body by increasing the rate of circulation and respiration. In a very hot climate all increase of heat is undesirable. Moreover, the excessive heat renders muscular action impossible, because the circulation is chiefly directed, in consequence of the activity of the skin, to the surface, in order that fluid may be furnished for evaporation, to keep down the heat of the body, to prevent the parching of the surface which other-

wise must ensue.

In consequence of the lessened muscular action, less of the albuminous constituents of the food are required to supply the waste of the muscles.

Hence, in hot climates, less of both kinds of food should be taken, and nature points this out in the absence of appetite; to force an appetite stimulants are taken, and then the system is overloaded with nourishment. The excessive perspiration requires an excess of liquids; but instead of water alone, sugar and spirit, the elements of respiration, are taken with the water as beer, and the

spirit, by its stimulating properties, is doubly injurious.

The excessive flow of blood to the surface (the consequence of the high temperature) no doubt prevents, for a time, the evils resulting from an excess of the two kinds of food. The chemical changes and evaporation going on in the skin draw the circulation to the surface of the body, just as the flame of the lamp draws the oil up the wick. Whilst the high temperature lasts this increased action of a flow to the surface is kept up. It is probable that the action of the heart is thereby made feeble by the excessive suction of the skin, as we see it frequently is temporarily by the perspiration bath. As soon as the temperature falls the blood ceases to flow in excess through the skin. According to the degree of cold it is almost driven from the surface. It accumulates within, and congestions and inflammations are produced. Free action on the inner or outer surface of the body for a time relieves the congestions, and enables the circulation to proceed.

After such alternations for years, the resident in the hot climate returns to a far colder home. There no heat leads the blood to the surface; it accumulates in the enlarged capillaries of the internal

viscera. The outward appearance is that of anœmia; whilst in

reality an internal plethora frequently exists.

The more frequent and severe the temporary congestions of the viscera have been, whilst in the hot climate, the more permanent does the internal congestion become, when the surface is constantly exposed to cold. Though the surface and extremities may look void of blood, the capillaries of the liver and spleen will often be full; and the enlargement of the internal capillaries which was the effect of temporary congestion for years, becomes permanent when the coldness of the atmosphere leaves the blood to be circulated by the enfeebled heart alone.

The consideration of the effects produced by migration, during a state of disease, from a cold to a warm and moist climate, says Dr. Copland, is of the utmost importance. Keeping in mind its influence on the healthy frame, chiefly in exciting the functions of the skin and liver, and diminishing those of the lungs, we are left

to prescribe it in various diseases.

In hæmoptysis this change is obviously beneficial, especially a warm and moist atmosphere, by this mode of operation, let the activity of the pulmonic circulation, and the disposit sanguineous exudation from the surfaces of the bronchi; broand tubercular phthisis are also often benefited, and the of the latter much delayed by this change of atmosphere, ewhen adopted early.

Chronic rheumatism is sometimes cured by this measuringly owing to its influence in promoting the biliary and

functions.

Dropsies, particularly anasarca and hydrothorax, a few instances, removed by a change to a warm whilst a moist state of the air is most serviceable in hæmorrhagic diseases, dry warmth seems more ber sies, dyspeptic affections, and hypochondriasis, evereffects in augmenting the insensible perspiration monary exhalation, and imparting tone to the cap

Besides these, he adds, gout in its early stag and scrofula in nearly all its forms, are benefit

a warm or even a mild atmosphere.

## THE RAINY SEASON.

Although medical authorities have not be estimate the effects of moisture, either action nation with heat, yet it is certain that this rious than either applied separately. In obesity and laxity of frame are induce

early observed; thence the proverbial acuteness of the Athenians,

and the sluggishness and stupidity of the Bœotians.

The effect of situation upon the habit may in some degree depend also on the gravity or weight of the atmosphere connected with locality. When the barometer is high, we feel vigorous and cheerful; when it sinks, languor and low spirits oppress us. Bichat states that, "accumulations of fat are said to take place in some animals in a few hours, in certain states of atmosphere. During a fog of twenty-four hour's continuance, thrushes, wheatears, ortolans, and red-breasts are reported to become so fat that they are unable to fly from the sportsman."

In Bengal, as on the West Coast of Africa and other unhealthy climates, the heat and moisture combined cause a vast increase of minute vegetable and animal life, while the decomposition of dead animal and vegetable matter is equally rapid, showing the aptitude of all substances to pass from the inorganic to the organic, and vice versa. The oxidation of metals proceeds with immense rapidity

also during the rainy season.

During the first month of this season, the temperature falls considerably, accompanied by a freshness of the air delightful to the senses, after the previous excessive and dry heat. The monsoon is steady and veers to the south and south-east; vegetation springs up with all the exuberance of a tropical climate, promoted, as is supposed, by an electric current of increased activity between the atmosphere and the earth; and the dust, so offensive at all

other seasons, subsides and is washed away.

From the 15th July to 15th October, and as the rains advance, we live in an atmosphere having all the properties of a tainted vapour-bath; and when the wind comes sifting through the Sunderbunds at south-east, we experience many of the inconveniences ascribed by Hennen to the sirocco of the Mediterranean, which " without affecting the thermometer or barometer in any remarkable degree," yet inflicts on the delicately sensitive human frame a feeling of indescribable languor and oppression, with an exhausting perspiration, much like what we suffer from in Bengal during the latter portion of the rainy season, and which a West Indian lady, speaking of the sirocco, described as giving "the feel as if she had been bathing in a boiler of syrup." This is the moist sirocco of Bengal. The mind, too, seems to partake in the general relaxation, being unfitted for vigorous or sustained effort; in short we here perceive the capiplenium, languor, et expletio, remarked by Petronius amongst the luxurious and dissolute Romans of his time. The muscular system, and that of the heart, are relaxed and weakened; so that, and after a time they become irritable and very defective in tone. These circumstances, together with the influences of malaria on the nervous system, appear to me to occasion the intermitting pulse so common to the old Indians.

heat in the system during the time of repose. But may not this greater susceptibility to disease be also promoted by the known increase of putrefaction in dead animal and vegetable substances during the night, resulting from the rapid abstraction of heat from the earth's surface by radiation, and the consequent production of moisture in the form of vapour, dew, or water—all productive within the tropics of vast increase of decomposition?

The abortions of the rainy season appear to arise from simple congestion and relaxation aided and promoted by diminished nervous power—the natural results of an excessive and exhaust-

ing humidity, joined to a high temperature.

Humid air, says Dr. Edwards, at an equal or even superior temperature, produces a peculiar sensation of cold which differs, not in its intensity, but in its nature. It is more profoundly felt, and seems to penetrate the whole system, and particularly disposes to paleness and shivering. By these characters I could v mistake a species of refrigeration, which consists in the dimi

tion of the power of producing heat.

In dry air, on the contrary, a sensation is experienced, whi called a sharp cold, and which designates rather the nature the degree of sensation; moreover, it is superficial, and why reduction of temperature is not too great, an increase of ac' experienced; the skin reddens, and in extreme cases the have a tendency to stiffen, instead of yielding to the and involuntary motions, which constitute shivering. seen by this comparison, and by what we have stated damp cold must tend to produce in individuals, wh developing heat is rather feeble, the series of action stitute the accession of an intermittent fever, esp are exposed to that action during sleep. The con will be found in the study of medical topography. number of cases these fevers are ascribed to ma fine weather, but others occur in places and at s the atmospheric constitution which we have minates.

A familiar but emphatic illustration of the e may be seen in its influence on the habitation structed of the finest known materials, wheth and of such solidity that in England they turies, and in Upper Egypt for a thousan through the destructive influence and sever alone, rendered in a score of years, or le crows; in much less time, indeed, they heap of rubbish covered with vegetation overflowed by the forest like the wave wet seasons, and the traces of man a productions of nature."

Of all the causes which thus render household and other property so surprisingly perishable, humidity would seem the most influential.

### THE COLD SEASON.

I believe it was Charles the First who described the best climate as that in which a man could bear exposure during the greater number of hours at all seasons. On this view our climate of Calcutta is assuredly one of the worst, for even during the cold weather, from the end of October to the beginning of February, an European cannot be exposed for any length of time with impunity; the hot sun and cold parching wind, with its evening and morning rawness, causing the most uncomfortable feelings of external dryness and internal fulness, unless it be in persons of youth and robust health, and under exercise sufficient to determine moisture to the surface. But, with all its disadvantages, this is the season during which the European soldier should be initiated into the climate of Bengal, and of India generally. This truth is now beginning to be understood by the authorities; and let us hope that we may never again witness the arrival of recruits from England, as of old, in the month of May, the hottest of the year. It was a cruel and destructive custom.

The unfavourable influence which the north-east monsoon exercises on the general health of persons of a feeble constitution, or who are ailing, may in part be ascribed to its relatively low electric state, or its being in a negative state, thus attracting the positive electricity of the animal frame, as well as that of the soil, If the powerful physical agent of electricity is that which, through the system of organic nerves, influences the various secretions, how much must our climate, and consequently our health be influenced by those causes, whether general or local, which affect the proportions of electricity in our atmosphere always greater and more uniform than in European climates. Coming down upon us directly from the frozen platform of Central Asia, the temperature of the monsoon is also relatively reduced so as to absorb much of our terrestrial heat, while, at the same time, its hygrometric capacity is much increased, parching up the soil along with the animal and vegetable fibre.

"The temperature of the blood is about 98°, at which it is sustained with some slight variations even when the body is immersed in an atmosphere of 29°, or of a still lower temperature. But the heat that is lost by the lungs and the skin is so great, that it is not supplied with sufficient rapidity in the young and the old; who,

to use a common expression, do not 'stand the cold,' so well as men in the prime of early manhood. The above numbers show that the power of cold on life varies according to definite laws; thus the mortality by cold is (35) twice as great under the age of 20 as the mortality (18) at 20—40; but after that turning point the power of resisting cold decreases every year, and men of 90 and men of 30 have suffered from the cold that we have experienced in the proportion of 100 to 1 (or of 1,749 to 17.5)."—Dr. FARR, February 27th, 1855.

People, on coming into Lower Bengal from the Upper Provinces during the cold season, perceive an extraordinary change in the condition of the atmosphere on first approaching the Delta of the Ganges; the bracing elastic cold of Upper India is exchanged for that of a damp cellar, and thus they invariably describe it.

At the commencement of the cold season, in October, the temperature and the winds are variable, the drying process is in full activity, and the unhealthiness is great. In Colonel Tullock's "Statistical Report on the West Indies," it is stated that, though the months previous to the cold ones have the most sickness, yet "the principal mortality is during the cold dry weather which generally prevails at Christmas." In Aracan also, during the Burmese war, it was found that, "in September and Octob when the rain began to abate, the fever was equally preval and still more fatal than in the preceding months; and in vember, when little or no rain fell, the disease appears to been at its height."

From the 1st of November to the end of February the is settled and agreeable to persons in health; but to the and sickly the altered balance of circulation and nervou occasions much discomfort. The monsoon keeps steadil east, the atmosphere during the day is dry, and a slight place in the barometer—the thermometer ranging from the nights are damp as well as cold during this dewing process being excessively productive; and to prevail occasionally are of a nature more dense the where seen, except in Pegu.

On the other hand, during the day, the cold absorbs moisture with extraordinary rapidity from animate and inanimate, over which it passes. From ade of the most seasoned wood, foreign or cracks audibly; plaster newly laid falls from rapidity of evaporation; the old Indian beer and shrivelled, with a sense of dryness in the so uncomfortable, as to give to some persons in whom the power of generating heat mylong residence in India, a constant sense

of the whole sentient surface, not to be described. "I can bear the chilling blasts of Caledonia," says a Scotchman, quoted by Ward, "but this-this cold, I know not what to do with it;" indeed, it requires a degree of equilibrium of health, not commonly enjoyed by persons of long residence in Bengal, to take kindly to the alternation from copious and incessant discharge from the surface, to its total suppression, and consequent abdominal and cerebral engorgements. It is only to the sound of constitution, who are temperate in all ways, and thereby able to bear the coldbath, or to such as are recently arrived from Europe, that our cold season is either agreeable or healthy. But here I would observe, that the degrees of health and disease are not to be measured by the thermometer, taken singly-such observations in connection with health affording little information to the medical inquirer; and it has been well observed in our own country, that the influence of the weather on the human frame is not to be thus estimated. An east wind, with the thermometer at 56°, will impress the body with a more chilling effect, than a south-west wind, when that instrument indicates a temperature 10° lower; and a foggy atmosphere, in like manner, much more injuriously than a clear one of equal cold. In the Statistical Report on the West Indies. already quoted, it is stated that in Jamaica the lowest mortality throughout a long range of observation, was in the three months prior to June 1827, when twelve deaths only were reported through the whole quarterly returns. The next quarter remittent fever broke out, and among a diminished force the deaths amounted to 252 in the same period, without the thermometer indicating any further increase of temperature than about 3°, and without any remarkable change in the weather.


The diseases more peculiar to the cold season in Bengal arecongestive fever of the continued form; intermittents, with the sequelæ of tumid or indurated liver or spleen; hepatic insidious sub-acute inflammation, terminating rapidly in abscess, if not promptly and vigorously treated; dysentery, frequently complicated with hepatic congestion-all more or less acute, according to individual habit, or length of residence in India. Catarrhs and bronchial affections are not so severe or so prevalent as the contrasted nature of the season would lead us to suspect. Hæmorrhoids with many persons follow immediately on the drying up of the surface, and consequent fulness of the internal vessels. It results also from the cold season that the blood is more venalized, a larger quantity of oxygen being consumed in the maintenance of the temperature of the body; and this condition of the blood, aided by congestion, would seem to dispose to apoplexy, the most fatal forms of which that have come under my observation in Bengal having occurred at this season; and paralytic affections are not unfrequent amongst the aged and infirm. Chronic tions, in an active state of development during the previou season and rains, now become suddenly repelled, accompani

headache and vertigo.

In old residents the appetite fails, accompanied by an or sive sense of abdominal fulness; and when this state is no by a suitable change of diet, clothing, and some medicine upon the skin and bowels, visceral congestion, ædema of the extremities, or some more active disease may ensue. New infants suffer materially from, and are sometimes destroye the impression of our cold season on the unguarded and de surface. The abortions of the cold months have alway peared to me connected with acute venous congestion, an intermitting states of the pulse and epigastric pulsations con to old Indians at this season are increased by this state great venous trunks. The kidneys act during the contin of the cold weather with diabetic violence, the urine limpid, and they only cease to do so on the return of a w season, and consequent equability of circulation, causing ture of the surface. The biliary excretion, in excess of hot and rainy seasons, is here diminished, as indicated whitish or clay-coloured state of the alvine discharges, ing the reverse order of what takes place in the instr the kidneys. The function of the liver, indeed, is now de and depraved.

The reader will not fail to observe that congestion by edly been mentioned as participating in the most de our diseases. That climate has a principal share in this unfavourable state, there can be no doubt; but this, we must consider also that the absence of all exercise of mind and body, with their animating, heathful influences on all the functions, predisposes aided by a too full and stimulating diet, to this end too, the European resident in tropical climates ha remedy against the evils of this double inaction, moderation in diet, which he will not adopt; for, and rainy seasons, the amount of exercise necessar temperate regions would here be impossible, and v even if possible. Those who would preserve the fore, must be temperate,\* use such bodily exerc will admit of, and relieve their minds from the me official duty, by the inexhaustible resources of and elegant accomplishment. It is only thus

<sup>\*</sup> It has been shown by experiment, that when alcohol has I quantities into the circulation, "the arterial blood retains the doubtless through the contamination of the blood chiefly the spirits proves so baneful in hot climates especially.



piness, and reputation can be insured under the disadvantages of our position in India, where too commonly the whole time is given up to business.

From the sketch now given of the locality and climate of Calcutta, it will be seen that, without taking the specific or malarious influences into account, we are there exposed to atmospheric changes to an extraordinary degree;—to an extreme of heat and dryness—extremes of heat and moisture—cold and moisture—cold and dryness. The European exile may well join in the "complaint of the black knight" of Chaucer.

"Nowe hote as fire, nowe colde as ashes ded;
Nowe hote for colde, now colde for hete again;
Nowe cold as yse, and now as coles red,
For hete I brenne.'

That these various influences alternately excite and depress the vital processes must be evident; and it is to their long continued application (even where disturbance of function does not amount to actual disease), that we must refer the attenuated condition and

general feebleness of the old Indian.

"Bengal has received a bad character for insalubrity," says Malte-Brun, "and certainly it is in an eminent degree exposed to a succession of violent extremes and vicissitudes; at one time to excessive rain, at another to hurricanes; then to scorching heat, and frequently to thick fogs; yet the English have, by dint of prudent regimen, accommodated themselves to the climate." When Malte-Brun wrote, I fear there was not much of that "prudent regimen" to which he ascribes such happy results; but let us hope that it is now about to be attained, and that it may no longer be a reproach to us that, not satisfied with choosing the worst localities, we also adopt habits of life the worst calculated for the accommodation of our constitution to them and their climates.

As illustrative of the conservative influence of good habits of life, it is worthy of remark that, of the European officers on the Bengal establishment, there died on an average of eight years, of the unmarried class, 3.77 per cent. annually; while, of the married class, the mortality for the same term was but 2.74. Of the European soldiery again there died annually, during the period here referred to, 7.38 per cent. This wide difference in sickness and mortality is referable almost solely to the difference of habits of life; for, in the important circumstances of age and constitution, the private soldier, on an average, has the advantage. On entering the service he may be taken generally to be of a stronger frame than the commissioned officer; and he very rarely serves to so advanced a period of life as the latter. With the exceptions of coarser diet, a habitation inferior to the officer, and harder work

on actual service, circumstances which do not generally tell much on the amount of sickness and mortality, the soldier has thus some real advantages over the officer—excepting in the all-important

one of habits of life.

Referring, lastly, to the influence of habits of life and of climate, in preventing thoracic diseases, I would quote Dr. Atkinson, of Wakefield, and others, who infer, that by the use of alcoholic liquids, " a state of the blood is produced, opposite to that which obtains in tuberculosis, and may thus prevent the development of it." This view is supported by Dr. Atkinson in a numerical statement, showing a mortality from phthisis, amongst the adults of Wakefield, " of rather less than one in three, and in publicans of one in twelve and a-half." Publicans are here placed in contrast with the general community as affording examples of harddrinking persons; but a moderate supply of alcoholic liquids, with a generous diet, is what Dr. Atkinson recommends as a prophylactic. This gentleman regards phthisis as "characterised by a excess of oxygen circulating in the system, and that when blood becomes too highly carbonized from various dise tubercles are not deposited."

If further observation should confirm this view of tuber and its causes, we may receive from it some assistance elucidating and determining the actual physiological ope tropical climates in preventing the development of pulm sumption. It has been seen that, during the hot weather especially, the blood is venalised, and malaria may per garded as having a somewhat similar influence; so that gonism alleged by the French military surgeons to eximate, after all, have more of foundation than did at first

STATISTICS OF THE INFLUENCE OF C SEASON IN PRODUCING SICKNESS AN IN CALCUTTA.

SICKNESS.

It now remains to trace in a more specithe influence of season in producing sich this purpose I have prepared several tab' ing this influence on the mortality of narequest, by Dr. Duncan Stewart; the seven years, viz., from 1831 to 1837, in

MONTHS.	there occurred	Of 1000 living Hindoos there died in different months.	Mussulmans	Total Natives : Of 1000 living, there died in different months.
In 7 Januarys	921	271	91	221
" 7 Februarys		121	9½ 8½ 8½ 8½ 9	164
,, 7 Marchs		184	81	15‡
, 7 Aprils		191	81	161
,, 7 Mays		171	9*	151
,, 7 Junes		141	84	13
" 7 Julys		18	114	164
,, 7 Augusts	901	26	101	214
" 7 Septembers	98	28	123	234
,, 7 Octobers	1041	30	12%	251
"7 Novembers		341	113	28
" 7 Decembers	1061	311	111	25%

This table exhibits in a remarkable manner the fatal influence of the months from September to January inclusive, and which would be even more striking if we possessed the means of deducting the deaths by cholera in the months of March, April, May, and June, so as to show only the ordinary endemic influence; as it is, however, the table is instructive and interesting.

The following table I have had collated from the records of the vestry; it exhibits burials in the Protestant burial-ground for twenty years, from 1819 to 1838 inclusive, and arranged in months:—

MONTHS.	Under five years of age.	Above five years of age.	Total.
In 20 Januarys	53	257	310
,, 20 Februarys	41	165	206
" 20 Marchs	70	260	330
,, 20 Aprils	105	307	412
" 20 Mays	104	421	525
" 20 Junes	99	298	397
,, 20 Julys	104	285	389
,, 20 Augusts	87	389	476
" 20 Septembers	74	379	453
" 20 Octobers	84	368	452
" 20 Novembers	60	379	439
" 20 Decembers	53	349	402
Total	934	3857	4791

Thus it appears, that though the greatest number of deaths occurred in the Mays—the worst of the cholera months—the period of greatest general mortality has been from August to January inclusive; and if the deaths by cholera during the hot months could be excluded, this result would appear very striking.

With a view to exhibit the simple endemic influence in the fairest manner, I again had recourse to the vestry records, from which the following table of Protestant burials has been framed for the twenty years just previously to the appearance of cholera as an epidemic, namely, from 1796 to 1815, inclusive:—

MONTHS.	Under five years.	Above five years.	Total.
In 20 Januarys	28	216	244
	26	153	179
	31	164	195
,, 20 Aprils	33	178	211
,, 20 Mays	25	235	260
,, 20 Junes	22	209	231
,, 20 Julys	34	219	253
,, 20 Augusts	44	288	332
,, 20 Septembers	42	296	33'
,, 20 Octobers	32 32 23	293 331 311	3'
Total	372	2893	

Thus it appears, that in twenty years, Novembers fatal months, and that the five months from Auginclusive, were more fatal than the whole seven

But further to ascertain the influence of se and mortality of the European soldiers and pared the following Table from a docume Medical Board:—it exhibits the totals of a in the Presidency General Hospital in each years; the item "other diseases," has some that I cannot help:—

		Т	OTAL	ADM	118810	NS.	TOTAL DEATHS.				
RANGE OF OBSERTION MONTHS		By acute diseases.	By chronic diseases.	By surgical diseases.	By "other diseases."	By all diseases.	By acute diseases.	By chronic diseases.	By surgical diseases.	By "other diseases."	By all diseases.
In 12 Januarys		899	64		245	1469	89	10	5	8	12
" 12 Februarys		557	46				67	6 5	2 5	13	88
" 12 Marchs " 12 Aprils		456 619		163 138	132 175		55 50	6	6	12	72
" 12 Aprils		722		161	178		67	11	7	12	97
" 12 Junes		791		184			64	4	4	11	83
" 12 Julys		843		176		1184	104	6	11	24	145
" 12 Augusts		732	42	180	146	1100	88	9	9	8	114
" 12 Septembers		683		148	174		112		11	3	144
" 12 Octobers		764		229	182		114	4	10	18	136
" 12 Novembers		1142		A 100 PM	325	1869	97	4	5	10	116
" 12 Decembers	40 40	948	87	247	233	1515	109	8	7	27	15]
Total		9156	727	2354	2253	14490	1016	81	82	153	1332

It appears from the Medical Board's Table, from which the above is framed, that out of 1,704 Europeans, who died in the twelve years, 372 died of cholera; 304 of dysentery; 58 of diarrhæa; 465 of remittent fever; 66 of intermittent; 50 of hepatitis; 88 of phthisis, so-called, but more than half were probably bronchitis; 19 of rheumatism; 6 only of splenic disease; 8 of apoplexy; 9 of delirium tremens; 6 of small pox, leaving 270 deaths from "other diseases."

The total admissions in twelve years are 15,293, and the deaths, 1,704.

The proportions of deaths to cases treated are as follows:-

Of Remittent Fever, 5,116 admissions and 465 deaths, or one in	11
Of Dysentery, 1,877 admissions and 304 deaths, or one in	61
Of Cholera, 853 admissions and 372 deaths, or one in	
Of Diarrhœa, 608 admissions and 58 deaths, or one in	104
Of Intermittent Fever, 501 admissions and 66 deaths, or one in .	7-7-
Of Hepatitis, 446 admissions and 50 deaths, or one in	8,9
Of Spleen diseases, 58 admissions and 6 deaths, or one in	93
Of Delirium Tremens, 124 admissions and 9 deaths, or one in .	137
Of Apoplexy, 27 admissions and 8 deaths, or one in	34
Of "other diseases," 2,253 admissions and 153 deaths, or one in	143

Average of deaths from all the diseases, is therefore nearly one in nine.

Of a hundred British soldiers who die in Bengal, Dr. Burke states that 26.8 are from fever; 7.3 from hepatitis; 30.5 from dysentery and other bowel complaints; 19.5 from cholera; 4.6 from pulmonic

diseases, 1.9 of which is from phthisis pulmonalis; leaving only

11 produced by other diseases.

From the Medical Board's Table of Seasons, it appears that the most healthy months comparatively, are February, March, April and May; that from June to January, the admissions range high, these also being by far the most fatal months. These results correspond nearly with observations on the climate of Jamaica, and others of the West Indian Colonies, wherein the most unhealthy as well as fatal months of the year extend from August to December inclusive; the months comparatively healthy being, from January to June. Such observations would seem to apply to most parts of the Northern Hemisphere, which generally possess the same character of climate: they also accord with the ancient maxim in the south of Europe which held the summer and autumnal to be the most sickly seasons.

Celsus, writing of the influence of season in the malarious city of Rome, says, "Igitur saluberrimum ver est; proxime deinde ab hoc, hyems; periculosior æstas; autumnus longe periculosissimus

From the researches of statistical writers, it appears that all over Europe the maximum of deaths occurs towards the close of winte

and the minimum towards the close of summer.

In order that the table should express only the result of ordir season and endemic influence, I have not included cholera, v epidemic is found to prevail chiefly in the months of April, and June.

October, November, December and January are the mor which there occurred the largest proportions of remitter cases, and the same applies to dysentery, diarrhœa, and a patitis; while Septembers alone give more intermittents whole of the other months besides.

In order to account for the unusual mortality under disease, it is proper to mention that the description treated in the General Hospital, are as follows :- Euro belonging to detachments from Her Majesty's and th Company's regiments; recruits for ditto; the sick of returning from service, as for instance, during the war, &c.; European and American seamen from townsmen, paupers, generally seamen; invalid s way to England, &c. &c.; -in short, a class of whom, besides being diseased and worn out, a salutary control which leads to prompt measu prevention and cure of disease. It thus happy stages of acute illness are neglected; an irred which must necessarily imply a larger proport

The following table exhibits the admission month, during the period included between

Hospital of Her Majesty's Regiment in the Garrison of Fort William.

	TOTAL ADMISSIONS.					TOTAL DEATHS.				
MONTHS.	By Fevers.	By Hepatitis.	By Bowel Complaints.	By other Diseases.	Total by all Diseases.	By Fevers.	By Hepatitis.	By Bowel Complaints.	By other Diseases.	Total by all Diseases.
In 12 Januarys	231 266 323 303 418 488 403 427 446 404 423 311	57 68 38 42 33	249 263 286 318 330 253 314 321 256 250 203 202	557 541 574 550	1395 1301 1228 1239 1083	7 1 9 6 8 5 16 18 18 23 19 16	3 2 1 2 2 2 2 5 4 4 0 4 1 1	20 16 16 19 22 18 28 25 30 21 25 16	10 5 10 9 12 13 10 8 10 6 9 10	40 24 36 36 44 38 59 55 62 50 57 43

It thus appears that in the 12 years there were 15,564 admissions by all diseases, and 544 deaths, or one death in every  $28\frac{3}{5}$  patients treated, being less by two-thirds than the general average afforded by the General Hospital returns for the same period—an emphatic example of the value of that internal discipline and economy in corps to which so much preventive and curative result is ascribed by all military and medical authorities. Out of 544 deaths, 256 are by bowel complaints—146 by fever, and 30 by liver disease, leaving 112 deaths by other diseases. If, from the column of deaths, we abstract, for the hot months, those by cholera, the comparative fatality of the months from September to January inclusive, will be rendered very remarkable.

The proportions of deaths to cases treated are as follows:-

By fever 4,443 admissions and 146 deaths, or one in 30 and two-thirds, nearly: by bowel complaints 3,245 admissions and 256 deaths, or one in 12½, nearly: by hepatitis, the very large proportion of 555 admissions and 30 deaths, or one in 18½: by other diseases, 7,321 admissions and 112 deaths, or one in 65 and ½rd, nearly.

In order to exhibit the difference of result in the two hospitals, I subjoin the following table:—

Diseases.	Ratio of Deaths in the Regi- mental Hos- pital.	Ratio of Deaths in the General Hospital.	Difference.	
Fevers	One in 30g	One in 10g	Of 20 nearly.	
Bowel complaints	" "12 <u>1</u>	" "6 <sub>10</sub> {	,, 6, ,,	
Hepatitis	,, ,, 181	,, ,, 8%	" 5½ " " 9½ "	
Other diseases	,, ,, 65 <del>1</del>	,, ,, 143	,, 50% ,,	

Under the head bowel complaints it is not known whether cholera, dysentery and diarrhoea are all comprehended. This laxity in framing the table, and the lumping of 7,000 odd cases under the vague designation of "other diseases," goes considerably to vitiate the regimental return; but both errors arose from iapprehension on the part of the compiler, who is at the dist station of Meerut.

The table, with all its faults, is not without value as bearir the three grand divisions of tropical disease,—viz., fevers, diseases, and those of the liver:—but comparison cannot, o' be made, to any just or useful purpose, between the two l as the one treats only a selected class of men whose mea 26; while the other treats the broken down and miseral ages, and in all the stages of their diseases.

The winter, all over Europe, is the season of the gretality; and not only is this a proved fact in statistics, winter is found to be attended by a diminished mort ratively. In the first quarter of 1846, the winter wild throughout England, and "the rate of morte than in the corresponding quarters of eight previous."

In the first quarter of 1847, on the other hand, was below the average, the severity of the weath out unusually great. The mortality correspond above the corrected average," for the 117 district in quarterly returns of the registrar-general.

"Winter," says Dr. Farr, "appears to be the is most natural to man to die." The same author that "as man progresses in life, the more not become to him; while the summer lengther his days."

Dr. Mouat, of the 13th Light Dragoons, in the Madras Presidency, out of 3,394 c

Statement of the average rate of Mortality per cent. among the different classes of Inhabitants in Calcutta per Census, and Table of Mortality.

Denominations.	Number of Inhabitants.	Total.	Average Mortality per annum.	Average Mortality per cent.	
English Eurasians	4840	7884	277	34 per ct.	1 in 28
Portuguese	3.00	,		347	
Western Mahomedans.		3341	417	124 "	1 in 8
Bengal Mahomedans . Moguls	45,067				
Arabs	0.00	59,622	1607	24 "	1 in 36
Western Hindus . Bengal Hindus .				11	
Mugs	683				
		157,418 636	9558 25‡	61/2 ", 41/2 ", 24 ",	1 in 16 1 in 25
Native Christians .		49 362	31	24 ,,	1 in 14
Jews		307			
27 3	: ::	40 55			

"The great difference in mortality between the Hindus\* and Mussulmans is striking, while the difference to be observed between the Portuguese, as compared with the English and the Eurasians, is equally so.

"Here is much room for speculation, and it cannot be said that as yet we have as good means of getting correct information upon this subject as they possess in Europe; nevertheless, we may approach as near as we can to the point we wish to ascertain, and

we may hope to improve in such statistical records.

"The Portuguese, among whom so great a mortality is shown, are a suffering race very subject to the catalogue of complaints enumerated in these papers; while the English and Eurasians are far more prosperous in life, and enjoy comforts and happiness in a very high degree, as compared with the former section of society. The mortality of English and Eurasians, 3½ per cent.

<sup>\*</sup> The difference of mortality amongst the Mahomedans and Hindus may be accounted for by the circumstance that the Hindus of Calcutta, consisting of families, include a much larger proportion of infant life. The same circumstance will explain the great difference between the average mortality amongst the Portuguese and the Europeans of Calcutta.—Ed. Journal Asiatic Society.

per annum, while that of the I very great. In 1830 I ascertain ings of Science,' the burials in from the year 1820, to show at pean population must have grea and burials had not increased; is acknowledged to have increa see upon referring to the first co the Protestant burials, for the deaths. The years 1833-4, the dations, show the greatest mor the native population those two tality. The two last years sho native population that healthi among the other columns of so nian, Hindu, Armenian and N twenty years, and I believe th Chinese and Jews keep no acco could not include them, and th population of this city."

"The native soldiers of Beng very healthy under ordinary circ inquiry that only one man die actual strength of the army. Proper to this class of natives, a vinces, that though only one-fe stationed in Bengal, the death moiety of the whole mortality re

That the climate of the count the Ganges, was not less fatal t is evident from Gladwin's transle. "In former reigns, Bengal, on air and water, was deemed inimicand other foreigners, and only the royal displeasure were stat which enjoys a perpetual spring as the land of spectres, the seat death."

"The Mussulman invaders," s
Hindustan, who afterwards estab
Delhi, considered this country, B
region; and whenever any of the
guilty of capital crimes, and the
mit their being beheaded, while
their removal necessary, they we
and water of Bengal were consid
death of the criminal."

being 121 per cent., is ublished in the 'Gleanof Protestant Christians that although the Eurosed, yet, that the deaths hat the same population naterially indeed, yet we ne of the tables, giving ty years, no increase of following the sea inunate years; while among w an extraordinary morof both European and estored. The mortality Catholic, Greek, Armeistian, are for the last e nearly correct. The ir burials; I of course minute portion of the

colonel Henderson, "are in the bus, is less than the bus, however, is Bengal and with the Upper Prohetroops exhibited are fourth are more than a

reat waters, the Delta of hommedan predecessors, a Persian document: f the inclemency of the constitution of Moguls bers who laboured under re; and this fertile soil sidered a strong prison, se, and the mansion of

ve writer, "of the west of mselves on the throne of be Dojakh, or an infernal or Courtiers, were found individual did not perthe same time, rendered ed to Bengal. The air I as to lead to the certain

For the following Table, showing the Mortality of the Native Population of Calcutta, from various Diseases, during the Years of 1832, 1833, 1834, 1835, 1836, and 1837, I am indebted to Dr. Dun

Į.	Average proportion of Deaths from various Diseases in every 100 Deaths.		Hindoos and Mahomedans.		23§ in 100.	
Duncan Stewart:	Proportions of Deaths from various Diseases in every 100 Deaths,		Hindoos. Mahomedans,		25 in 100,	
o Tri. Tomic			Hindoos,		22‡ in 100,	
Total Native Population, 217,095.			Annual Ratio.		7.6 per cent. 1.26 per cent. 3.94 per cent. 656 per cent. 6.6 per cent. 1.1 per cent. 224 in 100, or lin 13, or lin 15, or lin 15, or lin 90.	
	T Native P		Ratio in Six Years.		6.6 per cent.	or 1 in 15,
	Mahomedans, 59,622.		Annual Ratio.		656 per cent.	or 1 in 152,
Maho 59,			Ratio in Six Years.		3.94 per cent.	or 1 in 25,
Hindoos, 151,473.			Ratio.		1.26 per cent.	or 1 in 79,
		Datie in	Six Years.		7.0 per cent,	orlin13 nearly or lin 79, or lin 25, or lin 152, or lin 15, or lin 90.
		-	Diseases.		Cholera	

The following table is collated from the interesting and valuable article of Mr. H. T. Prinsep on the mortality for ages and births of Indo-Britons in the Lower Orphan School of Calcutta, during 40 years, and published in the Journal of the Asiatic Society for September, 1838.


		BOYS.			GIRLS.	
Age.	Number in the House, on 31st December of each year.	Total Deaths.	Ratio of Deaths per 1000.	Number in the House, on 31st December of each year.	Total Deaths.	Ratio of Deaths per 1000.
0 1 2 3	2·713 2·430 2·169 1·968 1·845	486 498 186 113 65	171·480 185·890 82·220 55·830 35·230	2·766 2·436 2·203 2·030 1·790	416 496 170 80	144-980 184-830 74-300 38-640 42-120
1 2 3 4 5 6 7 8	1·725 1·587 1·451 1·314	56 42 41 36	31·940 26·110 27·870 27·270	1·788 1·677 1·572 1·431	77 52 49 30 36	28-690 28-80/ 18-9/ 24-9
9 10 11 12 13	1·186 956 750 528 287	15 33 20 8 3	12·570 33·960 26·300 15·040 10·450	1·322 1·210 1·140 1·071 951	14 17 18 12 15	10
14 15 16 17	138 70 38 19	1 6 3 0	7·240 82·190 76·920	799 626 449 295	5 4 6	,
18 19 20	14 11 8	1 0 1	ii	194 131 81		
Total	21-207	1.614	908-510	25:970		
Mean	530-7	4014	22%	6494		

"N.B. In forming this ratio, as the number stated for the last day of the year, and there the deaths during the year, it has been ne deaths, on the calculation that each person of his life for half the time of those who life

By a statement forwarded by the Me mittee for the improvement of Calcu

For the following Table, showing the Mortality of the Native Population of Calcutta, from various Diseases, during

	Hin 151	Hindoos,	Маноп 59,6	Mahomedans. 59,622.	Total Native Population, 217,095.	Total ; Population, 217,095.	Proportions of Deaths from various Diseases in every 100 Deaths.	Proportions of Deaths rom various Diseases in every 100 Deaths.	Average proportion of Deaths from various Diseases in every 100 Deaths.
Diseases.	Ratio in Six Years.	Annual Ratio.	Ratio in Six Years.	Annual Ratio.	Ratio in Six Years.	Annual Ratio.	Hindoos.	Hindoos, Mahomedans, Mahomedans,	Hindoos and Mahomedans.
Cholera	7.6 per cent. 1.26 per cent. 3.94 per cent. 656 per cent. 6.6 per cent. orlin13 nearly or lin 79, or lin 25, or lin 15,	1.26 per cent. or 1 in 79,	3.94 per cent. or 1 in 25,	656 per cent. or 1 in 152,	6.6 per cent. or 1 in 15,	1.1 per cent. or 1 in 90.	22‡ in 100,	25 in 100,	23§ in 100.
Small Pox	1.77 per cent. '295 per cent. '768 per cent. '128 per cent. 1.5 per cent. or l in 55, or l in 339, or l in 130, or l in 782, or l in 66,	·295 per cent. or I in 339,	.768 per cent. or 1 in 130,	128 per cent. or 1 in 782,		·25 per cent. or 1 in 400.	5½ in 100,	6 in 100,	5‡ in 100.
eous		-	·11 per cent. 1.8 per cent.	+4	21.3 per cent.	21.3 per cent. 3.55 per cent,	72} in 100,	69 in 100,	70§ in 100.
Diseases	or lin 4 nearly, or 1 in 24,	1	or lin 9 nearly, or in 54½,	- 1	or 1 in 4\frac{2}{3},	or 1 in 28.		-10	
	34.5 per cent,	34.5 per cent, 5.7 per cent, 15.7 per cent, 2.6 per cent, 29.4 per cent, 4.9 per cent,	15.7 per cent. 2.6 per cent. 29.4 per cent	2.6 per cent.	29.4 per cent.	4.9 per cent.	100.	100.	100.



able rths ring for

> of per

0000

0000

10

is

he

n

during the thirty years, from 1st January, 1808 to 31st December, 1837, there were of admissions and deaths amongst the European patients of the Presidency General Hospital as follows:—

This, for thirty years, gives a mean ratio of  $10\frac{27}{100}$  deaths per cent. of the admissions.

The quinquennial ratios are as follows:-

						Admissions.	Deaths.	Ratio of Deaths per cent.
From	1808 to 1	1812		141		2.713	299	11
11	1813 to 1	1817				4.360	450	10,32
11	1818 to 1	1822				6.425	657	10 22
"	1823 to 1	827				9.560	1039	10 76
73	1828 to 1	1832				6.315	585	9 26
"	1833 to 1	1837				5.746	577	10,50
	A	fean			y.	5.8531	684%	10 27

This is all that the Medical Board can furnish; for it was not till 1808 that any records were used, even of admissions and deaths.

The following Table exhibits both the sickness and mortality of the British troops in garrison at the Presidency, together with the proportion of deaths to cases treated of the three principal endemic diseases of Bengal; viz., fever, dysentery, and hepatitis:—

Table of Admission into Hospital, and Deaths during 12 Years of the European Troops in Garrison of Fort William.

				Ratio per 1000	of Streng
Years.	Strength.	Admissions.	Deaths.	Admitted.	Died.
1822	866	1303	75	1,594-62	86-60
1823	828	1687	51	2,037-24	61:59
1824	736	2268	103	3,082-32	139-94
1825	902	2542	110	2,818.18	121.95
1826	863	1826	96	2,115.87	111:23
1827	893	1336	56	1,496.08	62.71
1828	913	1776	42	1,945.23	45.00
1829	885	1995	58	2,253.67	62.14
1830	808	1722	59	2,131.78	73.02
1831	831	1061	57	1,276.77	68.59
1832	771	1024	59	1,728.14	76.52
1833	687	1387	64	2,018.92	93.15
1834	608-	1166	57	1,917,76	93.15
1835	743	1211	33	1,629-87	44.41
1836	734	1245	25	1,696.18	34.06
1837 .	709	899	26	1,267.98	36.07
1838	633	694	22	1,096.36	34.75
Total	13:410	25-142	497		
Average	78814	1478 \$	29,4	1,833-33	73.26

The ratio of deaths to cases treated, was, during ten years, as follows :-

Table of Admission into Hospital, and Deaths during 12 Years, of the European Troops at Chinsurah (Depôt), 18 miles distant from Calcutta.

2 312		-	10	Ratio per 100	0 of Strengtl
Years.	Strength.	Admissions.	Deaths.	Admitted.	Died.
1826	1,083	818	86	755-31	79-40
1827	814	1,329	80	1,632-67	98.28
1828	200	542	20	2,710.00	100.00
1829	132	571	24	4,325.75	181-81
1830	366	822	24	2,245.95	65.57
1831	927	1,412	51	1,523.19	56.01
1832	737	1,182	41	1,603.79	55.63
1833	577	1,155	54	2,001.73	93.58
1834	784	1,308	32	1,668.36	40.81
1835	832	872	17	1,048.07	20.43
1836	743	1,428	39	1,921.93	52.35
1837	765	1,322	32	1,728·10	41.83
Total	7,960	12,761	500		
Average	6631	1,063-4	41%	1,930-40	73.72

The ratio of deaths to cases treated, was, during three years, as follows:—

 In Fever
 one in  $25\frac{\pi}{4}$  

 " Dysentery
 one in  $7\frac{25}{65}$  

 " Hepatitis, Acute and Chronic
 one in  $8\frac{10}{10}$ 

The above must be taken as the mortality on the spot only, and does not include the invalids, who died on their passage to England, or shortly after their arrival there, amounting to 3½ annually. If, as is well known also, the Medical Returns be taken at the highest strength in course of the year or quarter, or a tenth part higher than it ought to be, instead of the mean monthly strength, the ratio of mortality will be greatly increased. It is believed that a correction of all errors would bring the annual mortality to more than eighty per thousand.

That in India as elsewhere age materially influences the ratio of mortality will be seen from the following tables for the officers of the Bengal army, and those of the Civil Service. Out of 1184 deaths among officers, the proportion occurring annually in each rank, and at each age, has been as follows:—

1		Colonels, average age 61.	Lieutenant Colonels, average age 51.	Majors, average age 40.	Captains, average age 36.	Lieutenants, average age 18 to 33.	Cornets and Ensigns, average age 18 to 33.	General average at all ages.
-	Died annually per thousand of each Class	59.4	48-4	41.0	34.5	27.5	23.4	31-2

The mortality among the civil servants, for a period of forty years, from 1790 to 1836, exhibits almost precisely the sam sults, viz.

	Above 50 years of ageand 30 of service.	Age 40 to 50; service 25 to 30.	Age 40 to 45. service 20 to 25.	Age 35 to 40; service 15 to 20.	Age 30 to 35; service 10 to 15.	- 14 30;
Died Annually per thousand of each class	48.6	36.4	35.4	23.4	16.6	

Between ten and fifteen years' service is the per of absence is allowed to those who choose to return three years, which of course must have a matareducing the mortality of that class. With results are uniform for both civil and military are no less so when extended to the officers dencies.

The official results in regard to the mortalithe military officers and civil servants of the afford a convincing proof that in the East hitherto been derived from length of residuals are never employed out of India, an about the age of eighteen or twenty, the ranks may be assumed as a criterion for residence in the country. On that printed equal numbers of each rank, that the mortality and the mortality is a server of the country.

for the most part youths but recently arrived, is only twenty-three; while that of the Lieutenants, who must have been at least three years longer resident to have attained that rank, is twenty-seven; and that of the Captains, who must have been about twelve or thirteen years longer, is thirty-four per thousand, and so on in a corresponding proportion with the higher grades. In case it should be objected that this does not exhibit the precise operation of mortality during the first year or two of residence in the country, when the influence of acclimation is supposed to be most strongly manifested, the following information in regard to the Civil Servants in the Bengal Presidency will supply that defect.

	Numbers alive.	Deaths in first year.	Ratio of Deaths per thousand of living.
1st Year of Residence.	 975 933	19 22	19.5
2nd " " 3rd " "	 906	18	23.5
4th " "	 874	19	22.0

Here, then, we have traced the same individuals through four successive years of residence, with the liability to mortality constantly augmenting; and unless we are to suppose that a different law regulates the mortality among Europeans in the tropical climates of the eastern and western hemisphere, we are inevitably led to the same conclusions which we have already demonstrated from the previous numerical results.—Colonel Tullock's Reports.

The comparative annual per centage mortality of the officers belonging to the armies of the three Presidencies of Bengal, Madras, and Bombay, is stated by Colonel Henderson to be as follows:—

The subject of Statistics is one that has been altogether neglected by the medical authorities of the Bengal Presidency; and, though the example of the hospitals of her Majesty's army, from which very complete reports are made, has now been many years before them, this important branch—the very central point of medical science—can scarcely be said, until very recently, to have been approached. It is the deficiency here complained of that has caused the omission in this Article of all mention of the hospitals of Calcutta, and of the General Hospital in

particular :- an institution that has existed for more than seventy years, and in which tens of thousands of European soldiers have been treated under three or four different medical systems, yet no one fact, out of the numerous and important observations made during that long time is known to any one of us. Its surgeons, and those of the other public institutions-many of them able and experienced officers-have, through the neglect of the controlling medical authorities, been rendered, in respect to us and to science. no more than a set of dumb actors in the circle of a routine duty. Dr. John Macpherson forms a very honourable exception, his statistical investigations prepared at the General Hospital of Calcutta, being works of standard value. All this official neglect and want of discipline has been repeatedly urged by me in what I thought the right quarter, and in the most emphatic manner; but though my proposition met with no very flattering reception, 1 have yet the satisfaction to know that I produced some action, with the civil authorities, tardy perhaps, yet such as will lead to some ultimate improvement. We are in India continually kept in mind of that law of our nature, by which old men are disinclined from undertaking anything, however excellent, of which they cannot expected to see the end. Through the operation of climate; we have too often to lament the premature display of the tracting influences of age on the moral and physical constit of man, to the prejudice no less of public welfare than of bappiness.

From the circumstance that no general mode of reg and recording has existed till lately, on statistical pri which all should adopt, the great majority of experience has hitherto perished with individuals, and the valuterials of a long and active life have been thus for e

the public service.

The labour of analysing observations which have larly kept is immense—to some men impossible; whi a habit of observation is fostered and kept alive, thr creasing facility in the process, and the interest e subject. Such are the opinions of some of the al England; but they have as yet found no echo in our board, where we have still to regret the leaden is exclusive seniority principle of promotion.

To insure precision and uniformity in reports, of hospitals is systematic and strictly ordered in and this is one of the many advantages its play

possesses over ours.

As to the constantly recurring questions, do an improved climate, and is not the mortality of diminished of late years? It is not very easy torily, owing to the absence of statistical record. That the climate of the actual site of Calcutta is improved to a certain extent, there can be no doubt; and that European general mortality is likewise diminished, amongst the better classes especially, I believe to be true; but the chief cause will be found in the improved habits of the latter class; for, with the troops in garrison, notwithstanding the improved discipline and interior economy of modern times, it would not appear that mortality is much, if at all diminished, and the same may be said of the nearest military station of former

times-Berhampore.\*

The decrease in the proportional mortality of British troops in our various colonies, is stated by Robert Jackson to be only of recent date; and he adds, that it would not be safe to pronounce positively whether it is owing to improvement in medical management, with improvement in military economy, or to contingent and temporary change in the nature of morbid causes, producing a less aggravated form of disease than belonged to other times. The soldiers did not die in greater numbers in Germany and Holland in the war of 1793, 94, and 95, than they did in the war of 1803. Further, he says, that the mortality was not greater, cateris paribus in North America in the war of 1756, than in the revolutionary wars of 1775—or in the late war of 1813. It is in the navy, he might have added, that a great and progressive reduction in mortality has been effected by the measures of preventive medicine instituted in 1779.

Mortality has been nearly equal at all times in the West Indies, viz. in the war of 1756, in the war of 1778, the war of 1793, and

the war of 1803.

Of the earlier health history of the European troops in Bengal, we have little or no information. I must therefore content myself with the following tables for Fort William and Berhampore, made up from scattered records procured with much trouble, from the offices of the Auditor and Adjutant-General. After what has already been said, they require no comment.

<sup>\*</sup> Making the corrections stated at page 55, the table for Fort William will give 88·40, and that for Berhampore 102·69, as the ratio of deaths per 1,000. All the tables of the military have been framed from four official sources—viz. the Adjutant and Auditor-General's Office, that of the Inspector-General and the Medical Board; and where they differed I took the estimates of the Audit Office. More deaths were recorded in the Adjutant-General's returns than in any of the others. Slight errors have been made by the writer in calculating some of the Tables, but they in no manner affect the general accuracy of the whole.

Table showing the deaths amongst the European troops in Fort William during the years specified:—

Years.	Strength.	Deaths.	Ratio of deaths per 1000 of strength.	Remarks.
1790 1791 1797 1800 1801 1802 1805 1806 1807 1808 1809 1810	967 316 867 848 916 1020 927 1122 1198 1319 1182 1106	88 26 117 108 65 94 53 55 49 107 71 33	91·00 82·34 134·02 127·35 70·96 92·15 57·17 49·02 40·90 81·12 60·06 22·83	11 Months in Garrison. 7 Months in Garrison.
Total .	11,788	866		
Average	9824	72%	76-40	

Table showing the deaths amongst the European troops at the Station of Berhampore during the years specified:—

Years.	Strength.	Deaths.	Ratio of deaths per 1000 of strength.	Remarks.
1788 1791 1792 1793 1794 1796 1797 1798 1806	497 230 157 183 290 245 383 403 835	47 19 18 23 29 35 44 46 53	94·56 82·60 114·64 125·68 100· 138·77 114·88 114·14 63·47	11 Months at the Station.  10 Months at the Station.  9 Months at the Station.  5 Months at the Station.
1807 1808 1809 1810	848 726 721 512 6030	39 46 45 30 473	45·99 63·36 62·41 58·59	
Average	46311	365	90.69	

For the following important and interesting General Abstract of Admissions and Deaths by the principal Diseases among the European troops in Calcutta, Chinsurah, and Berhampore, in Bengal Proper, I am indebted to Colonel Tullock:—

	viz. from	n 182	10 years, 27 to 1836 lusive.	viz. fro	m Ju June, from	or 8 years, ne, 1826, 1827, 1830 to clusive.	Berhampore for 10 years, viz. for 1823, and from 1826 to 1834 each inclusive.			
Diseases.	Aggreg	782	Strength,	Aggre	gate 5 6531	Strength,	Aggre	gate s	Strength,	
	Admissions into Hospital.	Deaths.	Proportion of Deaths to Admissions.	Admissions into Hospital.	Deaths.	Proportion of Deaths to Admissions.	Admissions into Hospital.	Deaths.	Proportion of Deaths to Admissions.	
Feb. Intermittens ,, Remittens ,, Cont. Com	747 706 2390	6 63 53	1 in 1241 1 in 11 1 in 45	227 278 1413	4 26 24	1 in 57 1 in 10% 1 in 59	1044 1467 5324	11 120 90	1 in 95 1 in 124 1 in 594	
Pneumonia Hæmoptysis Phthisis Pulm Catarrhus	202 11 14 554	8 3 9 24	1 in 25 1 in 3§ 1 in 1§ 1 in 23		4 3 9 4	1 in 26 1 in 11 1 in 13 1 in 54		4 1 25 8	1 in 59 1 in 10 1 in 11 1 in 661	
Hepatitis, &c	490	21	1 in 23	228	19	1 in 12	636	34	1 in 183	
Enteritis	43 1603 974 358 142 240	11 137 26 3	1 in 4 1 in 113 1 in 374 1 in 119 0 in 142 1 in 34	475 200 90	119 24 1 	0 in 8 1 in 113 1 in 194 1 in 200 0 in 90 1 in 38	1593 356 117	7 191 18 1	1 in 3 1 in 13 1 in 88 1 in 356 0 in 117 1 in 4	
Apoplexia Paralysis Amentia	18 12 3	12	1 in 11 0 in 12 0 in 3		4 2	1 in 44 1 in 7 0 in 2		34	1 in 1 ls 1 in 7	
Mania Delirium Tremens Epilepsia	8 37 43	3	1 in 8 0 in 37 1 in 14	56 26	3 2	1 in 183 1 in 13	100 49	15 4	0 in 2 1 in 6 1 in 12	
Hydrops	23	8	1 in 3	20	4	1 in 5	40	12	1 in 3}	
Rheumatisms All other diseases	632 4556	26	1 in 158	3799	12	1 in 105 1 in 316½	665	18	1 in 665 1 in 376	
Total	13806	493	1 in 28	9228	332	1 in 274	22156	763	1 in 29	

From this record it appears that, out of an aggregate force of 25,431 British soldiers, stationed during ten, eight, and ten years respectively, between 1823 and 1836, at Calcutta, Chinsurah, and Berhampore, all in Bengal Proper, there occurred of sickness as follows:—

Fevers, re	mitte	ent, interm	ittent,	and	continued				13,596
Dysentery	and	diarrhœa	**		**				8,499
Hepatitis			**			**		**	1,354
Cholera		**				**	4.5		1,117

Total cases of four acute tropical diseases

There was a total of admissions into hospitals, on account of all diseases, of 45,170 cases, and a total of deaths from all diseases of 1,588 soldiers. It is true that these men were British soldiers, differing in habits from the higher classes of Europeans residing in the same localities—worse fed and lodged also, and that a few out of the few survivors of the Burmese war may have been admitted into the hospitals. But, making due allowance for all these circumstances, and for the bad climate of Lower Bengal comparatively, the actual sickness and mortality, not to speak of the proportionate invaliding, was enormous.

Out of that portion of the inhabitants of George Town, British Guiana (a country somewhat resembling Lower Bengal in its medical topography and climate), who were likely to resort to public hospitals—the total population of the town amounting to 20,000—there were admitted into the Colonial Hospital, from June, 1846, to June, 1847, 2,938 cases of remittent and intermittent fevers; and the annual consumption of quinine in Demerara and Essequebo averaged 3,000 ounces. But mere figures convey only a very faint impression of the calamities attending upon and resulting from tropical fevers, and other acute diseases of hot climates. To realise the full powers of such endemics we should be presented with the sequelæ of broken constitutions, and consequent invalidings, amongst British soldiers, and amongst all who are exposed to the widely-extended and unnatural influences here spoken of.

## SICKNESS AND MORTALITY OF BRITISH SOLDIERS AND SEAMEN.

The following summary deductions are made by M. Budin, from his admirable "Statistics of the Sanitary Condition and Mortality of Forces by Land and Sea, as influenced by Season, Localities, Age, Race, and National Characters." \*

These subjects are of such vast interest and importance to naval and military surgeons especially, that I hope to stand excused for here republishing the conclusions of our neighbour and worthy fellow-labourer of the French army:—

<sup>\*</sup> Vide Edinburgh Medical and Surgical Journal, No. 171—an article ably prepared by Dr. Balfour of the Grenadier Guards.

density of the population of the different quarters and streets of a large town should be seriously considered in selecting sites for barracks and hospitals.

"11. Numerous facts militate against the hypothesis of a progressive amelioration in the sanitary condition of Europea troops in warm climates in general, and particularly in tropics

regions, as an effect of length of residence.

"12. In a military point of view, the knowledge of the pathogenic march of the seasons in different parts of the globe, and of the relation of the sanitary condition of armies with the different meteorological influences, is of immense interest, and has not yet received the attention it deserves.

"13. The pathogenic influence of the seasons is in a strict dependence upon the quality of the soil, the latitude, longitude, an elevation of the places, their position in the northern and souther

hemispheres, and the nationality and race of the soldier.

"14. In all countries where the influence of age has be studied, the lowest mortality has been found to be that of soldies

from eighteen to twenty years of age.

"15. Nationality and race favour or neutralize the pathoge action of climate, so that, under precisely similar circumstatroops of different races and nations may suffer and die in proportions and of different diseases."

To the above we may add, of our own experience :-

16. That, in the first expedition to Rangoon, the mo British soldiers, in the first year of service, 1824-25, and thirty-five deaths from wounds, and 450 by disease, per strength; while, in the second year, 1825-26, the losse and by disease were about one-half of what occurred making a total mortality by wounds and by disease, 1,000 of strength, in the two years.

An expedition to Rangoon having been determ January, 1824, eattle were marched to Calcutta a stations, and slaughtered in the month of February, gree of heat so high that before the process of salticestablished, some degree of decomposition must have in the meat intended for the use of the soldiers. clared on the 5th of March, 1824. "From the missays Colonel Tullock," the troops had nothing to salt beef and biscuit of very inferior quality, and vegetables of any kind to counteract the effects of s

17. The other British expedition employed as in 1825-26, and employed in the province of Arameral Morrison, became speedily ineffective from fever of the country. Five thousand five hundred pean and native, were soon struck down with the structure of the country.

out reckoning the sick of the public establishments, and of the camp followers —soon, indeed, "every one who was not dead was in hospital."

"The Bengal infantry Sepoy suffered the most; in fact, they disappeared altogether." Of the original European force three-fourths perished, and the miserable survivors were ruined in con-

stitution-none living long thereafter.

18. It has been seen that in the first Burmese war, in Ava, owing to the neglect of every sanitary precaution, "the deaths nearly equalled the number of British soldiers originally employed on this service, and that but for the seasonable reinforcements which arrived, the whole must speedily have been annihilated."

To show, however, that Governments, like individuals, are slow to derive lessons from experience, we have cattle again marched to Calcutta, and slaughtered in the same heats of February, as stated by Major Hough, with the same consequences to the European troops employed in China, in 1840-42. It was the horrors

of Rangoon over again.

"The attempt to improve the Calcutta salted meat by fresh brine at Singhapore did not answer; and thus, in the Island of Chusan, between the 5th of July and the 22nd of October, 1840," "out of three thousand six hundred and fifty troops, only two thousand and thirty-six were fit for duty"—the 26th Cameronians and the 49th Regiment being the chief sufferers. These five regiments, says Major Hough, were reduced "to a perfect skeleton;" and, in the short space of three months, the former corps was reduced from upwards of nine hundred strong to two hundred and ninety-one.

There was here, in addition to the neglects of the Government, neglect and grievous ignorance in the officer placed in command on the Island of Chusan. "The sickness was comparatively mild amongst the officers, who had means of living on more generous diet." This, it must be confessed, reads very badly of the officer in command, who seems to have lived well, while he neglected

the comfort and welfare of the soldiers.

As a contrast to our former arrangements in Burma, in 1825-26, and in China, in 1840-42, may be cited the following:—
"The Marquis Wellesley began early in the cold season of 1800 to prepare for the expedition to Egypt, which sailed in April from Calcutta." The circumstance here recorded ought to be remembered to the honour of our great Indian statesman, as should also the admirable arrangements of the Marquis Dalhousie for the conduct of the last Burmese war.

Referring to the comparative circumstances of military service in various countries and in different climates, it may not prove un-

interesting here to quote the results of some battles, sieges, and campaigns in the East Indies, as compared to similar results in Europe. In Marlborough's battles the losses were as follow:—

	Blenheim .					5)	
1706.	Ramillies, about		,	1	22	16(	Allied armies included.
1708.	Oudenarde			1	33	36	Ameu armies merateu.
1709.	Malplaquet			1	**	5)	

At Waterloo,\* the Duke of Wellington lost in the proportion of . 1 in 6.

The Indian returns show the following ratios:-

	4							2.72 =	
1803.	Assaye		40	14				I in 3	
1804.	Dieg .							1 ,, 4	4
1817.	Mehedpore							1 ,, 6	,
	Sitabuldy							1 ,, 4	
	Korygaum							1 ,, 3	
	Maharajpore							1 ,, 6	
								- "	
1846.	Battles of th	ie S	utlej					1 ,, 5	,
1848.	Chilianwalla	h	-		-	3		1 7	1

"Here," says the *Edinburgh Review*, No. 197, "is no proof of cowardice on the part of the defeated, whose loss in every affair, except perhaps the last, greatly exceeded our own."

During the siege of Seringapatam, in 1799, it was stormed and captured "by 4,376 men, in two columns." The loss in the assault was as follows:—

	Killed.		Wounded.	Missing.
European officers	 22		45 .	-
" N.C.O. and soldiers .	181		122 .	22
Native soldiers	119		420 .	100

Making a total, killed, wounded, and missing, of 1,031 men.

Of the above officers, twenty-five were killed and wounded in the assault.

Lord Lake, with an original force of 9,000 men, augmented after the second storm by the force from Bombay, according to Major Hough, appeared before Bhurtpore in January, 1805. During four successive assaults, each increasing in desperation, Lord Lake was repulsed with the losses on each occasion, and, in the aggregate, as follows:—

First assault				456	men killed	and wounded.
Second ditto		*		573	"	n
Third ditto .				894	11	***
Fourth ditto .			*	987	**	**
Total			-	2910		

<sup>\*</sup> In the battle of Waterloo, out of a British force of 36,273, there were killed 1,417, making about 3.9 per cent.; but, including 362 men killed of the King's German Legion, the ratio rises to 4.9 per cent. The total force, British and Allied, under the Duke of Wellington, amounted to 69,686, out of which there was a total killed of 2,947, or a grand total killed of 4.2 per cent., British and Allied.

There were of officers killed, 15, and 95 wounded, making a total of officers killed and wounded, 110. "Major Thorn gives the loss in all the operations, at 3,100 men and 102 officers, killed and wounded."

During the expedition to Walcheren, in 1809, 1.67 per cent. of the entire force was killed in action, and 32.2 per cent. perished by disease, making a grand total of 34.69 per cent. in that fatal and ill-directed attempt.

Of our losses in the earlier campaigns of the French revolutionary war, we receive but the following general statements:—

In 1794, says Dr. William Fergusson, the French army in Flanders, composed principally of mere boys, many of them of five feet three or four inches in height, "kicked us before them like a foot-ball through Flanders and Holland into Germany, destroying in their course, full three-fourths of our army." The same authority, speaking of the same campaign, says that "by disease, by famine, by the rigour of the season, and by the sword, out of a host of fully 30,000 men, when the retreat from Flanders first began, scarcely 8,000 remained to witness its completion."

Figured statements more in detail would be desirable in respect of so important an operation as that described by Dr. Fergusson, but they are not now procurable; and in respect of the Continental wars of 1745 and of 1756, as well as of the first American war, it

were in vain to expect any more accurate record.

The hospital arrangement of the British army, in 1794, was very defective, and the experienced regimental surgeons were, by order of "an old, broken-down court physician of London," superseded by "graduates of the English Universities. The consequence was such as might have been foreseen.".... "Disorderly hospitals," says Dr. Fergusson, "will destroy an army faster than it can be recruited." We may infer, also, that in those days the sanitary arrangements were on a par with the hospital management; and where both these are deficient together, there can be no hope for the safety of any army.

In the Peninsular army, again, under the Duke of Wellington, taking forty-one months during which the war was carried on with the utmost vigour, an annual mortality of about 4 per cent. occurred in battle and from wounds, and 12 per cent. was from disease, "being nearly 16 per cent. of those employed;" whereas, in the first year of the first Burmese war,  $3\frac{1}{2}$  per cent. of the British troops were killed in action, and 45 per cent. perished by disease, "making a total loss of  $48\frac{1}{2}$  per cent., consequently each person employed throughout that year encountered more risk of life than in three Peninsular campaigns."

In the second year of the Burmese war, as already stated, the losses in action and by disease were about one-half of what occurred

in the first, making a total for two years of 5½ per cent. killed in action, and 67½ by disease, or a grand total for the two years of 72¾ of the European force employed under Sir Archibald Campbell. The official records exhibit a loss of 61 officers of her Majesty's army alone, killed, wounded, and died of disease,—"a very heavy loss, indeed," says Colonel Tullock, "considering that the average number of officers present did not probably exceed 150." \*

The expedition to Rangoon, during the first Burmese war, was therefore the most fatal of which we have, as yet, any record.

This result of a war with a savage race is the most remarkable circumstance of all, and it is well worthy of notice. The Burmese are held in light estimation as soldiers by the Europeans, but without reason. Warring is their natural fashion, in well constructed stockades, they inflicted on British troops almost the same proportionate amount of loss as did the legions of Napoleon, commanded by the marshals of France. Speaking of the destruction of the European force on this occasion, Colonel Tullock makes the following just observations:—"It seems essential to bring such facts as these prominently to notice, because there is no mode of estimating the severity of military service except by comparison, and it is of importance that the authorities with whom rests the ultimate reward of the soldier should have some means of knowing the risk of life and peril of constitution by which his pension has been earned."

A national mode of warfare always makes itself felt. It does not imitate, and it is not, therefore, subject to the moral inferiority, the weaknesses, inabilities, and incapacities resulting from the systematic routine of organized bodies, so cramping to the physical and moral energies of men. When the princes of Indisect aside their hordes of 80,000 to 100,000 horsemen, and imitate Europeans in forming brigades of disciplined infantry, and is stand-up fights, it was easy to foresee the speedy destruction

their armies, and the fall of their empires.

The Highlanders, in 1745, were sneered at by British office and especially by those of the cavalry, who made the least spectable figure in that remarkable contest. The British ar was then, as now, officered by gentlemen, and disciplined the Prussian system. The formation of general lines and moments in lines were then, as now, its characteristics; yet, conflict with peasants led by country gentlemen, it was quickly routed, and then destroyed. Even at Culloden, where mountain peasant was led in the usual manner, the result was same.

<sup>\*</sup> War Office Statistical Report. Presented to both Houses of Parliament by mand of her Majesty.

"The Scottish Highlanders," says Robert Jackson, "were badly provided with arms; they, notwithstanding, defeated the regular and experienced troops of the Crown, both at Prestonpans and at Falkirk; and there are grounds to believe, from the decided experiment that was made upon Barrel's regiment at Culloden, that they would have defeated them a third time, had there been union in council and accord in action."

These various circumstances of service are worthy the careful consideration of the military surgeon; as, for the conduct of his difficult and toilsome duties, it is not enough that he should have some foreknowledge of the probable amount of his sick, but he should also prepare some estimate of the probable amount of his wounded men. This tact in calculation forms, indeed, one of the requirements of an effective administrative medical officer.

Nor are the results of statistical investigations confined to military operations. On the contrary, they may be rendered subservient to, and suggestive of vast improvements in civil communities, as remarked by Colonel Tullock in the following im-

portant observations :-

"To ascertain the races of men best fitted to inhabit and develop the resources of different colonies is a most important inquiry, and one which has hitherto attracted too little attention, both in this and other countries. Had the Government of France, for instance, adverted to the absolute impossibility of any population increasing or keeping up its numbers under an annual mortality of seven per cent. (being that to which their settlers are exposed at Algiers), it would never have entered on the wild speculation of cultivating the soil of Africa by Europeans, nor have wasted a hundred millions sterling with no other result than the loss of 100,000 men, who have fallen victims to the climate of that country. In such questions, military returns, properly organized and properly digested, afford one of the most useful guides to direct the policy of the Colonial legislator; they point out the limits intended by Nature for particular races, and within which alone they can thrive and increase; they serve to indicate to the restless wanderers of our race the boundaries which neither the pursuit of wealth nor the dreams of ambition should induce them to pass; and proclaim, in forcible language, that man, like the elements, is controlled by a Power which hath said, 'Hither shalt thou come, but no further ! "

The worst enemy to the soldier has everywhere and at all times, been disease.

"Civilians, however, think that shot kills most soldiers, but Colonel Queach, a Peninsular officer of some experience, and an authority upon the subject, having served throughout the Peninsular campaigns with the old 95th Rifles, says, that 40,0 were killed in action or died of wounds—120,000 died of a great deal of which was rendered fatal by the want of medical attendance; whilst 120,000 more were, by diseased unfit for service."

It is a principle in modern war, well understood by the Marlborough, that the more rapid its course the more hum result; and Frederick of Prussia held that "all wars shown that any short and rapid; because a long war insensibly relaxes disappopulates the state, and exhausts its resources." Duri protracted operations of previous wars, the destruction "by famine and the ague" was horrible beyond description.

The losses in the great battles of Marlborough do not ap heavy when we consider the unusual nature of the obstate be overcome, the numbers opposed to him, and the old nature of the defence. This first of British commands careful, in an especial manner, of the health and comformen. It was the restless energy, the genius, the daring the impetuous spirit of Napoleon, which allowed no reghis own troops or to those of his enemies, that comple establishment of a system which set all tardy rules and pradefiance. He brought his men to make forced marches bivouac under every extremity of season—his principle be "legs win more battles than arms."

Mr. Hodge, in his "Report on the Mortality arising from Operations," states, that "Taking the whole of the casus action in the British service during the wars of the Frence lution, the proportion of those returned killed to the whole injured in naval engagements was 100 in 398, or rather moone in four; while in engagements on land it was 100 in rather less than one in five. It may perhaps be thought, a greater proportion of deaths were inflicted in the navy, tries suffered by the wounded would also be of a serious classical engagements.

but this is not necessarily the case.

"A large proportion of the wounds in naval actions are by round shot; of those in actions on land by musker injury to a vital organ from a round shot would be more prove fatal at once than one from a musket-ball; a man wounded by the latter might linger on for some time, a r

probable in the former case."

At first sight, there would appear no reason why arising from disease should be much increased in the war, as is well known to be the case in the army; but a nation of the facts proves that it was so. It is undou during the last war the risk of death to persons employ navy, from the accidental destruction of vessels alone, we five times greater than it has been since the peace.

Mr. Hodge adduces the following numbers as in excess of deaths caused by the war of twenty and a-half years, between 1793 and 1815:—

Casualties in action Destroyed and wrecked in ships From disease on board	 *	 		6,663 11,985 44,662
Total			-	63.310

"The following statement shows that the loss inflicted upon us in our contests with the navies of different nations, has been generally in proportion to the reputation of their seamen for skill and discipline, and is an additional proof of the importance of maintaining our superiority in those respects:—

			ortion of Brit illed and Wo	Number of Enemy's Ships.		
Action.	Enemy's Fleet.	To 1,000 British engaged.	To 1,000 of the enemy engaged.	To each ship taken or destroyed.	Engaged.	Taken or de- stroyed.
Cape St. Vincent	Spanish	32	19	75	25	4
Trafalgar	Franco-Spanish	100	78	94	33	18
Nile	French	112	91	82	13	11
Camperdown	Dutch	100	115	92	16	9

"The loss sustained in an action, is the price of the result obtained by it, and we must compare these elements with each other before we can arrive at a correct estimate as to either. By the above statement it is shown, that the casualties of the British in the battle off Cape St. Vincent, were only thirty-two per 1,000 engaged, while each of the enemy's ships taken or destroyed cost seventy-five men in killed and wounded. At the battle of the Nile each ship taken or destroyed cost only seven more, or eighty-two men in killed and wounded, although the proportion per 1,000 engaged was 112, or between three and four times as many as at Cape St. Vincent. From the general result, it appears, that out of 204 hostile vessels engaged, seventy-three were taken or destroyed by the British, whose total loss, in killed and wounded, was 7,349, or a fraction over 100 for each vessel lost to the enemy."

"From the various estimates that have been considered, the lowing has been drawn out, as the


General Summary of the Mortality in the Royal Navy among average force of 110,000 Men during  $20\frac{45}{100}$  Years of Hosties occurring in and between 1793 and 1815.

	Number o	f Deaths.	Estimated Number of	Excess cause War.		
Causes of Death.	Annual Ratio to 1,000 Mean Strength.	Total.	Deaths that would have occurred from the same causes during Peace,	Total.	Protein 100 De	
Casualties in action	3	6,663		6,663	10	
Drownedor destroyed in ships accidentally wrecked or burnt	6	13,621	1,636	11,985	11	
Estimated to have died from disease or ordinary accidents on board }	32	72,102	27,440	44,662	70	
Total	41	92,386	29,076	63,310	100	

"With reference to every description of armed force, it is a greatest importance to ascertain what is the average proportion men unable to attend to their duty from sickness. We have information of this kind as to the navy."

The following Table shows the Mortality in the Navy on the Indian Station for Fourteen Years ending with 184

			Annual	Ratio to Death	1,000 Meas arising	an Strength from
		Mean Strength.	Diseases.	Wounds and Injuries.	Accidental Drowning.	Unknown Causes
Peace	[1830 to 1836	1,849	15-10		2:20	
1 cacc	1837, 1838, 1839	1,883	14:34	2.48	4.42	0.53
War	1840, 1841, 1842	5,156	36-78	3.49	4.01	5.95



generations, acquire the sable aspect of the inter-tropical natives, by exchanging situations? Assuredly not. Where then is the superior pliancy of the human constitution? The truth is that the tender frame of man is incapable of sustaining that degree of exposure to the whole range of causes and effects incident to, or arising from the vicissitudes of climate, which so speedily operates a change in the structure, or at least the exterior, of unprotected animals.

But it is observed that, of those animals translated from a temperate to a torrid zone, "some die suddenly, others droop, and all degenerate." This is not to be wondered at, considering the disadvantages under which they labour. Man would not fare better, if placed in similar circumstances. Even as it is, the parallel is not far from applying; for of those Europeans who arrive on the banks of the Ganges, many fall early victims to the climate, as will be shown hereafter. That others droop, and are forced, ere many years, to seek their native air, is also well known. That the successors of all would gradually and assuredly degenerate, if they remained in the country, cannot be questioned; for already we know that the third generation of unmixed European is nowhere to be found in Bengal.

On the other hand, if we look at intertropical natives approaching our own latitudes, the picture is not more cheering. The African children, brought over by the Sierra Leone Company for education, seldom survived the third year in this country. "They bear the first winter," says Dr. Pearson, "tolerably well, but droop during the second, and the third generally proves fatal to them."

The object of these remarks, which, at first sight might seem irrelevant, will now appear. Since it is evident that Nature does not operate more powerfully in counteracting the ill effects of climate on man, than on other animals, it follows that we should not implicitly confide, as too many do, in the spontaneous efforts of the constitution, but, on the contrary, call to its aid all those artificial means of prevention and amelioration which reason may dictate, and experience confirm. In short, we should study well the climate, and mould our obsequious frames to the nature of the skies under which we sojourn.

The British youth leaves his native shores with vigorous health and buoyant spirits, for a foreign land of promise, where he is to meet with adventures, acquire fame, and realize a fortune. All the happy events, real or ideal, of his future journey through life, are painted by his ardent imagination in prominent characters, on the foreground of the scene; while reverses, sickness, disappointments, death itself, are all thrown into the shade, or, if suffered to intrude, only serve as incentives to the pursuit which has been com-

menced.

During the short space of existence to which man is doomed on earth, it is a merciful dispensation that youth anticipates no misfortune, and that, when the evil day arrives in after life, Hope comes on glittering wing, and gilds the scene even till the last ray

of our setting sun is extinguished.

That, so circumstanced, salutary precautions in matters affecting health are too often despised or neglected, need be no matter of surprise; and a quotation from the writings of a gentleman who resided more than twenty years in India, and whose talent for observation is peculiar, will place the fact beyond a doubt. "Nothing," says Captain Williamson, "can be more preposterous than the significant sneers of gentlemen on their first arrival in India; meaning thereby to ridicule, or to despise, what they consider effeminacy or luxury. Thus several may be seen walking about without chattahs" (umbrellas) "during the greatest heats. They affect to be ashamed of requiring aid, and endeavour to uphold, by such a display of indifference the great reliance placed on strength of constitution. This unhappy infatuation rarely exceeds a few days; at the end of that time, we are too often called upon to attend the funeral of the self-deluded victim."

Before proceeding to the individual disorders which prevail in hot climates, it may be well to allude briefly to some of those gradual and progressive changes in the constitution, and deviations from previous health and habits which, though predisposing and varying towards, yet fall short of actual disease. These are consequences which, on leaving their native soil, all must expect to feel more or less, and in which all are naturally and directly interested. For although a few individuals may occasionally return from even a long residence in hot climates, without having suffered any violent illness, or much deterioration of constitution, yet the great mass of Europeans will certainly experience the effects sketched out under this head, and many other consequences also which will be noticed in different parts of this work. It is, however, by the most scrupulous attention to these incipient deviations from health, by early arresting their growth, or at least retarding their progress, that we can at all expect to evade those dangerous diseases, to which they inevitably, though often imperceptibly, tend.

PERSPIRATION.—The [transition from a climate, the medium temperature of which is 52°, to one where the temperature is from 80° to 100°, or higher, might be supposed, a priori, to occasion the most serious consequences. The benevolent Author of our existence has, however, endowed man, as well as other animals, with the power not only of generating heat, and of preserving their

warmth in the coldest regions of the earth;\* but has also provided an apparatus for carrying off any superabundance of it that might accumulate where the temperature of the atmosphere ap-

proaches to or exceeds that of the body.

"We have seen," says Mr. Erasmus Wilson, "that the temperature of man varies very little in the whole extent of range between the tropics and the pole; that he can support the intense heat of the former without much elevation of his inward heat; that he can live where the mercury is a solid mass, like lead, with the most trifling depression of his vital warmth. But it must not be supposed that the constitution of the man is the same in these two opposite conditions; it is, indeed, widely different; in the one, he enjoys what may be termed a summer constitution: in the other, a winter constitution: and we all, without knowing it, have a summer constitution, to harmonize with the warmth of summer, and a winter constitution, to enable us to resist effectually the inclemency of that season.

"The regulation of the temperature of the body is only one of the purposes fulfilled by the perspiration; another, and an important one, is the removal from the system of a number of compounds noxious to animal life. It was estimated by Lavoisier and Seguin, that eight grains of perspiration were exhaled from the skin in the course of a minute, a quantity which is equivalent to thirty-three ounces in twenty-four hours. Of this quantity, a large proportion is, naturally, water; but nearly one per cent., according to Anselmino, consists of solid substances, of the latter, one hundred parts contain about twenty-three parts of salts, the remainder being organic matter. An analysis of one hundred parts of the solid matter of perspiration, according to Anselmino, gave the following

results :-

Osmazome,† combin Lactic acid salts, wi Animal matter, with Calcareous salts	th osn	nazon	ne .	,	:	 	48 29 21	parts.
Calcarcous saits		•					700	. "

"To which may be added carbonic acid gas, ammonia, and iron, and, in rare instances, copper. The peculiar odour of perspiration is due to organic matters, and its acid qualities and occasional acid smell to the lactic acid, an organic substance not far removed in composition from distilled vinegar."

† The peculiar animal principle which gives flavour to meats and part of the odour of perspiration.—A Practical Treatise on the Skin, 1845.

<sup>\*</sup> In the polar voyages the thermometer was noted as low as seventy degrees below zero, while throughout Hindustan it rises to 120° Fahrenheit, and upwards.

The talented author above quoted, adds that, "to obtain an estimate of the length of tube of the perspiratory system of the whole surface of the body, 2,800 may be taken as a fair average of the number of pores in the square inch, and 700, consequently, of the number of inches in length. Now, THE NUMBER OF SQUARE INCHES OF SURFACE IN A MAN OF ORDINARY HEIGHT AND BULK IS 2,500; THE NUMBER OF PORES THEREFORE 7,000,000, AND THE NUMBER OF INCHES OF PERSPIRATORY TUBE 1,750,000, THAT IS. 145,833 FEET, or 48,600 YARDS, OR NEARLY TWENTY-EIGHT MILES." Well may the author exclaim, "What if this drainage were obstructed." This again brings us to the practical conclusion of Moseley, that "cold is the cause of almost all the diseases of hot climates to which climate alone is accessary." Even in Europe. the summer night affects us with a chill, while the same temperature, a few months later in the winter season, would feel oppressive from its heat. It is during sleep that alternations of heat and cold most seriously affect us.

We are no sooner beneath a vertical sun than we naturally begin to experience the disagreeable sensation of unaccustomed warmth and as the temperature of the atmosphere, in the shade, now ad vances to within ten or twelve degrees of that of the blood, and in the sun actually exceeds it, the heat constantly generated in the bod cannot be so rapidly abstracted as hitherto, by the surrounding a and would thus soon accumulate so as to destroy the very fur tions of life, did not Nature immediately open the sluices of skin, and by a flow of perspiration reduce the temperature of

body to its original standard.

When we contemplate the admirable provision of Nat against what might appear to us an unforeseen event; when survey the resources and expedients which she can command all emergencies, her power of supplying every waste, and o straining every aberration of the constitution, we would be a tempted to conclude that man was calculated for immort But alas!

" Nascentes morimur, finisque ab origine pendet."

Till, at length, this wonderful machine, exhausted by it efforts at preservation, and deserted by its immaterial tenant and is resolved into its constituent elements!

But, reverting to the function of the skin: we must not co that this refrigerating process, adapted by Nature to preven serious mischief, is in itself unproductive of any detriment constitution, when carried to excess. Far otherwise is the and we may conclude for certain that those Europeans wh direct solar exposure, and who are temperate in all wa especially in the use of wine, beer, and spirits, will be in

ment of health and vigour, and capable to undergo the greatest fatigues, mental and bodily, without injury. One familiar example

will illustrate this subject.

We will suppose two gentlemen to be sitting in a room, in the East or West Indies, just before the setting in of the sea-breeze, both complaining of thirst, their skin hot, and the temperature of their bodies 100°, or two degrees above the natural standard. One of them, pursuant to the instructions of Dr. Currie, who never was in a tropical climate, applies to the negus, beer, or brandy-andwater cup, and, after a draught or two, brings out a copious perspiration, which soon reduces the temperature to 98°. It will not stop here, however, nor will the gentleman, according to the plan proposed; for, instead of putting the bulb of the thermometer under his tongue, to see if the mercury is low enough, feeling his thirst increased by the perspiration, he very naturally prefers a glass or two more of the same stimulating draught "to support the discharge," still, however, "stopping short of intoxication." Now, by these means the temperature is reduced to 97° or 961°, in which state even the slight and otherwise refreshing chill of the seabreeze, checks more or less the cuticular discharge, and paves the way for future maladies.

Let us now return to the other gentleman, who pursues a different line of conduct. Instead of the more palatable and stimulating drinks, he takes a draught of plain cold water. This is hardly swallowed before the temperature of the body loses by abstraction alone, one degree at least of its heat. But the external surface of the body immediately sympathising with the internal surface of the stomach relaxes, and a mild perspiration breaks out, which reduces the temperature to its natural standard of 98°. Farther, this simultaneous relaxation of the two surfaces completely removes the disagreeable sensation of thirst; and as the simple "antediluvian beverage" does not possess many Circean charms for modern palates, there will not be the slightest danger of its being abused in quantity, or of the perspiratory process being carried beyond its salutary limits. Nor need we apprehend its being neglected; since from the moment that the skin begins to be constricted, or morbid heat to accumulate, the sympathising stomach and fauces will not fail again to warn us, by craving the

proper remedy.

Taken, therefore, as a general rule, the advantages of the latter plan are numerous—the objections few. It possesses all the requisites of the former in procuring a just and equable reduction of temperature, without any danger of bringing it below the proper level, or of wasting the strength by the profuseness of the discharge.

During, or subsequent to violent exertion, under a powerful sun,

or in any other situation in a tropical climate, when profuse perspiration is rapidly carrying off the animal heat, and especially when fatigue or exhaustion has taken place, or is impending, then cold drink might prove dangerous, on the same principle as would cold externally applied. In persons who have been for some time in the climate, and whose digestive organs are enfeebled, some weak wine and water may not be objectionable; but such indulgence is by no means necessary in the young and vigorous, and it should be reserved for ulterior residence and more advanced periods of life. I may here mention that, during the first Burmese war, while serving as surgeon to the Governor General's Body Guard of cavalry, I found warm tea, after the most severe marches in the sun, by far the most refreshing beverage.

The habit of Napoleon under great fatigues was as just physiologically, as it evinced the sagacity of that wonderful man. Finding that excessive fatigue constricted the skin, and produced a sensation of cold, he went into a warm-bath, and on getting out he took a cup of strong coffee—a sensorial stimulant. He thus became in a few minutes fit for renewed exertions, both mental and

physical.

It is hoped that these details may not appear prolix, or unnecessarily minute, to the medical reader; but, considering that this is generally the first erroneous step which Europeans take on entering the tropies, and that the function of the skin, here under consideration, is more intimately connected with other very important ones than is commonly supposed, it has been thought proper to set them right in limine. The probability of future suffering will rarely deter the European from indulging in present gratification; but when stimulating liquids are represented, from high authority, as not only innocent but salutary, it will require some strength of fact, and of argument, to persuade young men to relinquish their use, or to check the wide-spreading evil.

SYMPATHY OF THE SKIN WITH INTERNAL ORGANS.

—In attempting to delineate the influence of tropical climates on the European constitution, although we may endeavour "to chain the events in regular array;" yet it must be confessed that Nature spurns all such artificial arrangements.

As matter of fact, however, there does exist between different parts of the body a certain connection or relation, so that when one is affected by special impressions, the other sympathises, as it were, and takes on an analogous action. Of the sympathies here referred to, none is more universally remarked than that which subsists between the external surface of the body and the internal surface of the alimentary canal. This, indeed, seems more easily comprehended than many others, since the *latter* appears to be a

continuation of the *former*, with the exception of the cuticle. I have remarked that, when we first arrive between the tropics, the perspiration and biliary secretion are both increased; and that, as we become habituated to the climate, they both *decrease*, pari passu.

In the first section an instance was given of the skin sympathising with the stomach, where the cold drink was applied to the latter organ. Had the water been applied to the exterior of the body, on the other hand, the stomach would have sympathised,

and the thirst would have been assuaged.

Mr. Erasmus Wilson justly observes, that the sensibility of the skin conveys to the mind a knowledge of the temperature of the body, thus constituting a "thermometer of vital heat, the degrees upon the scale being computed by the expressions agreeable and disagreeable, in place of the terms of the common thermometer. The value of these expressions to health is not, however, sufficiently estimated, but it is, nevertheless, certain, that a disagreeable impression of temperature in the skin is a warning note of something mischievous to health, acting either within or without the economy." The same author states the rule of health to be by food, by raiment, by exercise, and by ablution, to maintain and preserve an agreeable warmth of the skin. Everything above this is suspicious; everything below noxious and dangerous. For good or for evil, as it may be, the function of the skin is highly important.

It will be found on experience that the loss of tone in the extreme vessels of the surface, consequent on excessive or long-continued perspiration, is accompanied, or soon succeeded, by a consentaneous loss of tone in the stomach, and will fairly account for that anorexia, or diminished appetite, which we seldom fail to perceive soon after entering the tropics, or indeed during hot weather in England. Now this, though but a link in the chain of effects, seems a most wise precaution of Nature, to lower and adapt the irritable, plethoric European constitution to a burning climate, by effectually guarding against the dangerous consequences of repletion. This view of the actual subject will set in a clear light the pernicious effects of stimulating liquids, operating on an organ already debilitated, probably for salutary purposes, and goading it thereby to exertions beyond its natural power, producing a temporary plethora and excitement, with a great increase

of subsequent atony.

"An important medical law," says Mr. Erasmus Wilson, "is founded on the continuity and similarity of structure of the investing and lining membrane of the body. This law resolves itself into three expressions—namely, that disease affecting a part of a membrane is liable to spread to the whole, secondly,

that disease of the mucous membrane may spread to the skin, and vice versa; and, thirdly, that disease of a part of a mucous membrane may become translated to a part of the skin. and vice versà." A remark which every person of observation must have made, even in this country, during the summer, but particularly in equatorial regions, will further elucidate this subject. If by walking, for instance, or other bodily exercise in the heat of the sun, during the forenoon, or near the dinner hour, the perspiration be much increased, and the extreme vessels relaxed, we find, on sitting down to table, our appetites almost entirely gone, until we take a glass of wine, or other stimulating fluid, to excite the energy of the stomach. Observation and personal feeling have taught that in hot climates, perhaps during hot weather, in all climates, an hour's cool repose before dinner is highly salutary; and if, on commencing our repast, we find we cannot eat without drinking, we may be assured that it is Nature's caveat to beware of eating at all. This will be deemed hard doctrine by some, and visionary by others; but it is neither the one nor the other, and those who neglect or despise it may feel the bad consequences when it is too late to repair the error.

There are several other causes also which operate in conjunction with the above, to impair the appetite; one of which is, the absence of rest at night. After a disturbed and unrefreshing sleep, but too common in tropical climates, the whole frame languishes next day, and the stomach participates in the general relaxation. The means of obviating and managing these effects will be pointed

out in the prophylactic part of this work.

INFLUENCE OF TROPICAL HEAT ON THE BILIARY FUNCTION.—The effect of a tropical climate on the function of the liver is universally allowed to be an *increase* of the biliary secretion. This is so evident in our own country, where the summer and autumn are distinguished by diseases arising from superabundant secretion of bile, that it would be waste of time to adduce any arguments in proof.

Were this a question of mere curiosity, or theoretical specultion, I should pass it by unnoticed; but from long and attention observation, as well as from reflection, I believe that I have docovered a connection between two important functions of animal economy, which will let in some light on this subject, a lead to practical inferences of much importance. The argume and facts adduced in support of this connection will be for under the heads Hepatitis and Dysentery, and in other parts of

work; and, in the meantime, I shall merely state in a few we the result of my observation, leaving the reader to give credit, or not, as he may feel inclined.

There exists, then, between the extreme vessels of the

portæ in the liver, and the extreme vessels on the surface of the body-in other words, between the biliary secretion and perspiration—one of the strongest sympathies in the human frame, although entirely unnoticed hitherto, so far as I am acquainted, That these two functions are regularly, and to appearances equally increased, or at least influenced by one particular agent, atmospheric heat, from the cradle to the grave, from the Pole to the Equator, will be readily granted by every observer; and that this synchronous action alone, independent of any other original connection, should soon grow up into a powerful sympathy, manifesting itself when either of these functions came under the influence of other agents, is a legitimate conclusion in theory, and what I hope to prove by a fair appeal to facts. This last consideration is the great practical one; for it is of little consequence whether this sympathy was originally implanted by the hand of Nature at our first formation, or sprung up gradually in the manner alluded to, provided we know that it actually exists, and that by directing our operations towards any one of the functions in question, we can decisively influence the other. That is what I maintain, and what in a future part of the work I hope to prove by facts and cogent arguments.

I shall here offer one practical remark, resulting from this view of the subject, and which will be found deserving of every European's attention on his emigration to southern regions; namely, that as the state of the perspiratory process is a visible and pretty fair index to that of the biliary, so every precautionary means which keeps in check, or moderates, the profusion of the *former* discharge, will invariably have the same effect on the *latter*, and thus tend to obviate the inconvenience, not to say disorders, arising from redundancy of the hepatic secretion. To this rule I do not know a single exception, consequently its universal appli-

cation can never lead astray in any instance.

It is well known that if any organ be stimulated to inordinate action, one of two things must in general ensue. If the cause applied be constant, and sufficient to keep up this inordinate action for any length of time, serious injury is likely to accrue to the organ itself, even so far as structural alteration. But if the cause be only temporary, or the force not in any great degree, then an occasional torpor or exhaustion of the organ takes place, during which period its function falls short of the natural range. To give a familiar example, of which too many of us are quite competent to judge:—thus, if the stomach be goaded to immoderate exertion to-day, by a provocative variety of savoury dishes and stimulating liquors, we all know the atony which will succeed to-morrow, and how incapable it then will be of performing its accustomed office. It is the same with respect to the liver. After

great excitement by excessive heat, violent exercise in the sun, &c., a torpor succeeds, which will be more or less according to the degree of previous excitement, and the length of time during

which the stimulating causes have been applied.

When Europeans first arrive in tropical climates, the degree of torpor bears so small a proportion to that of preceding excitement in the liver, that it is scarcely noticed; particularly as the debilitated vessels of this organ continue, like the perspiratory vessels on the surface, to secrete an imperfect fluid for some time after the exciting cause has ceased; hence the increase of the biliary secretion occupies our principal attention. But these periods of torpor, however short at first, gradually and progressively increase, till at length they greatly exceed the periods of excitement; and then a deficiency of the biliary secretion becomes evident. This is not only consonant to experience, but to analogy. Thus, when a man first betakes himself to inebriety, the excitement occasioned by wine or spirits on the stomach and nervous system, far exceeds the subsequent atony, and we are astonished to see him go on for some time, apparently without suffering much detriment to his constitution. But the stage of excitement is gradually curtailed, while that of atony increases, which forces him not only to augment the dose, but to repeat it oftener, till the organ and life are destroyed!

It is remarkable that this alternation of redundancy and deficiency, or *irregular* secretion in the biliary organ, should pass unnoticed by writers on hot climates. They, one and all, represent the liver as a colossal apparatus, of the most herculean power, that goes on for years performing prodigies in the secreting way, without ever being exhausted for a moment, or falling *below* the range of ordinary action, till structural derangement, such as scirrhosity or tuberculation, incapacitates it for its duty! A very attentive observation of what passed in my own frame, and those of others, has led me to form a very different conclusion; and the foregoing statement will, I think, be found a true and a natural

representation of the case.

Here, then, we have two very opposite states of the liver and its functions;—first, inordinate action, with increased secretion—the periods gradually shortening; and, secondly, torpor of the vessels of the liver, with deficient secretion—the periods progres sively lengthening. In both cases the bile itself is vitiated. We can readily conceive how this last comes to pass, by compariso with what takes place in the stomach during and subsequent to debauch. In both instances the chyme passes through the pylor into the duodenum, in a state less fit for chylification than during a season of temperance and regularity; so, during the increase secretion, and subsequent inactivity in the liver, the bile pass

out into the intestines deteriorated in quality, as well as supera-

bundant, or deficient, in quantity.

In what this vitiation consists it is certainly not easy to say. In high degrees of it with hurried secretion, both the colour and taste are surprisingly altered, since it occasions all the shades between a deep bottle-green and jet black; possessing at one time an acidity that sets the teeth on edge; at other times, and more frequently, an acrimony that seems absolutely to corrode the stomach and fauces, as it passes off by vomiting, and, when directed downwards, can be compared to nothing more appropriate than the sensation which one would expect from boiling lead flowing through the intestines. Many a time have I experienced this, and many a time have my patients expressed themselves in similar language. The slightly disordered state of the hepatic functions, which we are now considering as primary effects of climate, and within the range of health, may be known by the following symptoms:-Irregularity in the bowels, with motions of various colours, and fætid or insipid odour; general languor of body and mind, slight nausea, especially in the mornings, when we attempt to brush our teeth; a yellowish fur about the back part of the tongue; unpleasant taste in the mouth on getting out of bed; a tinge in the eyes and complexion from absorption of bile; the urine high-coloured, and a slight irritation in passing it; the appetite impaired, and easily turned against fat or oily victuals; irritability of temper; dejection of mind; loss of flesh; disturbed sleep. These are the first effects then of increased and irregular secretion of bile, and may appear in all degrees, according as we are less or more cautious in avoiding the numerous causes that give additional force to the influence of climate.

If, for example, I use more than ordinary exercise, expose myself to the heat of the sun, or drink stimulating liquids to-day, an
increased and vitiated flow of bile takes place, and to-morrow
produces either nausea or sickness of stomach, or a diarrhœa, with
gripings and twitchings in the bowels. But a slight degree of
inaction or torpor succeeding, both in the liver and intestines,
there will probably be no alvine evacuation at all the ensuing day,
till a fresh flow of bile sets all in motion once more. These irregularities, although they may continue a long time without producing much inconvenience, especially if they be not aggravated
by excesses, yet they should never be despised, since they inevitably, though insensibly, pave the way for serious derangement
in the biliary and digestive organs, especially in hot climates,
unless counteracted by rigid temperance, and proper prophylactic

measures.

peans escape from this the most primary effect of a hot climate, for it can hardly be called a disease.

From mosquitoes, cockroaches, ants, and the numerous other tribes of depredators on our *personal* property, we have some defence by night, and in general a respite by day; but this unwelcome guest assails us at all, and particularly the most unsea-

sonable, hours.

The sensations arising from prickly heat are perfectly indescribable, being compounded of pricking, itching, tingling, and many other feelings for which there is no appropriate appellation. It is usually, but not invariably, accompanied by an eruption of vivid, red, pimples, not larger in general than a pin's head, which spread over the breast, arms, thighs, neck, and occasionally along the forehead, close to the hair. The eruption often disappears to a certain degree when we are sitting quiet, and the skin is cool; but no sooner do we use any exercise that brings out perspiration, or swallow any warm or stimulating liquid, such as tea, soup, or wine, than the pimples become elevated, so as to be distinctly

seen, and but too sensibly felt!

Prickly heat being rather a symptom than a cause of good health, its disappearance has been erroneously accused of producing much mischief; hence some of the early writers on tropical diseases, harping too much on the humoral pathology, speak very seriously of the danger of repelling, and of the advantage of "encouraging" the eruption, by taking warm liquors-as tea, coffee, wine whey, broth, and nourishing meats. Even Dr. Moseley retails the puerile and exaggerated dangers of his predecessor Hillary. "There is great danger," he says, "in repelling prickly heat; therefore cold bathing and washing the body with cold water, at the time it is out, is always to be avoided." Every naval and military surgeon, however, who has been a few months in a hot climate, must have seen hundreds, if not thousands of seamen and soldiers plunging into the water for days and weeks in succession, covered with prickly heat, yet without bad consequences ensuing.

The eruption is seldom repelled by the cold bath, which rather seems to aggravate it and the disagreeable sensations belonging to it, especially during the glow which succeeds the immersion. It certainly disappears suddenly sometimes on the accession of other diseases, but there is no reason to suppose that its disappearance occasioned them. At the same time cold bathing and repellents are not to be recommended in this eruption, even in persons of robust constitutions, recently arrived in the country, and who are in the enjoyment of good health. Where any organ happens to be weak, on the other hand, or any tendency to disease exists, the repulsion of an eruption is by all means to be avoided, and

the more so if the sufferer have been long resident in the climate.

Hair-powder, lime-juice, and a variety of external applications have been used for the removal of prickly heat, but with little or no benefit. The truth is that the only means productive of good effect in mitigating its violence, till the constitution becomes assimilated to the climate are—light clothing, temperance in eating and drinking, avoidance of all exercise in the heat of the day, open bowels; and, lastly, the use of the punkah, or large fan, during the night, as is now the ordinary practice in Bengal. The punkah is always safe; and, unlike the thorough-draught or external breeze, it removes the heated air surrounding the body, without exposing it to the dangers arising from sudden night changes in the temperature and humidity of the atmosphere.

## POSTSCRIPT.

In the foregoing sections I have adopted the order and the language of Dr. James Johnson, offering occasional additions and illustrations. I have endeavoured to place before the reader a fair abstract of the physiological doctrine—the cutaneo-hepatic sympathy of the author. This doctrine was propounded by Dr. Johnson in early life, in the East Indies, with a limited range of observation in years, and in the subjects for observation—namely, the crew of a ship of war. But with all these disadvantages it will, I think, be admitted that, practically considered, in relation to the nature, causes, prevention, and cure of disease, there is much in the theory to "lead to practical inferences of considerable importance;" for it is to the restoring of the lost balance of circulation, of the nervous functions, and of secretion, that our main efforts have been, and always must be directed in the treatment of tropical disease, whether it be Fever, Dysentery, Hepatitis, or Cholera.

For all the articles that follow, I am individually and exclusively responsible—J. R. M.

## THE PREVENTION OF DISEASE.

Descartes must have had in view the science of PREVENTION OF DISEASE, or of STATE MEDICINE, when he declared:—"S'il est possible de perfectionner l'espèce humaine, c'est dans la medicine qu'il faut en chercher les moyens." This philosopher felt that

whatever tended to improve the physical constitution of man, must promote general happiness and virtue, and thus prevent crime.

Nor did Bacon hold this greatest department of medicine in less estimation; for he exhorts the profession in his time to "exert themselves for the general good, and raise their minds above the sordid considerations of cure; not deriving their honours from the necessities of mankind, but becoming ministers of the Divine power and goodness both in prolonging and restoring the life of man." It would accord with the views of these great men that adequate public provision and arrangement should be made to enable the medical profession to render the services desired; but, in this country, it has hitherto been employed by the State individually, and chiefly in its curative or therapeutic capacity only; while it might be employed collectively, in a higher and nobler way—namely, in its preventive or prophylactic office.

It is surprising that in so civilized a state of society as ours, we should still be so far behind some of the ancient peoples. Moses imposed sanitary laws upon the Jews; the Greeks and Romans held their officers of health in high consideration. "In the Cyclopædia of Xenophon," says Dr. Farr, "the preservation of health

is declared a noble art worthy of Cyrus himself."

In the army and navy of England the medical officers have for ages past occupied themselves, spontaneously, and without prospect of reward, in establishing important principles in this great branch of medical science—the prevention of disease;—and the sister science of Medical Statistics may be said to be founded, in this country at least, on the labours of such men as Pringle, Lind, Blanc, Robert Jackson, Trotter, Hennen, and Henry Marshall.\*

<sup>\*</sup> I take leave here to state that, while serving in the East Indies, in March, 1835, I submitted to the then Governor-General of India, Sir Charles, afterwards Lord, Metcalfe, a systematic and detailed plan for requiring from all competent medical officers throughout our eastern possessions, reports on Medical Topography and Sanitary Statistics of the districts, stations, or cantonments, whether fixed or temporary, with the localities of which each officer may in the course of service be best acquainted. In the following year, 1837, I placed before the Government of Bengal a detailed Report on the Medical Topography, Climate, and Diseases of Calcutta, with suggestions for removing the defects, local and structural, which there injuriously affected the health of its inhabitants, European and native. This report also contained suggestions for the removal of the European troops from the plains to the mountain ranges throughout India. The first measure here mentioned, though it received little or no support from the medical authorities of Bengal, was ordered by a direct act of the Governor-General, Sir Charles Metcalfe, to be a standing regulation throughout the three Presidencies of India, and it has stood so ever since. The second proposal for the sanitary improvement of Calcutta has for several years been practically carried out under legislative enactments ;- the works are still in progress, under an official board, and will continue to their completion. Dr. Farr, of the Registrar General's Office, speaking of the measures above mentioned, says-" We look upon the well-conducted sanitary inquiry, commenced in Bengal, as one of the most important undertakings of

Dr. Alison, in his History of Medicine in the present century, observes that, "Next to the additions which have been made since the end of last century to our knowledge of physiology, we may place the numerous important observations by which our information as to the external causes of disease has been rendered

more extensive and more precise.

These observations and the inferences from them demand the more attention from physicians, that they necessarily involve a kind of evidence essentially different from that on which we proceed in other medical inquiries, and if we durst hope that the progress of human wisdom and virtue would bear any proportion to that of human knowledge, we might expect that the lessons to be drawn from these inquiries would prove of even greater importance to the future happiness of mankind than any which we can gather from the history and treatment of disease.

These inquiries have in some instances been prosecuted by individuals in civil life; but the opportunities of making decisive observations on some of the causes of disease which occur in the experience of medical officers of fleets and armies, who are perfectly informed of the whole circumstances of the organized bodies of men under their observation, and often see these circumstances suddenly altered, or have even the power of altering them at pleasure, are much superior to those which other practitioners enjoy; and the peculiar value of such observations has never been so well

understood as during the last war."

The unobtrusive but important services of our brethren in the navy and army, alluded to by Dr. Alison, are not written in the rolls of fame, but they will be remembered and appreciated by the philanthropist and the statesman, for ages after other and more brilliant services are forgotten. But in public rewards or emoluments our brethren must not as yet expect to share; for "in medicine alone," according to Marshall Hall, "improvement is without recompense." Referring to "the progress of human wisdom and virtue," it may reasonably be supposed that by rendering men more healthy they may become more happy, and that thus the bonds of our common humanity may be drawn the closer,

the age in India—useful to science, to England, and creditable alike to Mr. Martin, with whom it originated, and to the Government."

Lord Hardinge, referring to the same subject, says—" I am happy to bear testimony to the services you rendered to our Indian Empire in originating those valuable measures of sanitary improvement which have so materially contributed to the comfort and prosperity of an immense population. The importance of these benefits will be ever felt; and I am aware, from the interest that, as Governor-General, I took in this subject, that their inestimable results are honourably associated with the active part you took in their first introduction." Thus, although my endeavours were but coldly received by the superannuated members of the Bengal medical board, they are approved by persons the most competent to judge of their value, whether scientific or practical.—J. R. M.

so that the uncertainties, which are the same to us all, may eventually be regarded as the true and natural bond of union amongst men.

From what has been stated, it must be obvious that, for the advancement and perfectioning of the science of Prevention of Disease, we must command the co-operation of the medical profession generally, and, in particular, that of the medical officers of the army, the navy, the police, hospitals, dispensaries, lunatic asylums, parochial unions and workhouses, vaccination establishments, prisons, factories, mines, provident societies and other

institutions, and public works.

The medical profession has so invariably spurned all selfish considerations in originating and promoting objects of great sanitary utility, that I would here allude only to the obvious advantages which will result to medicine, and the increased estimation in which its professors and practitioners will be held, both by the Government and the country at large, from the realization of the projects here contemplated. "One of the first steps," says Heberden, "towards preserving the health of our fellow-creatures, is to point out the sources from which diseases are apprehended."

The Medical Topographer should investigate all the circumstances which tend to deteriorate the human race, and to lower its vigour and vitality; all that relates to the external causes of diseases, their propagation, and their prevention; all plans for improving the physical, and, through it, the moral condition of the people. He should cultivate more extensively the medical topography of the empire. The natural features and peculiarities of every locality affect materially the life and health of the inhabitants. Any general system of sanitary inquiry should, therefore, embrace information respecting the surface and elevation of the ground, the stratification and composition of the soil, the supply and quality of the water, the extent of marshes and wet ground, the progress of drainage; the nature and amount of the products of the land; the condition, increase or decrease, and prevalent diseases of the animals maintained thereon; together with periodical reports of the temperature, pressure, humidity, motion, and electricity of the atmosphere. Without a knowledge of these facts it is impossible to draw satisfactory conclusions with respect to the occurrence of epidemic diseases, and variations in the rate of mortality and reproduction.

The result of such inquiries would also, doubtless, show that agriculture is everywhere the most powerful improver of climate, and that its advancement ministers not only to the support of man in the production of his food, but in a greater degree to his health

and vigour, by purifying the air he breathes.

The practical connection, therefore, of agriculture with the nutrition and growth, the sickness and health, the birth and death, of man and of the lower animals, is inseparable from a compre-

hensive consideration of the public health.

The science of localising the population, and of constructing towns, cantonments, barracks, and hospitals, is much neglected in this country, and throughout its various intertropical possessions. It is proved over and over again, in the records of the Registrar-General that, owing to defects in the sites of towns, in the erection of buildings, in the selection of healthy soils and levels, in the supply of pure air and water, in the drainage and cleansing of streets and houses, a mortality of from a fifth to a third part greater occurs in large towns than in small communities and rural districts.

The amount of sickness occurring in cities, towns, and villages, and thinly populated districts, requires a much more careful and extended investigation than has ever yet obtained. Evidence should be obtained, on an extended scale, of the moral and physical deterioration of mankind when aggravated in low and crowded towns. "The causes," says the Registrar-General, "that destroy the lives of so many people, degrade the lives of more, and may ultimately, it is feared, have a very unfavourable effect on the energies of a large proportion of the English race. Here

is a wide field for salutary and beneficial reform.'

Amongst those whose habits and pursuits in life qualify them to be consulted in the objects of State Medicine, the following may be enumerated:—The clergy, the statesmen, colonial authorities, philanthropists, naval and military commanders, the legal profession, the medical profession, quarantine authorities, the landed proprietary and the agricultural body, magistrates, poorlaw guardians, actuaries, engineers and architects, shipowners, manufacturers of all classes, the several trades. Professors of State Medicine in foreign countries should be solicited to forward, by their information and experience, the objects in view; and the various methods which the Government of those countries have adopted to investigate and preserve the health of the people, should be ascertained and carefully considered.

It will thus appear that among the more immediate objects of

State Medicine, may be enumerated the following:-

1. To direct public attention to the increased security of life and property, which must result from a sustained and organized investigation, by competent persons, of all those agencies and circumstances, moral and physical, which deteriorate, through local or general influences, the public health.

2. To disseminate knowledge amongst all classes of the community, on the most obvious and well-ascertained causes of injury to public health, and as to the most effective and economical measures of sanitary improvement, both in town and

country.

It is believed that the union of the medical, clerical, and lay bodies, for the collection of scientific and practical information, for the consideration of sanitary measures within the United Kingdom and its foreign dependencies, and for establishing fixed and determinate principles in sanitary affairs, could not fail to enlarge the sphere of usefulness of the medical profession, for the most extended and beneficent pursuits, and to promote the moral as well as the physical improvement of the population.

"In the science of health," says Dr. Farr, "there are more exact demonstrable truths than in the science of disease; and the advantages of 'prevention' over 'cure' require no proof." I need, therefore, proceed no further to explain why I here give the priority and the preference to the subject of THE PREVENTION OF

DISEASE.

The first duty of the naval and military surgeon is, to preserve the health of the men committed to his charge—a service far higher and more useful than the cure of disease and wounds, how-

ever necessary this last function.
"Hygiene," says Cabanis, "teaches the means of preserving health, and forms an important branch of moral as well as of medical science. Ethics, being in fact but the science of life, how is it possible for this science to be complete, without a knowledge of the changes which the subject to which it is applied may experience, and without a knowledge of the means by which their changes are effected. Hygiene, therefore, and, consequently, some concise notions of anatomy and physiology, should form part of every system of education." The study of nature, he says, is, in general, a study of facts, and not of causes, a knowledge of these last not being necessary. "For studying the healthy and diseased states, for tracing the progress and development of any particular disease, we have no occasion to know the essence of life, or that of the morbific cause. Observation, experience, and reasoning are sufficient for our purposes-we require nothing more."

That the subject of Public Health has not as yet been taught in our schools is matter of much and general regret, and those of the medical profession, who have devoted their time and attention to this branch of medicine, have always done so, in the army and navy at least gratuitously, and under great disadvantages and difficulties. I can say with truth, that I first became aware of the power of external causes in the production of disease, under circumstances of general and personal suffering of the most painful character, in the marshes of Orissa and Gondwana, and afterwards in those of Rangoon. Our men were dying a miserable, a

dismal, a nameless death, by thousands. We lost more mon its gloriously than would have made the most bloody compaigns. I have gone through so many storms of febrile suffering, on so many occasions, that I often wonder how I came to survive thom. I was young, and inexperienced in the "art of preserving booth;" but I laboured hard, and, like others in the public service, I may be supposed to have acquired some knowledge essential to the welfare of the soldiers under my charge." But suffering, general and personal, was always the school in which I received my instruction.

At the moment of writing, the Legislature and Government of this country have ordained that a Board for the management of the health concerns of England shall be constituted without any person of the medical profession having a seat in its deliberations. That for this unhappy state of things the medical profession is itself principally to blame is, I think, manifest; for had "the art of preserving health" been taught in our schools of medicino, and been thus placed on its proper footing in the public estimation, no Government, in these days of free discussion, could have resisted the just claims of science. As matters stand at present, there is no remedy;—but the remedy will eventually be found, when medical science shall be represented in the councils of the nation.

It is not so much owing to the defects of our ancient sanitary laws, great as these defects confessedly were, as to the utter neglect of their provisions, that matters relating to the public health in this country have been allowed to fall so immeasurably behind the wants, the science and spirit of the age; and it is feared that popular ignorance, national indifference, prejudices, and parsimony, will continue yet a while, as of old, to neutralise better measures of sanitary regulation than even now exist! Englishmen, even of classes above the labourer, will agitate and combine for political privileges of no value when secured. They will ery for cheap bread, for holidays, for shorter time of labour, and for higher wages; but they will neither reflect upon, nor combine, for any matters affecting their sanitary improvement. Here, in our cities, as in Paris, the thing must be done for them, by an authority which they may not question. The truth I believe to be that, in respect to measures of prevention of disease, medical men have always been ready and capable to direct to good purpose, but society was not, and is not yet, anywhere possessed of sufficient knowledge to co-operate with them. I happen to possess some

<sup>\* &</sup>quot;The art of preserving health," says Dr. Farr, "is taught in no regular course of lectures in any of the great schools of medicine in the United Kingdom. Yet the classical sanitary works of Pringle, Lind, Blanc, Robert Jackson, Johnson, and Martin, have been framed from observation in the British navy and army."—Report on the Mortality of Cholera. London, 1852.

official as well as professional knowledge on this head; for I was appointed, in 1843, member of the Royal Commission, ordered by her Majesty, " for inquiring into the present state of large towns and populous districts in England and Wales, with reference to the cause of disease amongst the inhabitants." As one of the said Commissioners, I examined and reported on the sanitary condition of six of the manufacturing towns of England, and two thousand copies of my reports were printed by order of the Government. The Commission was composed of thirteen gentlemen, including two Cabinet ministers, and its deliberations and inquiries extended over two years. The final "Recommendations" contained in the Report of the Commission to her Majesty, formed the groundwork of the clauses of a Bill, subsequently passed by Parliament, for regulating the health concerns of England and Wales. The Recommendations were, with a very few exceptions, of a permissive character. To this principle I ventured on sundry occasions to object, urging that, on the score of experience, whatever was allowed to be permissive was sure to be neglected; whereas, whatever had been made imperative, like party-walls to prevent the spreading of fire, had surely been obeyed. I urged that Englishmen yield a ready obedience to Acts of Parliament, however stringent, but that to the requisitions of a sanitary Board they would render neither respect nor obedience. I think that my anticipations have been amply verified in the workings of the first unhappy "Board of Health," and that its short-comings have arisen fully as much from defective legislation as from the intrinsic defects of that illsorted Board.

I feel convinced that legislation in sanitary affairs to be effective must be IMPERATIVE, and based on the principle that no man can be allowed a right to use his property so as to cause injury to the health of his neighbour. In the "Builder" newspaper, I find at the moment of writing, the following common sense view of this question as regards one out of our many nuisances: "The total abolition of cesspools must be made compulsory; landlords will not make the necessary outlay unless compelled to do so, and the sooner it is done the better." The State does interfere, and has always interfered, in all sorts of ways, in the prevention of crime, and on many kindred subjects. Centralization may be carried too far, but so also may respect for local government. The first principles of justice, and the first sentiments of humanity, would seem to be common to all mankind; but who would trust the framing and the administering of a criminal code to local or municipal management? Self preservation requires that we should stop an evil where we can, if we must stop it somewhere. We cannot give up the right and duty of society to stop crime in the bud; nor

should we hesitate to assert the same right in respect of disease.

The simple circumstance that amongst the destitute poor we must do the cure, and that the cure is very costly and difficult, imposes on us also the duty of prevention. The polite or permissive system of legislating for the public health may do when danger is believed to be suppositious or remote; but in the presence of actual danger it will never answer any good purpose, in this or in any other country.

The right to use one's property is one thing, and the right to abuse the privilege is another thing. It is of this last that I com-

plain.

The facts and circumstances here detailed will not be considered irrelevant when the important principle involved in them is taken into account. The establishment of sound principles in sanitary affairs must precede the arrangement of details—a fact not always present to the minds of those who frame sanitary regulations.

If it be the business of Government to prevent and to punish crime—to secure the public peace—to enforce industry instead of rapine, and the settlement of disputes by appeal to reason instead of by fraud and violence; if the well being of the subject be, in short, the main object of legislation, then would it appear the special duty of the ruling power to secure the health and the life of those who, of all others, stand most in need of its protection against the invasions of individual or corporate caprice, ignorance, or stolid avarice. Here we perceive a moral and political duty of necessity. On this subject it should ever be borne in mind that where there is disease, there also will be found the seat of poverty and crime. Disease, poverty, and crime, in their worst forms, are constantly and everywhere found together. The truth then is, that misery and crime produce disease, and disease produces misery and crime, in a circle which revolves, in the same calamitous monotony, from year to year of the brief existence of the masses crowded in the worst quarters of our manufacturing cities.

Whilst men are in the lowest state of physical destitution, surrounded by filth, vermin, privation, and squalor of every conceivable kind—familiar with sickness and death, and strangers to every comfort—with the mind continually on the rack, or absorbed in striving against physical necessity—or with the animal spirits broken down by its pressure; how is it to be expected that obedience to the laws, that morals, education, or religion should find a place? How can a man whose mind is ruined even more effectually than his body—the man by whom moral degradation and physical suffering have done their worst—how can such a man be expected to give a passing thought ever to such matters? The thing is impossible. But, not to speak of those higher considerations, I should say that the benefits of the surrounding civilization

even are not for the occupants of the lanes, courts, and alleys, through which I have of late passed. \* Even amongst what are called the better classes of society we find that in the false pride of a quasi science, or of civilization, we continually neglect measures of precaution known to the rudest people; as if we thought ourselves not subject to the laws of organization which regulate the functions of the inferior animals. The nature of brute animals, indeed, we study attentively, and adapt our conduct to their constitution, whilst of our own we continue ignorant and neglectful, "If one-tenth of the persevering attention and labour bestowed to no purpose in rubbing down and currying the skins of horses," says Dr. A. Combe, "were bestowed by the human race in keeping themselves in good condition, and a little attention more paid to diet and clothing, colds, nervous diseases, and stomach complaints would cease to form so large an item in the catalogue of human miseries."

MILITARY HYGIENE, OR PRESERVING OF THE SOLDIER'S HEALTH.—M. Thiers remarks that in ordinary histories of war we see only armies completely formed and ready to enter into action; but it can scarcely be imagined what efforts it costs to bring the armed man to his post, equipped, fed, trained, lastly, cured, if he has been sick or wounded. All these difficulties are increased in proportion to the change of climate, or the distance to which the army moves from the point of departure. Most generals and governments, he says, neglect this kind of attentions, and their armies melt away visibly. Those only who practice them with perseverance and skill, find means to keep their troops numerous and well-disposed.

Of the efforts and difficulties here announced, those of the medical officer are not the least, although nowhere else, if he were fully trusted, could his services be so effective as in the preserving of the soldier's health. As it is, his disadvantages are endless. He is consulted, perhaps on active service, through ignorance or caprice, often when it is too late, but more generally not at all; and thus the soldier always suffers. In the British armies, at home and abroad, the rule is to leave everything, even of a sanitary nature, to the proper authorities, as they are called; and thus much is on all occasions found to be wanting. There is nothing so difficult to deal with, whether in public or private life, as the troubles we create for ourselves.

When a commander by sea or land is invested with authority, he

<sup>\*</sup> These two last paragraphs are transcribed from my Official Report on the Sanitary Condition of the Labouring Population in our Manufacturing Towns, as Member of the Health of Town's Commission. The observations are believed to be generally applicable to all persons and places similarly circumstanced; and they are here presented in order to show what were my sentiments on this most important subject. The principle I believe to be the same in respect to sanitary Regulations for civil and military communities; they should be compulsory for both.

always desires it to be understood that he is complete master of the situation—that he knows everything, and will do everything. But when such a man knows little, and does less, as too frequently happens, then the results are sure to be disastrous, whether we look to considerations of health, or to the completion of warlike enterprises. The truth is that Governments may confer material power on governors, admirals, and generals; but if they

do not possess moral power, it is as nothing.

Nor is ignorance and neglect of matters affecting the health of the soldier confined, as is supposed in civil life, to blundering and ill-informed lieutenants; for far otherwise is the case. When very young, and serving in one of the most pestilential countries known in India, I made a topographic examination of the localities, and reported the result to my commanding officer, suggesting at the same what I regarded as the most suitable arrangement for encamping the men, against the coming rainy season, when it was well known that a great increase of deadly fever would result. The answer was—"I'll be d——d if I do."

Now, here was no blundering lieutenant, but, on the contrary, one of the most able and well-informed field-officers I have ever known; yet, such was his treatment of a grave matter of duty, and the neglect of which, before the year was over, cost him his life.

Again, on landing at Rangoon, during the first Burmese war, I was credibly informed that the superintending surgeon of the Bengal division there had warned the officer commanding that, without fresh animal food and vegetables, the European soldiers must perish from scurvy. The answer was characteristic, and somewhat more civil than that granted to me. It was this:—
"Medical opinions are very good, sir, when they are called for."

Here also was no blundering subaltern, but a gray-headed general of reputation, and one who received much honour and

profit, after his army had perished miserably.

If it be asked how come these things, I answer that it has always been so, and that it always will continue to be so—until it is ordered otherwise.

In all our wars pestilence will everywhere be found to form their most dismal and affecting page; yet it need not have been so. In none of our most fatal expeditions—in that of Rangoon, for instance—was there any actual necessity for serving decayed rations to our men, or for exposing them through the hot and rainy seasons, to excessive fatigue, to the rank air and damp encampment—there to be destroyed, without notice and without remembrance.

Governments and communities sympathise readily enough with the soldier's triumphs, but not with his perils and sufferings. Of sympathy with his sufferings from hunger and thirst—from long and forced marches under the burning sun and the heavy dew —from the cold bivouac—from the long suspense attending field operations; with these, and with his sufferings from inscrutable pestilence, there is but little sympathy; yet they are, after all, the most ordinary as well as the hardest trials of war. For one soldier who is struck down on the field of battle, ten, sometimes twenty, are lost to their country through the unappreciated hardships here briefly recounted. England, often engaged in great wars, is, of all nations, the most slow to learn the real nature of war and of its requirements; and when peace arrives, all the experiences of former wars are forgotten in the pursuits of commerce. When we complain of inhumanity, military errors, and disastrous national failures, political adventurers and sophists are always ready to declare that "we have not suffered more than we have invariably done before on many similar occasions." Such men always have an eye to fair seemings, hollow consistencies, and present conveniences, rather than to truth.

As a familiar example of the results of mismanagement on the part of Governments and commanders, I shall here only notice the hospital arrangements during former wars, and especially the general hospital system. Pringle, writing an hundred years ago, says: "Among the chief causes of sickness and mortality in an army, the reader will little expect that I should rank the hospitals themselves, though intended for its health and preservation, and that on account of the bad air and other inconveniences attending them." Robert Jackson is even more detailed and emphatic in his condemnation of general hospitals, and he quotes the military of George the Second's time who characterised this institution as "the destroyer of the army." The truth is that the physical ills they produced by generating new diseases, tending to fatal relapses and counteracting recovery, repressing zeal in the regimental surgeons, and causing delay in early treatment, were not surpassed by their moral ills which went to destroy the best feelings of the man, and by consequence the highest qualities of the soldier. Here, if anywhere, one would have expected some deference to medical opinion, but the system continued.

The general hospital system sprung up afresh for the use of the infantry, and was employed on a large scale in Holland, in 1794-95, and the mortality was enormous, ending in "dreadful destruction;" while the cavalry, which "traversed the same fields, and lived in the same air," but which carried its own sick and treated them regimentally, had "little or no mortality in the whole course of its service." It was the same with some few corps of infantry that adopted the regimental plan of management. In the parliamentary report of the Military Commission of 1808, the general hospital system is described as "attended with the most destructive consequences to the sick soldiers, and that it has produced expenditure and waste of every kind." Again, we find "the accumulated horrors of ill-arranged hospitals," in other

words, the terrible consequences of Hospital Miasm, frequently alluded to during the Peninsular war by the historian Alison. "The military hospitals," he says, "charged sometimes with twenty thousand sick at a time, fostered contagion rather than cured disease;" while the government of England, that lavished its millions on the vain and irregular efforts, prompted by the presumptuous folly of the Spaniards, left their general no funds "to pay for hospital necessaries," for the use and comfort of his sick and wounded soldiers.

A correspondent writing from the Crimea, in December 1854. says, "All the officers here concur in describing the general hospital at Balaclava as a perfect pest house." Thus, from the days of George II, general hospitals have in no way improved on acquaintance; if, indeed, so bad an institution be capable of improvement.\*

\* "The Army at the census of 1851 consisted of 142,870 officers and men, of whom 66,424 were stationed in the United Kingdom, 2,948 on passage out or home, and 73,498 abroad in the colonies and in the East Indies. The annual mortality of men in Civil life at home of the corresponding ages is at the rate of 9 in 1,000, but the mortality of the troops at home probably exceeds 15 per 1,000, and the mortality of the troops abroad, and chiefly in the tropical climates, is such that the mortality of the whole Army is said to be at the rate of 30 in 1,000 in time of peace. At these rates, 3,290 officers and soldiers die abroad annually, of whom about 2,193 belong to England, whose names, whatever their connection with property may be, never appear on the English registers. In the time of war the deaths in the Army abroad are raised in two ways; by the augmentation of the forces, and the increased rate of mortality from wounds and from the diseases that have hitherto been incidental to warfare in the field. Thus the mean strength of the British force, officers and men, in the Peninsular, was 66,372; the deaths during the forty-one months that ended May 25th, 1814, were 35,525, of which only 9,948 happened in battle or as the consequences of wounds. 225 per 1,000 of the 61,511 men, were, on an average, upon the sick list, and their annual mortality was at the rate of 161 per 1,000.

"To the ordinary deaths of officers and soldiers abroad in 1854 must be added the excess of deaths in the war, which have been caused partly by the extension of the same epidemic of cholera that has prevailed in England, and partly by diarrhea, dysentery, and other diseases that, like cholera, are made fatal by lying on the ground, by the use of impure water, by dirt and damp, by privation, and by the substitution of salt pork, rum, and biscuits, for the fresh meat, vegetables, bread, fruit, ale stout, or wine, that

officers and men, like the rest of the people, live on at home.
"Sixty-one thousand of the deaths in England during the year 1854 are referable to the imperfect operation of the sanitary organization of our towns. And the same causes exaggerated certainly, with the absence of the comforts and necessaries that are supplied

at home, has led to the deplorable destruction of life in the Crimea.

"The deaths in an average year among 54,000 men in the town and country population of England at the same ages as the men in the Army is 486, or nearly 41 monthly, and about 972 are constantly sick. All the deaths and sickness in excess of these numbers, except the deaths and wounds from battle, are, like the excess of deaths and sickness in our towns, referable to conditions that, in the present state of engineering, chemical and medical science, may be removed to a considerable extent in ordinary climates, even in the field and in the presence of an enemy; for the art of preserving life has, since the Peninsular campaigns, made as much progress as the manufacture of arms, and if skilfully applied, our Army will never again endure the mortality from disease that so much impaired its efficiency once in the Peninsular, and again, after the lapse of more than forty years, in the Crimea."-Dr. William Farr, Reports of the Registrar-General.

When general hospitals must of military necessity be established, even for a short time, their inmates should consist only of such sick and wounded men as cannot be removed to better quarters, and crowding ought above all things to be carefully avoided; the position, construction, and arrangement being, at the same time, of the most approved kind; the charge and control being conferred on medical officers of rank and experience, in order that the various abuses, moral, medical, and financial, inherent in the General Hospital system, may be guarded against as far as possible. It ought also to be made an understood matter in the establishment of all general hospitals, that they should be but temporary, and that they should be abolished at the earliest possible moment.

When the scene of war is in Europe, the sick and wounded should invariably be sent home. With our command of shipping this may always be done, and there is no just reason for neglecting this duty to the soldier. It is never neglected in the instance of the officer.

As illustrative of this subject I may mention that, previously to enlarging the accommodations of the Smallpox Hospital in London, "Erysipelas, Typhus, Malignant Cynanchy," and other formidable diseases were common; but "by spreading the same number of cases over double the extent of surface," these terrible consequences of crowding have been altogether avoided.

Respecting the ill-regulated general hospitals for seamen and soldiers, the ill-situated and ill-ventilated hospitals of old in our great cities, one might reasonably have asked with Ponteau, "are hospitals, then, more pernicious than useful to society?" So much for one item only of neglect in matters affecting the preservation of the soldier's health. Those who desire to see further into the health history of the British Army will find other numerous sources of disease and death in the land forces, narrated in the famous speech of Earl Grey, delivered in the House of Lords on the 8th of April, 1854.

This noble address went far to inform and confirm public opinion; and public opinion in this country need rarely be mistrusted, if it is but formed on sound knowledge, and pronounced with due care; but the subject is a difficult one.

"The administration of an army," says Audouin, "is sometimes called a trade, but those who understand the subject know that it is a science, and one of so complicated a character that the study of one of its branches might occupy a long life. To comprehend even the elements of the administration of an army, it is necessary to know its origin, progress, and rules; the various means employed to raise soldiers, organize, arm, equip, pay, and put them in motion; to subsist them in health and in sickness; to command them in the field of battle; to profit by their success:

to remedy their failures; to reward and to punish them; and to preserve an account of their fortunate and unfortunate operations." Histoire de l'Administration de la Guerre.

Referring more especially to the preserving of the soldier's health, I would quote the sensible observations of Hennen, who, speaking of the prophylactic value of medicines, says that he does not question their proper use in the cases of reflecting individuals; "but I don't hesitate to say, first-that they cannot be generally applicable to a whole corps or garrison. Secondly, that although the soldier may submit in passive obedience, he will invariably make himself amends, as he supposes, for the restriction by subsequent excess of one kind or other; and thirdly, although military officers are sufficiently enamoured of any favourite theory originating with themselves, they view the proposals of medical officers but too often with a jealous eye, especially when the advantages to be derived from them are merely prospective. We possess the power, by means of the established medical inspections, to meet the approaching disease as early as possible, but I question the prudence, in a military view, of anticipating it before its arrival by a general administration of medicine throughout the garrison; because nearly thirty years' experience has convinced me that no power on earth will reconcile British soldiers to taking physic en masse, when they are not sick, nor will they ever view the man who orders it in any other light than that of a speculative experimentalist.

The true preventives to disease are shelter from the heat of the day, and from the dews and cold of night, avoiding the neighbourhood of marshes and other unhealthy spots in military exercises, mounting guards at such an hour that the least possible number of fatigue-parties may be employed in conveying dinners, &c.; timing duties in such a way that the men may enjoy their natural sleep, regulating the messes so that the soldier shall always have a due proportion of vegetables, and especially a comfortable breakfast before going on morning duties; furnishing every man with flannel waistcoats, or cotton shirts,\* enforcing personal cleanliness by frequent bathing, and by daily washing the feet, &c.; but, above all, regulating the canteen, so that access can be had to liquor only in the evening, and then taking every precaution that the bad spirits and sour wine of the country be rigidly withheld. We may refine as much as we choose, and we may modify our plans according to circumstances with critical precision, but these are the basis on which health is founded, so far as the soldier is individually concerned."

<sup>\*</sup> During the hot and rainy season in Bengal the shirt should be changed after exercise, and friction with a dry cloth used at the time of changing. The bedding should also be daily exposed to the sun.

Of all the causes which tend to the premature destruction of the British soldier in our intertropical possessions, none are so powerful as the neglect in selecting proper localities for camps and cantonments, together with the neglect of suitable structural arrangements in his barracks and hospitals. As compared to these neglects the other ills, although great, sink into comparative insignificance.

The dieting of the soldier is another question which greatly concerns his comfort, health, and efficiency, but which has not received from authority a tithe of the attention due to it; and with the exception that breakfast has been allowed him, I understand that the coarse sameness of his food, and the same inferior cooking of it, remain as in 1793. This subject demands the attentive consideration of competent persons—a complete revision and reform—so that the soldier's food should be ordered to suit his wants, and changed so as to be adapted to the various climates in which he serves. He should also be instructed to dress his food in a palatable wholesome manner; and these things can be done with little cost or trouble.

The same indifference of authority is still, after ages of heavypriced experience, manifested in the ordering of the soldier's clothing and head dress throughout our intertropical colonies, no adequate arrangement being made for climate, for peace or war service, for the night and day duties.

All the best officers concur in the recommendation of Robert Jackson, to permit the soldier to do for himself whatever he can do without injury to his health and discipline; and I feel quite assured that as regards his food, as one example out of several, there is much truth in the suggestion.

As already stated, I do not believe that there is any one cause so effective to the destruction of the British soldier, all over the world, as that of the neglect of fitting positions and other arrangements for the temporary and the permanent encampment of our troops.

In Holland, in 1794; at Walcheren, in 1809-10; on the Guadiana, in 1810; in Varna, in 1854, the same errors were committed. To station our soldiers in the plains, at the embouchures of rivers and low harbours in our Eastern and Western intertropical possessions, is to place them where they are sure to die without any necessity. There never can, indeed, exist under any circumstances, any reasons of military or political necessity for destroying an army; it is an absurd subterfuge and misrepresentation, under which no ignorant or negligent commander should be allowed to shelter himself for a moment. If a disgraceful capitulation is punishable with death, surely some punishment should attach to the wanton sacrifice of an army from causes easy of prevention.

So far back as 1835, I urged on the supreme government of India, as part of a measure sanctioned and carried at my sole

instance and recommendation by Sir Charles Metcalfe, then Governor General of India, for collecting Sanitary reports from all districts, stations, and cantonments in the three residencies of

India, the following preliminary clause, namely, that

"The topographical reports when forwarded to each Presidency by the superintending surgeons, shall then be collated by a committee of three medical officers, nominated by the medical boards, and that such as are approved shall be printed and formed into a memoir, a copy of which shall be furnished to all staff surgeons and officers of the Quarter-Master General's department."

By this clause I intended that information on the all-important question of medical topography should be imparted to the officers of the Quarter-Master General's department;—information which

they could not acquire of themselves.

In furtherance of this object I recommend that a medical topographer, an important officer of health, an officer of trust and of rank, be permanently attached to the Quarter-Master General's department, in the British as well as in the Indian army. In time of peace he should inspect and report on the sites of camps, cantonments, and stations, and on the condition of barracks, and hospitals, and transport ships.

In time of war, in addition to the above duties, he should accompany the Quarter-Master General in the field; and, where military reasons do not control, his opinion should be obtained on the sites of encampment, and upon every matter relating to the prevention of disease; all the suggestions of matured sanitary experience being put in requisition for the protection of the soldiers'

health.

This great conservative branch of medicine has not even yet been called into sufficient exercise in our fleets and armies; yet it is on the perfection of its civil establishments that the safety and

efficiency of our sea and land forces entirely depend.

Had such an officer existed in 1809 the soldiers employed on the expedition to Walcheren, would not have been permitted to encamp in a pestilential site, involving as this did the destruction of the army, and the sacrifice of a million sterling, per annum, in perpetuity; and if Wellington had had such an adviser he would not perhaps have decimated his army on the Guadiana.

But it is in our intertropical possessions that the chronic waste and destruction of men and money takes place; and could we count the hundreds of thousands of British soldiers, and the millions sterling there lost unnecessarily, even the humanity of

commanders and of governments would be shocked.

With a little attention to matters very easy of being understood, and quite as easy of being executed, we may obviate, for the future at least, those dark passages which tarnish the histories of our colonial administration, and of all our wars.

In unhealthy climates especially, and during the prevalence of epidemics in all climates, the medical inspection of the troops should be regular and frequent. This rule tends greatly to the preservation of health, for it is a standing order of all well regulated troops that every individual soldier be fit for duty in the full sense of the word, or on the sick-list under regimen and medical treatment.

It is a remarkable fact, and one often observed, that in malarious countries especially, and where vicissitudes are great, most diseases have their origin in night exposure. "During natural sleep," says Dr. W. F. Edwards, "there is a diminution in the power of producing heat, and this explains why a damp cold air, or a dry and piercing air, which is borne without inconvenience while the individual is awake, even without the aid of exercise, may be hurtful during sleep." Speaking of the yellow fever, William Fergusson says, "It is my belief that malaria can only prevail upon the body during the passive state of sleep; in fact, that to sleep is the danger." It may be remarked, in addition, that the effect of exposure to cold during sleep must necessarily vary according to the power of producing heat. As a means ofguarding the system against the effects of atmospheric vicissitudes I know of none so influential as the cold bath, provided always that the subject be healthy and of temperate habits.

The unnecessary and vexatious restrictions on the soldier's freedom of action, such as we constantly observe in garrisons and cantonments in India, are irksome to his mind and injurious to his health. He should be allowed to do for himself whatever he can do without prejudice to health or discipline; and this ought to be made matter of standing regulation, not depending, as now, on the mere will or the uncertain notions of individual commanding officers. "It is easy to conceive," says Pringle, "that the prevention of disease cannot depend on the use of medicines, nor upon anything which a soldier shall have in his power to neglect, but upon such orders as he himself shall not think unreasonable,

and such as he must obey."

The flannel is still a debatable question with military surgeons; and William Fergusson considers it necessary only for the bivouac. As an article of military wear he considers it one of which the healthy, hardy soldier (and there ought to be none others in the army) can never stand in need. The recommendations of this distinguished officer are worthy of attention on another important point—the night-covering and accommodation of the soldier. He urges that, in every part of the British empire the men should sleep in separate hammocks, which are cool, soft, and elastic, and can be washed like a garment, while they cannot be used without raising the body off the ground, or the hard boards, and they require no aid from flocks or straw to make the

occupant comfortable, an additional blanket in cold climates being all that is necessary. The hammock-railings have the great advantage of preserving freedom of ventilation, even during the night, and of being altogether out of the way during the day. Wherever a couple of stakes can be driven into the ground it will serve all the above-mentioned purposes in the bivouac; and the creation of hammock-railings, consisting of uprights with cross-

posts and hooks, would be as simple as economical.

The preservation of the British soldier's health in our various colonies is a matter of paramount importance as regards our political condition, little as the subject has hitherto engaged the attention of our legislators. "No circumstance," says a popular reviewer, "has so checked the progress of English rule as the maladies peculiar to tropical climates. And it is no exaggeration to affirm, that any means by which this intense mortality could be diminished would more effectually secure our dominion than the most brilliant victories we have ever achieved. Not only does the expense involved in conveying troops to fill the vacancies in the decimated legions fall heavily on the parent country, but another result, more lamentable than any pecuniary loss, invariably follows. A prestige of insalubrity hangs like a cloud over our colonial possessions. The soldier and the emigrant leave home with a melancholy foreboding at the probable speedy termination of their career. Hence the colonies unhappily become the resort of the desperate and the reckless. The worst features of English society are there perpetrated with few of its excellencies. This affects the condition of our army to an immense extent."

The rulers of France have not considered this question in its just view, or they never could have conceived it possible to colonise the Algerian provinces under an annual mortality of seven per cent. The result, years ago, has been the sacrifice of a hundred millions sterling, and the loss of a hundred thousand men.

Colonel Tulloch justly observes that military returns, properly organised and properly digested, afford one of the most useful guides to direct the policy of the colonial legislator. Nor is the value of well-ordered numerical returns confined to military questions.

"Whether we wish to appreciate the value of symptoms," says Louis, "to know the progress and duration of diseases, to assign their degree of gravity, their relative frequency, the influence of medical constitutions upon their development, to enlighten ourselves as to the value of therapeutical agents, or the causes of diseases, it is indispensable to count."

Cormenin, a celebrated French writer, says truly, that "details govern the world;" and the aphorism is especially true of armies. It is different with civil communities, in which the defects of law and of details, are compensated by the moral attitude and conduct

of the people. Mr. Macaulay expresses this fact when he declares that the Scotch would be well governed without any laws, or under the worst laws, while the Irish would be ill governed under the best laws. But it is not so with an army—an organised and artificial body, in which good regulations and "details" are everything.

The details here presented by authority of the French War Minister, for the service of the army to be employed on the Danube, may at first sight appear a kind of truism; but everything must be made plain to the soldier, and everything ordered

in detail.

"SUMMARY OF THE PRECAUTIONS TO BE TAKEN FOR THE PRE-SERVATION OF THE HEALTH OF THE TROOPS IN THE EAST,

"The Council of Health has been charged by the Minister of War to examine the precautions most proper for maintaining the health of the troops in the country to which they are called to make war. This Council has drawn up detailed instructions, from which the following particulars are extracted as necessary to be observed, with the greatest care:—

"1. It is necessary to be always so clothed as to be proof against the sudden chills to which one is liable, at all seasons, from the abrupt changes of temperature which very frequently

happen in nearly all parts of this country.

"2. In summer the best protection against sun-strokes, which are often very dangerous, is never to leave shelter without having the head covered.

"3, Cleanliness of persons, clothes, and dwellings is imperi-

ously required by the nature of the climate.

"4. Whenever practicable, the face, and particularly the eyes, should be frequently washed daily, after exposure to dust.

"5. The feet should not be washed with cold water, especially

when heated after a march.

"6. The greatest care is needed for protection against the freshness of the nights, even when the heat is extreme; it is dangerous to remain clad only with the shirt during the night. At the biyouac, and in the tent the soldiers should be very care-

fully covered.

"7. When the camp is pitched near a marsh, a tank, pools of stagnant water, or a valley, the chief openings of the barracks or tents should be in the opposite direction. In these bivouacs every possible means should be employed to counteract the invariably noxious vapours exhaled by such *foci* of infection. At night the openings of the barracks, with the exception of those indispensable for ventilation, should be closed.

" 8. It is wrong to sleep in immediate contact with the ground;

perfectly dry substances, not easily permeated by moisture should be interposed. For this purpose fresh branches or vegetable matters should never be used.

"9. Water, drunk in large quantity, is always injurious. If, after a fatiguing march, a stream of water is met with, thirst must be sparingly satisfied, and the water reserved for subsequent use.

"10. When only a small quantity of water is procurable, instead of swallowing it, the mouth should be gargled as long as

possible, and the water rejected as soon as it is warm.

"11. When reduced to the necessity of drinking stagnant water, it should, by way of precaution, be strained through a cloth, to separate leeches, imperceptible from their smallness, and which it is very dangerous to swallow.

"12. A mixture of wine and water, brandy and water, or infusion of coffee and water, is always an excellent drink, taken in moderation. It should be mixed at the time of use, and not prepared beforehand, as it in that case becomes heated, changes, and no longer fulfils its purpose.

"13. When salted meat and fish are substituted for fresh meat, they should be soaked before cooking, and, when practicable,

mixed with a certain amount of vegetables.

- "14. Condiments in small quantity are good seasoning; in excess they irritate the stomach, and render thirst more difficult to bear.
- "15. Saffron increases the digestibility of rice and flour; it is particularly useful with maize.

"16. Food should always be taken before a march.

- "17. Before and after guard-mounting, it is very useful to drink a moderate quantity of warm water, with a little brandy or infusion of coffee in it. Night guards should always be well clothed.
- "18. Sickness should at once be reported to the medical officer.
- "19. These recommendations shall be published in general orders. Every commanding officer will take care that they are read to the troops, once a week at least, and that the officers superintend and direct their execution."—British and Foreign Medico-Chirurgical Review, April, 1855, page 424.

The tente-abri of the French—the suggestion of necessity in their African campaigns—is the most useful article yet introduced amongst soldiers. It is three feet high, four and a half long, and is composed of three pieces of canvas, attached by buttons and strings. Two sticks placed at each extremity serve to support it, and it is fixed by pins to the earth. Three minutes suffice to set it up, and two to take it down. Each tent accommodates three men; each of the three carries a portion of the canvas, and they

divide the sticks and pins among them, and so they march. The first thing they do on arrival at their ground is to fix the tent, and to dig a trench to carry off the water in case of rain. At night they sleep wrapped in blankets, and with their knapsacs for pillows. They are thus preserved from damp and rain.

The tente-abri thus proves a protection for health and a guarantee for discipline; it preserves an effective force and

assures the duration of the army.

In camps and fleets, the officers in command should by every means promote and encourage all kinds of innocent and salutary games and exercises, music, dancing, &c. The monotony of naval and military life, and the varying and sometimes disappointing circumstances of active service, lead to mental despondency and consequent ill health. The fact has often been exemplified; so much so, that the medicina mentis should ever hold a high

place in naval and military prophylaxis.

The late Dr. James Johnson relates, that "His Majesty's ship Russell, 74, sailed from Madras on the 22nd of October, 1806, and arrived at Batavia on the 27th of November; the crew healthy, and their minds highly elated with the sanguine expectations of surprising the Dutch squadron there. Such, however, was their sudden disappointment, and concomitant mental depression on missing the object of their hopes, that they began immediately to fall ill, ten, twelve, or fourteen per day, till nearly two hundred men were laid up with scurvy, scorbutic fluxes, and hepatic complaints! Of these, upwards of thirty died before they got back to Bombay, and more than fifty were sent to the hospital there. The Albion did not fare better. The Powerful fared worse: so that in these three ships only, in the short space of a few months, full an hundred men died on board, and double that number were sent to hospitals, many of whom fell victims to the disease specified and aggravated, and in a great measure engendered, by mental despondency."

With equal truth Sir Gilbert Blane, referring to the converse influences of high spirits, says:—" When the mind is interested and agitated by warm and generous affections, the body forgets its wants and feelings, and is capable of a degree of labour and exertion which it could not undergo in cold blood. The quantity of muscular action expended in fighting a great gun for a single hour, is perhaps greater than what is employed for a whole day in ordinary labour; and though performed in the midst of heat and smoke, and with little bodily refreshment, yet the powers of nature are not exhausted or overstrained—even the smart of wounds is not felt!" . . . "It is stated that, when the fleet under Admiral Matthews, in the year 1744, was off Toulon, in the daily expectation of engaging the combined fleets of France and Spain, there was a general suspension of the progress of sickness, par-

ticularly of the scurvy, from the influence of that generous flow of spirits with which the prospect of battle inspires British seamen. But if the mere prospect and ardent expectation of battle, without any happy result, could have such a sensible result, what must be the effect of the elevation of mind created by the exultation of victory, a victory in which the naval glory of our country was revived and retrieved, after a series of misfortunes and disgraces, which had well nigh extinguished the national pride and spirit in every department of the service. The plain and honest, though unthinking seaman, is not less affected by this than the more enlightened lover of his country. Even the invalids at the hospital manifested their joy by hoisting shreds of coloured cloth on their crutches."

In accounting for the smallness of the mortality amongst the soldiers in the first month of service at Walcheren, the distinguished author here quoted says, that "an excited tone of mind, as well as youth and robust health, had a share in keeping down the mortality at this period." And so it was with the health of Hill's division when it surprised that of Gerard at Aroyo de Molinos, in Spain. Dr. Luscombe states, that during the week in which this surprise was being executed, the rain was heavy and almost incessant, and the men passed two nights in bivouac without fires; yet the sick in Hill's division was less during that and the subsequent week than in any equal period during that year. Dr. Luscombe, in almost the very words of Blane, assigns as the reason of the immunity of the soldiers from the effects of fatigue and exposure to cold and wet, that they were under "exercise and mental excitement."

Finally-it cannot be too often repeated, that on the perfection of the so-called civil establishments-the Medical and the Commissariat departments-must depend, not only the efficiency, but the very existence of our fleets and armies. I say "so-called" because, in respect to the medical corps at least, it is a complete misnomer; for neither by sea or on land is there nor can there be any absolute protection, for the surgeon, during the hour of battle, from the fire of an enemy. The recent calamitous losses in the Crimea have demonstrated another fact, viz., that so long as the departments above-named are dependent and subordinate, and under the management and control of officers holding but an inferior rank, consideration, and power, so long will they remain insufficient to their great purposes; but let them once be ruled and directed by responsible officers of a rank and station, adequate to command an instant attention to their respective wants-let such officers be largely and immediately responsible-and then, but not till then, will the two most important establishments of our fleets and armies rise so as to be equal to all requirements, whether during peace or in war.

TROPICAL HYGIENE. — Hints for the Preservation of Health in all Hot Climates.

Certain popular means of prevention of disease present themselves to the common sense of most nations to a certain degree, if they would but choose to reflect on their importance, and act upon them. But it is with communities as with individuals; hygienic rules are better known than regarded. They are like the vital points in religion and morals;—all men agree in them, yet how easily are they forgotten! In order to think seriously on matters relating to health most men require to suffer from disease, the lessons derived from such experience being longest remembered.

But there is one circumstance which ought to be impressed everywhere on individuals and communities; and that is, that however useful medicine may be in moderate and judiciously administered doses, under occasional circumstances of change of climate or season, or during the prevalence of certain epidemics, it is yet more on the proper selection of localities, the avoidance of day and night exposure, care in diet, clothing, exercise, &c.; in short, on the adoption of all those well-known measures of avoidance, whether affecting individual habit of life, or those more general predisposing causes of disease now so well understood, that the prevention of disease depends, and not on a system of self-quackery with calomel and other mercurial preparations, such as many persons pursue in England, and in India too, to their great injury, for the removal of what they call "biliousness."

Many is the robust habit I have seen destroyed by this senseless custom; and I have known several lives lost, and others put in jeopardy by the use of saline purgatives during seasons of cholera.

Another source of destruction to health I must here mention, as it has come frequently under my notice, both in India and in England; -I mean the long-continued use of aperient medicines containing the mercurial preparations. Patients frequently obtain from their physicians aperient pills, for instance, containing some blue pill, or calomel. They may have been given with a particular view, or for a special occasion only; but it often happens that the patient continues for months, or even for years, that which was intended to be used for days or weeks. The results are very lamentable. I have seen persons in a state of nervous irritability bordering on insanity, from this cause, with sub-acute inflammation of the mucous digestive membrane, and chronic ptyalism—all resulting from long-continued and unconscious use of mercury. A field-officer used blue pill and colocynth for two years and a half, and an American gentleman took the same preparations, with ipecacuanha, during a voyage from Madras to America, and back to Calcutta. It is quite needless to detail how utterly ruined was the health of both.

"Ingenuity," says Dr. Copland, "cannot possibly devise a more successful method of converting a healthy person into a confirmed invalid, of destroying many of the comforts of existence, and of occasioning hypochondriasis and melancholy, than the practice of prescribing large doses of calomel on every trifling occasion, or when the bowels require gentle assistance; or because the patient erroneously supposes himself to be bilious, or is told so by those who should know better." The same distinguished writer ascribes the lapse of occasional indigestion into confirmed stricture of the rectum, and of hæmorrhoidal affections into fistulæ, to the frequent and injudicious use of calomel for the removal of mere occasional derangements of health. The unfortunate word "bilious," as applied to slight disorders, is the scape-goat of the ignorant.

Dr. Paris assures us that, "if the truth were told, a large portion of dyspeptics seek the advice of a physician, not so much for the better adjustment and regulation of their diet as for the means by which they may counteract the ill effects of their indulgences—hence the popularity of those "antibilious remedies, which promise to take the sting out of their excesses, and enable the unhappy dupes to fondle and play with vice as with a charmed

serpent."

The following is an abstract of the excellent special rules prescribed by Dr. James Johnson, regarding dress, food, drink, exercise, sleep, bathing, and the conduct and regulation of the passions. For what has been withdrawn, altered, or added, I alone am responsible.—J. R. M.

There is no situation where the stranger European may not obviate, in a great measure, the first and most dangerous effects of a tropical climate, by a strict observance of two fundamental rules, TEMPERANCE AND COOLNESS. The latter indeed includes the former; and, simple as it may appear, it is in reality, the grand principle of intertropical hygiene, so well understood and practiced by all classes and descriptions of natives of the East, and which must ever be kept in view, and regulate our measures for the preservation of health.

Common sense, independently of all observation and reasoning on the subject, might, a priori, come to this conclusion. From heat spring all those effects which originally predispose to the reception or operation of other morbific causes; and how can we obviate those effects of heat, but by calling in the aid of its antagonist—cold. To the sudden application of the latter, after the

former has effected its baneful influence on the human frame, I have traced most of those diseases attributable to climate: nothing therefore, can be more reasonable than that our great object should be to moderate by all possible means this heat, and to habituate ourselves, from the beginning, to the impressions of cold; so that we may thus bid defiance to the alternations or vicissitudes of both those powerful agents.

This is, in truth, the grand secret of counteracting the influence of tropical climates on European constitutions; and its practical application to the common purposes of life, as well as to particular exigencies, shall now be rendered easy and intelligible.

DRESS.—When Europeans enter the tropics they must bid adieu to the luxury of linen—if what is uncomfortable, and indeed unsafe in these climates, can be styled a luxury. The natives of the country, from the lowliest to the highest, wear none but cotton clothes, and those of them who can afford luxuries wear them in

the largest quantities.

The waterman of Bengal may be taken as an example of a man who works very hard, on wages of from two to fourpence per day. His clothing, like his diet, is scanty and precarious in the extreme; the former amounting only to a narrow piece of cloth passed between the thighs, and fastened before and behind to a piece of stout packthread that encircles the waist. Thus unprotected, he exposes his skin to the action of an intense tropical sun-to a deluge of rain, to the dews of the night, and to a cold piercing N. E. monsoon-with equal indifference; and with perspiration issuing at every pore, he darts overboard when necessary, and wades through puddles and marshes-this moment under water, and the next in the open air, with rapid evaporation from the whole body. It is true that he, in some degree, supplies the defect of clothing by the assiduous and regular use of oil friction to the entire surface of the body; but, independently of all and every habit, Nature has, by race and from birth, done much to secure the hardworking boatman, by forming both the colour and the texture of his skin, in such a manner that the extreme vessels of the surface are neither violently stimulated by the heat, nor easily struck torpid by sudden transitions to cold. Certain it is also, that the action of perspiratory vessels is different from those of Europeans, the fluid exuded in the Native being more oily and tenacious than the sweat of the former.

The dress of the shepherds again, throughout Bengal, who are much exposed to all weathers, consists of a blanket gathered in at one end, so as to rest on the head, the rest hanging all round like a cloak. This answers the triple purpose of protection from the hottest sun—of a tent in the rainy season to throw off the wet—and of a coat in the cold season to defend the body from the

piereing cold. Thus our ridicule of the Portuguese and Spaniards for wearing their long black cloaks in summer, to keep them cool, and in winter, to keep them warm, is founded on prejudice rather than on correct observation.

But if we look beyond the hardy labouring classes of natives of Hindustan, we observe both Hindu and Mahomedan guarding most cautiously against solar heat, as well as cold. The ample turban and khummerbund meet our eye at every step; the former to defend the head from the direct rays of a powerful sun—the latter for the purpose of preserving the important viscera of the abdomen from the deleterious impressions of cold. The khummerbund is certainly a most valuable part of the dress, and one that is extensively imitated throughout India, by Europeans, in the form of a cotton or flannel waistband, worn generally next the skin.

The cotton dress, from its slowness of conducting heat, is admirably adapted for the tropics. It must be recollected that the temperature of the atmosphere, sub dio, in the hot seasons, exceeds that of the blood by many degrees, and even in the shade it too often equals, or rises above the heat of the body's surface, which is always, during health, some degrees below 97°.

Here then we have a covering which is *cooler* than linen, inasmuch as it conducts more slowly the *excess* of external heat to our bodies; but this, though a great advantage, is not the only one.

When a vicissitude takes place, and the atmospheric temperature sinks suddenly far below that of the body, the cotton covering, faithful to its trust, abstracts more slowly the heat from it, and thus preserves to the wearer a more and more steady equilibrium. To all these advantages must be added the facility with which the cotton absorbs perspiration. While linen so circumstanced would feel wet and cold, under a breeze, and even occasion a shiver, the cotton dress, as stated, would maintain an equable warmth.

That woollen and cotton dresses should be warmer than linen, in low temperatures, will be readily granted; but that they should be cooler in high temperatures, will, perhaps, be doubted. But let two beds be placed in the same room during the day, when the thermometer stands at 90°; and let one be covered with a pair of blankets, the other with a pair of linen sheets. On removing both coverings in the evening, the bed on which were placed the blankets will be found cool, the other warm: the linen transmitted the heat of the surrounding air to all the parts beneath it, while the woollen covering, as a non-conductor, prevented and obstructed the transmission of heat from without.

From this view of the subject flannel might be supposed superior to cotton, and indeed at certain seasons, or in particular places where the mercury often takes a wide range in a very short time, the former is a safer covering than the latter, and is adopted by

many experienced and seasoned Europeans. But, in general, flannel is inconvenient for three reasons. First, it is too heavy—an insuperable objection;—secondly, where the temperature of the air ranges pretty steadily a little below that of the skin, the flannel is much too slow a conductor of heat from the body. Thirdly, the spiculæ of the flannel prove too irritating, and increase the action of the perspiratory vessels on the surface of the body, where our great object is to moderate the process. From the second and third objections, indeed, even cotton or calico is not quite free, unless of a fine fabric, when its good qualities far

counterbalance any inconvenience in the above respects.

The great object of tropical prophylactics being TO MODERATE, WITHOUT CHECKING THE CUTICULAR DISCHARGE, I would here enter a caution against a too frequent changing of the body linen. a habit confined to newly arrived Europeans principally. To change morning and evening is enough for all and every purpose. even in the hot and rainy seasons; and to change oftener is simply injurious. The property which frequent change of linen has, in exciting the cuticular secretion, and the effects resulting from the sympathy of the skin with the stomach, liver, and lungs, may account, in a great measure, for the superior health which accompanies cleanliness in our own climate; and, on the contrary, for many of the diseases of the indigent and slovenly, which are so frequently connected with, or dependent on, irregularity or suppression of the cuticular discharge. But though this is true, by the injudicious, nay, injurious habit of too frequent change of linen in a tropical climate, the fluids on the surface of the body, already in excess, are thus powerfully solicited, and the action of the perspiratory vessels, with all their associations, morbidly increased, instead of being restrained.

The necessity which tyrant custom, or perhaps routine, has hitherto imposed on us, of continuing to appear in European dress particularly in *uniforms*, on almost all public occasions, and in all formal parties, under a burning sky, is not one of the least miseries

of a tropical life!

It is true that this ceremony is waived in the more social circles that gather round the dinner-table, where the light, cool vestures of the East supersede the cumbrous garb of northern climates. But it is laughable, or rather pitiable enough, to behold a set of griffinish sticklers for decorum, for some time after each fresh importation from Europe, whom no persuasions can induce to cast off their exuviæ, even in the most affable company, pinioned in their stiff habiliments, while the streams of perspiration that issue from every pore, and ooze through various angles of their dress, might almost induce us to fear that they were on the point of realizing Hamlet's wish; and that, in good earnest, their

"Solid flesh would melt— Thaw, and resolve itself into a dew,"

To the above observations on dress may be added, that no European should voluntarily expose himself, at any season, to the direct rays of the sun. If forced to be out of doors, the chatta should never be neglected, if he wish to guard against coup-desoleil, or some other dangerous consequence of imprudent exposure.

FOOD.—All must agree with Celsus, that "sanis omnia sana—and with a late eminent physician, that, an attention to quantity is of infinitely more consequence than the quality of our repasts; and also that an over fastidious regard to either will render us unfit for society, and not more healthy after all; yet, when we change our native climate for the torrid zone, many of us may find, when it is too late, that we can hardly attend too strictly to the quantity and quality of our food during the period of being accomodated to the new climate; and that a due regulation of this important matter will turn out a powerful engine in the preservation of health.

It is now well known, from dire experience, that instead of a disposition to debility and putrescency, a congestive, and sometimes an inflammatory diathesis, with tendency to general or local plethora, characterises the European and his diseases, for some years at least after his arrival between the tropics; and, hence, provident Nature endeavours to guard against the evil by diminishing our relish for food. But, alas! how prone are we to spur the jaded appetite, not only "by dishes tortured from their native taste," but by more dangerous stimulants of wine and other liquors, as well as by condiments and spices, which should be reserved for that general relaxation and debility which unavoidably supervene during a protracted residence in sultry climates. Here is an instance where we cannot safely imitate the seasoned European; for there are no points of hygiene to which the attention of a newcomer should be more particularly directed, than to the quantity and simplicity of his viands. They are practical points, entirely within his own command, and a due regulation of them is not at all calculated to draw on him the observation of others-a very great advantage.

That vegetable food is, generally speaking, better adapted to a tropical climate than animal will be admitted, especially in the instance of the unseasoned European; not that it is quicker or easier of digestion, for it is slower, but it excites less commotion in the system during that process, and is not apt to induce plethora afterwards. These are the considerations which should give preference to vegetable over animal substances in the diet of

the unseasoned European, and which should induce him to be at

least sparing in the use of the latter.

Every valetudinarian, and particularly, the hectic, knows full well the febrile movement which follows a full meal; and the same takes place, more or less, in every individual, whatever may be the state of health at the time. How cautious should we be then of exasperating those natural paroxysms, when placed in situations where various other febrific causes are constantly impending over, or even assailing us! The febrile stricture which obtains on the surface of our bodies, and in the secerning vessels of the liver, during the gastric digestion of our food, as evinced by diminution of the cutaneous and hepatic secretions, will be proportioned to the duration and difficulty of that process in the stomach. and to the quantity of ingesta; and as a corresponding increase of the two secretions succeeds, when the chyme passes into the intestines, we perceive the propriety of moderating them by abstemiousness, since they are already in excess from the heat of the climate alone, and this excess is one of the first links in the chain of causes and effects that leads ultimately to various derangements of function and of structure in important organs, as exemplified in the fevers and dysenteries, in the hepatitis and cholera of tropical

The newly arrived European should content himself with plain breakfast of bread and butter, with tea or coffee, and avoid indulging in meat, fish, eggs, or buttered toast. The latter alone often disagrees, and occasions rancidity, with nausea at stomach, while it increases the secretion of bile, already in excess. A glance indeed at the *Bawurchee*, buttering our toast with the greasy wing of a fowl, or an old dirty piece of rag, will have more effect in restraining the consumption of the article than any didactic precept which can be laid down; and a picturesque sight of this kind may be procured any morning by taking a stroll into the purlieus of the

kitchen.

In regard to dinner, were the European master of both time and circumstance, the early part of the afternoon should be that of his principal meal; but the great majority of men of business, whether official or mercantile, are unable to disengage themselves during the day, and thus from seven to eight o'clock becomes the settled hour.

It is true that military men, excepting such as hold staff offices, may choose their hour of dinner, but fashion and routine rule this and other habits; and the mess hour is too late. Naval and military men are, beyond all others, devoted to system, and to a certain extent justly so. But system, like everything in this world, may be abused so as to degenerate into something even worse than mere routine. As matter of fact, regiments in

general have luncheon ("tiffin") on table at one o'clock, with wine and ale, and a heavy dinner at seven with more wine and ale. Thus any officer who is so disposed may become the subject of a daily feverishness, soon to be followed by disease; and even such as are temperately disposed commit excesses in eating and

drinking, often without being aware of it.

An old staff officer in Fort William used to say that he had known more duels, courts martial, and dismissals, to result from the "tiffin" alone than from any other cause; but what were the other results, in the olden times, of the tiffin and the dinner, from year's end to year's end, there is no man alive now-a-days to tell! Although a great deal that was objectionable in the double meal has given way to modern improvement in the general habit of society throughout India, still too much of both tiffin and dinner remains. Both meals are greatly too much after the European fashion; while wine and ale are too liberally circulated, at least for health.

He then who consults his health in the East will beware of late and heavy dinners, particularly during the period of probation, but will rather be satisfied with a light and early repast as the principal meal, when tea or coffee at six or seven o'clock will be found a grateful refreshment. After this his rest will be as natural and refreshing as can be expected in such a climate, and he will rise next morning with infinitely more vigour than if he had crowned a sumptuous dinner with a bottle of wine the previous

evening.

Let but a trial of one week put these directions to the test and they will be found to have a more substantial foundation than

theory.

A limited indulgence in fruits, during the first year, is prudent; and there is little reason to believe that, when ripe and used in the forenoon, they dispose to irritation of the bowels. Particular kinds of fruits have peculiar effects on certain constitutions; thus, mangoes have sometimes a stimulating and heating effect, which not seldom brings out a plentiful crop of pustules, or even boils, on the unseasoned European. The pine apple, though very delicious, is not the safest fruit to make too free with at any time. The orange is always grateful and wholesome, and good ripe shaddocks are so likewise, owing to their cooling subacid qualities. The banana is wholesome and nutritious, whether undressed or frittered.

The use of spice or condiment, as already stated, should be reserved for those ulterior periods of our residence in hot climates when the tone of the constitution is lowered, and the stomach participates in the general relaxation. They are then safe and salutary.

DRINK .- In a vast empire, such as that of India, held by the frail tenure of opinion, and where the current of all the various religious opinions of the natives runs strong against intoxication, it was early found necessary from motives of policy rather than of health, to discourage the acquisition of habits at once dangerous and disgraceful. Hence the inebriate has always been justly considered as not merely culpable in destroying his individual health, but as deteriorating the European character in the eyes of the natives, whom it is on all accounts desirable to impress with a just sense of our superiority. It is thus that our power may acquire material substance from the gradually improving convictions of the natives themselves. Happily what has been promotive of our own interests has been also preservative of our health, as well as conducive to our happiness; and the general temperance which now characterises the European circles of society in the East Indies is most gratifying. We shall no longer hear from medical or lay writers of "gently stimulating liquids," used during the forenoon "for supporting perspiration, and for keeping up the tone of the digestive organs," for all experience has proved such habits to be actually pernicious. The European who should now take to such unworthy indulgences would soon find himself

excluded from good society.

It has already been observed that the great physiological rule for preserving health in hot climates is-TO KEEP THE BODY COOL. The strong sympathy that subsists between the skin and the several internal organs, as the stomach, liver, and intestinal canal, has also been emphatically referred to. The common sense of mankind would seem indeed to point out the propriety of avoiding heating drinks, for the same reason that we endeavour instinctively to guard against a high external temperature. But the truth is that, until men begin to feel the corporeal ill effects of intemperance, a deaf ear is turned to the most impressive lectures against the most deplorable of propensities, that which Napoleon declared to be the least compatible with greatness. With the feeble and irresolute the magic bowl, which this moment can raise its votaries into heroes and demi-gods, will, in a few hours, sink them beneath the level of the brute creation. Moralists and philosophers have long descanted on this theme, but with little success, as few will attempt to prove that water is the simple and salutary beverage designed by nature for man; seeing that every nation, even the most refined, has practically repudiated the doctrine. Let the medical profession, however, do its duty in portraying truthfully the ill effects of the abuse of drink, in tropical climates especially. The truth is that as drunkenness leads, in a moral point of view. to every crime, so, in a physical point of view, it promotes the invasion, and retards the cure, of every tropical or other disease.

It may be received as a truth that, during the first two years of residence at least, the nearer we approach to a perfectly aqueous regimen in drink, so much the better chance have we of avoiding sickness; and the more slowly and gradually we deviate from this afterwards, so much the more retentive will we be of that in-

valuable blessing, HEALTH.

In speaking of the means of preserving health, which depends so much on the tone of the digestive organs, we must not omit the recent luxury—the American ice—which has now become an article of necessity in the East. It would be out of place here to speak of its uses in the treatment of various diseases, and of fevers especially, as those of the cerebral and gastric nature. "Nothing," says Dolomieu, "is more salutary during the sirocco than iced beverages; they revive the spirits, strengthen the body, and assist the digestion." Those who have now, during several years past, made use of this real luxury in the East Indies, need no arguments in favour of promoting so remarkable an instance of American enterprise.

I would caution the newly arrived European against a very common and very dangerous mistake, namely, the acting in matters of diet and exercise on the supposition that he may with impunity do as the elder residents; for it is consonant with experience, as with theory, that the *latter* class may indulge in the luxuries of the table with infinitely less risk than the former. To think and act otherwise is to confound all discrimination between very different habits of body which the seasoned and un-

seasoned possess.

One other circumstance should always be held in recollection, to wit, that when a course of temperance is fully entered on, and the pleasures of temperance fully enjoyed, no consideration should induce us to commit an occasional debauch, especially during our seasoning, for we are at those times in infinitely greater danger of both endemic and epidemic attacks than the habitual bacchanal.

In the last article it has been stated that the subacid fruits are both grateful and wholesome, and so will it be found with moderately acid drinks, such as sherbet. Nature seems to point out the vegetable acids in hot climates, as grateful in allaying drought and diffusing a coolness from the stomach all over the body. To the temperate and the healthy they will generally prove salutary, and I have never seen or heard of any ill effects from their use.

We hear much amongst vulgar habitual topers of the supposed prophylactic influence of spirits and cigars against night exposure, malaria, and contagion; but no medical observer, in any of our numerous colonies, has ever seen reason to believe in any such delusive doctrine, nor is there in reality the smallest foundation for it. All excitement is followed by a corresponding depression of the vital functions, and it is then that the toper is doubly liable to suffer.

EXERCISE.—This is one of the luxuries of a northern climate to which we must, in a great measure, bid adieu between the tropics. The principal object and effect of exercise in the former situation appear to consist in keeping up a just balance in the circulation, in supporting and maintaining the functions of the skin, and promoting the various secretions. But the perspiration, biliary, and other secretions being already in excess in equatorial regions, a perseverance in our customary European exercises would prove highly injurious, and it often does so, by promoting and aggravating the ill effects of an unnatural climate. As such excess very soon leads to debility, and to diminishing action in the functions alluded to, and to a corresponding inequilibrium of the blood, so it is necessary to counteract these by such active or passive exercise as the climate will admit at particular periods of the day or year; a discrimination imperiously demanded if we mean to preserve health. When the sun is near the meridian for several hours of the day on the plains of India, not a leaf is seen to move, every animated being retreats under cover, and even the "Adjutant," or gigantic crane of Bengal, soars out of reach of the earth's reflected heat, and either perches on the highest pinacles of lofty buildings, or hovers in the upper regions of the air, a scarcely discernible speck.

At this time the native retires instinctively to the innermost apartment of his humble shed, where both light and heat are excluded. There he sits quietly in the midst of his family, regaling himself with cold water or sherbert, while a gentle perspiration flows from the skin, and contributes naturally and powerfully to his refrigeration. Mr. Twining justly observes that the natives, both Hindoo and Mahomedan, seem to suffer much from the hot season in Bengal. During the latter part of this season those natives whose circumstances enable them to act as they wish, avoid any exertion as much as possible, reducing the quantity of their food, and eating certain fruits which they consider cooling. In the afternoons they drink the fluid contained in the unripe cocoa-nut, or a very simple sherbet, or some sugar and water, deeming the latter especially cooling; and in the mornings they take an infusion of Nalta Pat, or leaf of the Corchorus Olitorius, which they say has a cooling effect, at the same time that it acts as a mild tonic, promoting digestion, and preventing lassitude. He adds that the natives, though adapted by nature to bear the climate, take more care to moderate the effects of heat than Europeans, especially in their light clothing, abstemious food, and tranquil habits.

After this example of the salutary habits of all classes of natives of India, it is hardly necessary to urge the injuriousness of every kind of active exercises to Europeans under tropical heats, and especially during the heats of the day, yet hundreds perished annually from this very cause, particularly in the West Indies, upon each influx of European troops during the late war. Happily it may now be said that, owing to the removal of the European troops from the plains to the mountains, the horrors resulting from the climate of Jamaica have been exchanged for health and comfort. This beneficent act is due to the generous and noble nature of the late Lord Metcalfe, when governor of that island.

Who would expect to find dancing a prominent amusement in a tropical climate? The natives of the West Indies are exceedingly fond of this exercise; but those of the East are wise men still, for instead of dancing themselves they employ Natch girls to dance for them.

Gestation of every kind, whether in palankeens or spring carriages, is a species of passive exercise exceedingly well adapted to a tropical climate. The languid state of the circulation of the blood, in those who have resided long there, is pointedly exhibited in the disposition which they evince to raise the lower extremities on a line with the body when at rest; and this object is completely attained in the palankeen, which indeed renders it a peculiarly agreeable vehicle.

On the same principle we may explain the feeling of satisfaction and the utility of shampooing, where the gentle pressure and friction of the soft Asiatic hand over the surface of the body, but particularly over the limbs, invigorate the circulation after fatigue, as well as after long inaction, and thus excite the insensible cuticular secretion. The Keesa, or hair glove of India, is an admirable means of giving additional effect to the practice of shampooing, a practice which, to the indolent wealthy natives, proves a real and effective substitute for exercise.

The swing might perhaps be rendered useful in the hot and rainy season in the East Indies. In chronic disorders of the viscera it could hardly fail to be grateful and salutary, by its tendency to determine to the surface and relax the subcutaneous vessels, which are generally torpid in those diseases. It might be practised in the early mornings and evenings within doors, whenever the weather or other circumstances do not admit of gestation in the open air.

BATHING.—Moseley's caution to the European residents of his day in the West Indies, may be recorded for the benefit of all who think that, in hot climates, they may do what they please with cold

water. "I DARE not recommend cold bathing; it is death with intemperance, and dangerous where there is any fault in the viscera. It is a luxury denied to almost all, except the sober and abstemious females, who well know the delight and advantage of it." The cold bath is death, not during intemperance but in the collapse which follows a debauch, or indeed any other great fatigue of mind or body. It is also dangerous under every form of visceral disease, but the healthy and temperate of habit may as safely partake of the "delight and advantage" of the cold bath as Dr. Moseley's "sober and abstemious females." The truth is, that the cold bath is a prize due to, and gained by, the temperate. To all else it is eminently unsafe, as any naval and military surgeon of tropical experience can testify. To the European resident in tropical elimates whose constitution is sound, and whose habits are temperate, no more efficient means exists of obviating the most unpleasant effects even of the cold season; for he who reacts well under the cold bath will not be troubled with dry skin, and sense of internal fulness. To persons of ordinary health, but who are not robust, the cold bath will be found tonic and agreeable from the beginning of March, in India generally, to the end of September. The temperature ranges high in these months, and the determination to the surface is such as to ensure a sufficient reaction. It is a common error to think that before using the cold bath we must get cooled first, while the very opposite rule is the correct one. To the delicate, indeed, immersion in a warm bath for a few minutes is an excellent preliminary, followed at once by affusion of some three or four vessels of cold water. A glow over the whole surface of the body will immediately follow. This is a safe and excellent mode of bathing to all who shrink from or who feel doubtful of salutary reaction from the use of cold water.

We hear warm and cold bathing recommended without reference to the state of health, to the season, or to regularity of habit, although these circumstances should form essential preliminaries to the choice. It may be concluded for certain that, to persons who have suffered from tropical disease, or who are affected with visceral congestions, or with visceral enlargements especially, the results of fevers or dysenteries, the warm bath is the only safe one at all seasons. The same rule applies to the dissipated, and to such as are in the habit of keeping late hours. In such persons the balance of circulation is already disturbed, and the effect of cold water is to throw the blood with force on organs already irritated by irregular courses of life, the abdominal viscera especially.

Under such unfavourable circumstances it is not to be expected that the "conservative energies" should be capable of being "roused to successful resistance;" with the whole external surface parched, and with the digestive mucous surfaces in a state of irritation, how can it be otherwise?

As many persons have but an imperfect knowledge of the relative temperatures of the several baths in ordinary use, the following scale is here presented:—

Cold Bath, from			60° to	75°
Tepid Bath, from			85° to	92°
Warm Bath, from		4	92° to	98°
Hot Bath, from			98° to 1	12°

By the healthy and temperate European the use of the *cold* bath should be regularly and daily persevered in, from the moment of his entrance within the tropics; and when from long residence there, the functions above referred to begin to be irregular, or defective, he may prudently veer round by degrees to the use of the *tepid* bath, which will then be found a most valuable part of

tropical Hygiene.

The use of the cold bath being a passive operation, unattended by any exercise, it may be used at any period of the day, although the mornings and evenings are generally selected by Europeans in the East; immediately after leaving their couch and before dinner. On both occasions the bath is very refreshing, and it powerfully obviates that train of nervous symptoms so generally complained of by our countrymen in hot climates. Before dinner it seems to exert its salutary influence on the surface of the body, and by sympathy on the stomach, removing the disagreeable sensation of thirst, which might otherwise induce a too free use of potation during the repast. It is always imprudent to bathe while the process of digestion in the stomach is going on, as it disturbs that important operation.

SLEEP.—When we bid adieu to the temperate climate of Europe, with its "long nights of revelry," and enter the tropics, we may count on a great falling off in this "solace of our woes." The disturbed repose which we almost always experience there, has a greater eventual influence on our constitutions than is generally supposed, notwithstanding the silence of authors on this subject. Whatever we detract from the requisite period of our natural sleep will surely be deducted, in the end, from the natural range of our existence, independently of the predisposition to disease which is thus constantly generated. This is a melancholy but a true reflection, and it should induce us to exert our rational faculties in obviating so great an evil.

The great object of the European is to sleep cool, and obtain complete protection from Mosquitoes. Happily both advantages may be secured to the temperate by the large mosquitoe frame and

curtain with punkah suspended from the ridge, as now prevalent throughout Bengal. This is not only luxurious but safe, the gentle agitation of the punkah removing the heated air surrounding the body without exposing the person to the dangers arising from sudden night changes in the temperature or humidity of the

atmosphere.

The European is thus enabled to procure more and sounder rest than he could possibly do otherwise; and by giving his frame a more thorough and complete respite from the great stimulus of heat, he imparts to it tone and vigour, so necessary to meet the exhaustion of the ensuing day, as well as to repair that of the preceding. Early hours are here indispensable; for the fashionable nocturnal dissipations of Europe would soon cut the thread of our existence within the tropics. The order of nature is never inverted with impunity, even in the most temperate climates; beneath the torrid zone it is certain destruction. The hour of retirement to repose should never be protracted beyond ten o'clock; and at daylight we should start from our couch to enjoy the cool, the fragrant and salubrious breath of morn.

Without some artificial aid, such as that above mentioned, a great waste of strength—indeed of life—may arise from our inability to obtain *cool* repose at night. The cold and rainy seasons, heavy dews, or exhalations from contiguous jungles or marshes, often render it impossible with safety to sleep in the *open air*; a practice, during the *hot season*, fraught with refreshing benefit, where the obstacles mentioned do not prevent its execution.

In Bengal Proper, in the plains of Upper India, and on the Coromandel coast, except during the hot land winds there, or at the change of the Monsoons, Europeans may generally indulge, during the hot and dry season, in the luxury of sleeping in the open verandahs, not only with safety, but with infinite advantage. It is an old habit throughout most parts of India, especially of the military classes; and the judicious Captain Williamson says justly that, while it is attended with the greatest refreshment, enabling them to rise early, divested of that most distressing lassitude attendant upon sleeping in a close oppressive apartment, communicating a febrile sensation, very few instances could be adduced of any serious indisposition attending it. Healthy and temperate persons, and who are habituated to the use of the cold bath, need be under no apprehensions, excepting at the places and seasons alluded to, as to transitions from the scorching heat of the day to the serenity of night, for indeed it is gradual and easy.

Those who habitually exclude themselves from the breath of heaven, whether from inclination or necessity, become languid from the *continued* operation of heat and want of repose:—even the slightest aerial vicissitude, or admission of a partial current of

cool air, unhinges the tenor of their health, and deranges the functions of important organs. These are they who require the afternoon siesta, and to whom indeed it is necessary, on account of the abridged refreshment and sleep of the night; while the others are able to go through the avocations of the day without any such substitute—a manifest and great advantage.

A few words on the incubus, or night-mare—a troublesome visitor to the tropical couch—may properly conclude this section.

From the results of treatment it would appear that the primary cause has its seat in the digestive organs, in whatever way it may act; and that the night-mare originates in defective digestion, producing heart-burn, flatulence, griping and eructation, with a train of dyspeptic complaints.

Of all medicines, the carbonate of soda will be found the most efficacious, taken in scruple doses at bed time, or night and morning, in some aromatic water, such as the peppermint. This medicine not only neutralizes the acids of the stomach, but it promotes the elimination of the bile, and the evacuation of slimy dis-

charges from the bowels.

There are few people with whom particular kinds of food do not disagree, and, being known, these should be avoided. Thus, chestnuts or sour wine will almost always produce incubus in those predisposed to it, as was observed by Hildanus:—" Qui scire cupit quid sit incubus? Is ante somnum comedat castaneas, et superbibat vinum fæculentum." In this country, cucumbers, nuts, apples, and flatulent kinds of food are the articles most likely to bring on night-mare.

THE CONDUCT AND GOVERNMENT OF THE PAS-SIONS.—Most of the precepts that apply to the regulation and government of the passions in cold climates will be found to apply to them in tropical climates; but it is necessary to correct at once an erroneous impression that there is something peculiar to the tropics which excites certain passions in a higher degree than in

temperate regions.

Dr. Moseley says that, "there is in the inhabitants of hot climates, unless present sickness has an absolute control over the body, a promptitude and a bias to pleasure, and an alienation from serious thought and deep reflection. The brilliancy of the skies and the beauty of the atmosphere conspire to influence the nerves against philosophy and her rigid tenets, and forbid their practice among the children of the sun." However true this description may be in respect of "the children of the sun," it does not accurately exhibit the condition of the stranger European, in whom such a course of relaxation would be very immoral; for such a view would furnish the dissolute libertine with a physical

excuse for his debaucheries, when the real source might be traced to laxity of religious and moral principle. We would ask Dr. Moseley, if the "promptitude and bias to pleasure" be increased in hot climates, why the ability to pursue or practice it should be

lessened? a fact well known to every debauchee.

It has been asked, says Hennen, what has the medical topographer to do with the morals of the natives of a country? and it has been asserted that their immoralities cannot affect the health of troops quartered amongst them, if proper discipline be observed. Such opinions are founded upon a very superficial view of the subject; for the soil and inhabitants, he adds, always react on each other. A sober and industrious race of people will, for example, have a greater desire to improve their country than men of a contrary character, and will also possess greater physical power to carry their desire into execution. Place such a body of men in a district overrun with noxious weeds and timber, and fast degenerating into a morass, and can there exist any rational doubt that they will clear it sooner, and preserve it longer in that improved state than men of a different disposition? Place in a similar situation, or even in a district thus improved, a body of men who are idle and intemperate, and the immediate result will be that the soil will deteriorate for want of proper care, the weeds will reappear, the drains will become obstructed, the edible products of the earth will lessen in quantity, and diminish in their nutritive quality; the inhabitants will become unhealthy from the bad state of their grounds; and the diminution of their physical powers thus produced will disable them progressively more and more from remedying the causes of the evil. Many of these effects will doubtless first be felt in their own persons, but it is undeniable that they must ultimately operate on their visitors. On this obvious principle is founded the axiom in Medical Topography-" that a slothful, squalid-looking population invariably characterises an unhealthy country."

Whether the founders of the Hindu faith acted designedly or otherwise, there can be no doubt that, by depressing all the physical energies, by a diet purely vegetable, they fastened with a firmer and stronger band the bonds of Brahminical domination on the people of India; until a chain, forged only by superstition, became in progress of fifty centuries, and through the most

powerful of moral and physical agencies, strong as death.

"The use of certain kinds of food and drink," says Cabanis, "may tend to confirm or impair certain moral habits. Sometimes it may operate directly, and by the immediate impressions which it produces; at other times, by the different states of health or disease which it occasions, or by the changes in the fluids and solids which result from it; for all these different alterations in

the system soon manifest themselves more or less distinctly in the ordinary dispositions of the will and understanding."

The same great authority in the physiology of climates adds that, "the poor diets prescribed by the legislators of various religious orders, have never had the effect of diminishing the venereal appetite, but have, on the contrary, inflamed these propensities the more, or disordered the imagination in diminishing the physical forces; and thus men have been rendered more feeble,

more unhappy, and more easy of domination."

But let us hope that, in respect of the natives of India, the all-powerful benefits of education, and the example of their European governors, may enable them to conquer the influences of climate, and of the depraving religious and political habits of ages. To expect, however, that such changes can be speedily effected were contrary to reason and experience. The caste is of unknown antiquity. "Neither the proselytising sword of the Mussulman, nor the mild light of Christianity, has had any influence upon it, and the Hindu still worships before the altars of his gods with the same devotion as when Orpheus charmed the wild beasts by the sounds of his lyre, and when Moses ascended Mount Sinai. Religion, manners, customs, costume, civilization, all have remained immovable, as the temples hewn out of the granite rocks of Ellora."

After what has been here stated, it cannot be questioned that the habits and character of the people amongst whom we live, during many years, must have a powerful influence on our morals. But to return to the condition of the stranger European. The removal of religious and moral restraint, the temptations to vice, the facility of the means, and the force of example, are the real causes of this bias to pleasure; and in respect to the effects of licentious indulgences between the tropics, the reader may be assured that he will find, perhaps when too late, how much more dangerous and destructive they are than in Europe.

The nature of the supposed "propensity" has been explained to him; and as the principal cause resides neither in the air, nor in the "brilliancy of the skies," but in his own breast, he has no excuse for permitting it to grow into the wild luxuriance of un-

bridled excess.

The monotony of life and the apathy of mind so conspicuous in hot climates, together with the obstacles to matrimony, too often lead to vicious and immoral connections with native females, which speedily sap the foundations of principles imbibed in early youth, and involve a train of consequences not seldom embarrassing, if not embittering every subsequent period of life. It is here that a taste for some of the more refined and elegant species of literature will prove an invaluable acquisition for dispelling ennui—the moth of mind and body.

## THE REMITTENT FEVER OF BENGAL.

This fever has been long known to, and carefully described by. British writers on the diseases of the East. Some of the older observers termed it Putrid Intestinal Remitting, and Putrid Remitting Marsh, Fever; perhaps without being sufficiently aware that in those days, owing to the salt diet, crowding, and other sanitary neglects in their then long voyages, a general scorbutic taint existed, amongst the newly arrived Europeans especially. Since then, and under the various designations of Jungle, Hill, Teraèe, Bilious Remittent, Marsh Remittent, or Malarious Fever, it has been described by more modern writers. Robert Jackson terms it Gastric, or Bilious Remittent, and describes it as a fever belonging to all countries; but as endemic in the West Indies at all seasons of the year. Though common to all countries, and observed occasionally at all seasons of the year, he justly adds that it prevails more generally in warm countries, on extensive alluvial plains, and in the autumnal season.

Remittent fevers will be found almost everywhere throughout the East Indies, varying in their intensities, and in their complications, as they may occur in the Deltas, along the marshy banks, or in the embouchures of rivers; in the plains extending from the bases of mountain ranges, termed teraées; in partially inundated or irrigated lands, or in such as are traversed by percolating streams, or by canals; in wooded districts, termed jungles; or in certain hill districts. The seabord, especially where there is jungle, or salt marsh, and the adjacent islands, when of a jungly or marshy nature, are peculiarly pestilential; and so are often found the drying-up marshes, and the drying-up beds of rivers.

The British army has often been fever stricken, and occasionally destroyed, in low, dry-looking, sandy plains, such as those of Walcheren and of Rosendaal; on the elevated, rocky, and dry-looking countries, and in the half-dried water-courses of Portugal; on the plains of Spain, along the course of the Guadiana; but everywhere there was "water to within a few inches of the surface;" the ravines were always "half-dried;" and the Guadiana itself consisted of but "lines of detached pools."

Whether on the coast of Kent, about Dungeness, "during the hot summer and autumn of 1807," or in the West India Islands, the same topographical conditions were found to exist by William Fergusson;—there was generally the presence of actual marsh, or damp; and always the antecedents of heavy rain, or of "flooding in the rainy season." "The fevers of Cadiz, Carthagena, Gibraltar, and Zealand," says Dr. James Johnson, "may compete in respect to virulence and fatality, with those of Batavia, Bengal,

St. Domingo, and Philadelphia."

Varying somewhat in type and in complication, each depending on locality, on constitution, and on habit of life, the essential characters of the remittent fever are still everywhere the same; and, whether as cause or effect, this disease has much to do with almost all the derangements of health of Europeans in the East. It is also by far the most prevalent of Indian endemic diseases; and on the right understanding of its movements, peculiar nature, complications, and just treatment, will always, and in a great measure depend, the usefulness of the Indian medical officer.

According to a table furnished to me by Colonel Tulloch, out of an aggregate European force of 25,431 men, of her Majesty's army, serving in periods of eight and ten years respectively, between 1823 and 1836, in the stations of Calcutta, Chinsurah, and Berhampore, all in Bengal Proper, 13,596 cases of fever, remittent, intermittent, and continued, occurred;—whereas, out of the same aggregate force, 8,499 cases of dysentery and diarrhœa occurred—thus establishing the far greater prevalence of the malarious fevers over diseases of the bowels.

In the history of the remittent fever of Bengal, during the last hundred years, the first observation that presents itself is—the great difference as to its intensity in the present as compared to former times; and, secondly, the causes of this difference.

The earliest account we possess of the state of public health, and of the season of greatest mortality in Calcutta is, that of Captain Hamilton (1688—1723), wherein he mentions 460 burials out of 1,200 British inhabitants, from August to the en-

suing January.

Of Major Kilpatrick's detachment of 240 men, mostly Europeans, stationed at Fultah, Ives tells us, that "not thirty of the whole were left alive between August and December, 1756, by one of these epidemics." He adds that, "the number of men buried in Bengal amounted to more than half of all who died in the several hospitals in India, during the whole term of Admiral Watson's command, a period of three years and one month." Dr. Bogue, who also served in Watson's fleet, says that, "out of three ships of the line and a twenty gun ship, and those not fully

manned, we lost in six months upwards of two hundred men, most of whom died of these fevers."

Dr. James Lind was surgeon of the Drake Indiaman, and writes of the fever "which raged in Bengal, in the year 1762," and which he terms Putrid and Remitting Marsh Fever. This fever "raged more or less in different places, according as the soil was more or less swampy;" and, "became so violent," during the rainy season especially, "as to end in death during the third fit, which is generally the case." Others, he says, "were exposed to the danger of dying at every fit;" and "when the disorder continued for any time, without a change, it generally ended in death; while the weather grew better it sometimes, in the space of a few days, from a common fever became an intermittent one, and the patient recovered, unless his liver, which was sometimes the case, happened to be affected."

Stavorinus, the Dutch naval commander, speaking of "the sort of sickness or fever" which prevailed amongst the European inhabitants of Calcutta, during his visit to that city in 1766—1771, says that "it generally sweeps away those who are attacked by it in the space of three days."

Dr. John Clark, who visited Calcutta between 1768 and 1771, says that "the fever and flux" were very fatal in the former year. Of the fever, he states that "it frequently carries off the patient in twelve hours." . . . "During the sickly season in Bengal, the uncertainty of life is so great that it frequently happens that one may leave a friend at night in perfect health who shall not survive the following day. There have been several melancholy instances of persons who have returned home in a state of perfect health, from performing the last duties to a deceased friend, and have next day been numbered with the dead."

The same authority records that out of 189 cases of fever treated in ships trading to the various ports in the East, "105 recovered, and 84 died." Again, "out of 876, the complement of men belonging to eight ships, 78 died in Bengal, and 55 at sea or nearly one in six. Both the fever and flux, if obstinate, have an equal tendency to terminate in abdominal obstructions, particularly in fatal swellings and suppurations of the liver." From this truthful record, and looking to the sanitary condition of the sufferers, and to the results of the treatment by bark, we are constrained to infer that the ultimate recoveries must have been but few, out of all who were the subjects of the fever or the flux.

Mr. Magennis states that in 1784, out of the crew of his ship, the Valentine, and six others stationed at Khidgeree, there died of fever and dysentery, 170 men, the usual period of their stay in the Hooghly being from August to January. Curtis writes that about the same time, out of two companies of the 98th and 100th regi-

ments embarked in England for India, there died during a "suffering and tedious passage of exactly eleven months, 75 men, 40 of them being from fever, 18 from dysentery, and the remainder from scurvy and cachexy." It was on this occasion that the 2nd battalion of the 42nd regiment alone suffered a loss, by the time it landed in Calcutta, of five officers and 116 men, all from fevers, bowel complaints, "and the scurvy." It would appear that the unfortunate companies referred to by Curtis, had "some occasional fillings up from the other ships as the numbers decreased;" but, making allowance for this circumstance, the loss of life was horrible—especially, when we reflect that each ship in a fleet shared a like mortality.

The ships were, in the language of Clive, "crammed with soldiers;" so that, along with salt rations, there was crowding, and consequent filth and want of ventilation—making altogether a combination of the most unfavourable circumstances to the soldier's

health.

That a scorbutic taint was very general in these times may, I think, be admitted; and this circumstance will account for the general term "putrid," as applied by the older writers to the endemic fevers and dysenteries: this unfavourable complication will go far also to account for the enormous mortality. Curtis states, that in the open well-ventilated Naval Hospital at Madras, containing from four to five hundred men, "the great bulk of the cases were ulcers;" indeed he constantly refers to the scorbutic taint as prevalent amongst both soldiers and seamen, a fact almost entirely overlooked by modern writers when treating of the earlier results of treatment in tropical diseases. Dr. James Lind, in his excellent "Treatise on the Putrid and Remitting Marsh Fever of Bengal in 1762," refers expressly to the prevalence of scurvy amongst the crews of all the ships off Calcutta. So recently indeed as the beginning of the present century, James Johnson states that, in the small portion of the river running between Calcutta and Khidgeree, full 300 European seamen, or more than a fourth of the ships' crews, fell annual victims to the ravages of remittent fever.

But happily we have no longer to record any such fearful ravages by endemic diseases in the Capital of the East; neither do such sweeping epidemics as that recorded by Clark in 1770, with its cold stage of twelve hours, occur, carrying off 80,000 Natives and 1,500 Europeans. Such pestilences seem to have gone from us; and we find that here, as in the Western Hemisphere, the malignant fevers of former days, if they have not-disappeared, are at least mitigated. Even in Jamaica, although severe epidemic fevers sometimes recur, they do not now, as formerly, destroy "to the amount of the whole number of its white

inhabitants once in five years."

Of a just and useful comparison, so far as relates to the means of cure, and the results, in former as compared to recent times, there can therefore be little or none. But this much may be conceded in favour of the position of modern Indian medical officers; that owing to the better sanitary condition of their patients, to the better understanding of the value of depletory means, of a cooling regimen, and of the power of improved antiperiodic means, we not only save many more lives, but we leave far fewer of the sequelæ to fevers and dysenteries than of old. For instance, out of 284 Europeans treated by Dr. Paisley, in the Military Hospital at Madras, in 1782, 182 men suffered from some form of "visceral obstruction;" in other words, from organic disease of the liver or spleen, or of both, rendering a death more or less speedy a necessary result.

The object of these remarks is, principally, to excite the attention of the younger members of the profession in the East to the superior efficacy of measures of prevention, to those of cure of disease. In the first-named department of our profession, although a new science, the progress has been immense; whereas in the last, although a science of some two thousand years, or more, the results are by no means so signally useful, whether as regards

the benefits to the individual, or to the community.

With all the advantages of modern medical science, no one could, even now, pretend to any very great success in the cure of Europeans suffering at once from the united influences of the most concentrated malaria and of scorbutus. We know, in fact, that in the olden times, owing to the joint influences of the marsh poison, and of the blood depraved beforehand by a salt diet, and by crowding in ships and barracks, the seaman and soldier were destroyed by fever and dysentery within a few hours of their admission into hospital. Such men were in truth almost beyond any, and all, medical means of cure:—no amount of cinchona could have cured them; but very easily devised, and very obvious means of prevention MIGHT HAVE PRESERVED THEM IN HEALTH.

The causes of the present comparative improvement in public health must be of the highest interest and importance, especially to communities living within the tropics; and, with all just confidence in modern medicine, guided by the lights of an improved physiology, and those also of pathology, I cannot yet agree with those who would ascribe the whole of the difference here spoken of to superior modes of medical treatment, great as these confessedly are. It is not through modern improvements in the treatment of disease, as contrasted with the older modes, that public health has been so much amended, as through the great measures of prevention of disease, consequent on the progress of

improvement of localities, institutions of police, &c. It is to the preservative power of knowledge, to the reciprocal actions of the social state, and of political events upon each other, and upon medical science, that the advancement of public health is most indebted, and so it will continue to be, although these circumstances are not sufficiently weighed by some of us when, in our hurry to praise ourselves, we forget what is due to our predecessors of old, and that those last had frequently to treat a violent and complicated form of disease which we have never seen, and with whose fatal severity we are consequently unacquainted.

It is justly observed by a popular writer, that there never were any specifics discovered against the plague, the sweating sickness, or the leprosy; and yet the leprosy, the sweating sickness, and the plague, are now among the things unknown to us. They disappeared, not before any marvels of medicine, or any perfection of chemical science, but before the gradual amelioration of

our condition through sanitary improvements.

PREDISPOSING CAUSES.—The predisposing causes of fever will be found in all those conditions of the body, and in all those habits of life, which tend to spoil the blood, and which lower the tone of the nervous system. The depressing passions of the mind also tend greatly to the advent of fever, whether endemic or contagious; and here we are carried back to the voyages of thirteen months to Bengal, to the crowding of soldiers and seamen, to the mental despondencies, to the scurvies, to the mortal "putrid remitting Marsh fevers" of Europeans on landing in India, in the days of Clive and Hastings.

Extremes of every kind—extremes of supply and deficiency, of repletion and of privation;—excesses of every kind, especially in the use of spirituous liquors, wine, fermented drinks, and of tobacco, each and all of them dispose to fever; and so does exposure to extremes of heat and of cold, to heavy dews and fogs, to night air, to changes of season and other influences. Amongst seamen and soldiers proceeding to our various intertropical possessions, the salt rations and the spirits served to the men during the voyage used to be to a great extent, and are still, to some degree, very injurious to health, and very predisposing to fevers.

dysenteries, and hepatic diseases.

Excessive labour and the consequent fatigue, whether from the operations of sieges, or from long marches, powerfully dispose to fevers amongst soldiers; and when to labour and fatigue we add the absence of proper diet, the want of necessary cover from tents, want of sufficient clothing, and want of sleep, we shall be at no loss to account for the ravages of disease in camps—espe-

cially when to all the above causes we superadd the abuse of

stimulating liquors, and the malarious influences.

After the brief sketch here offered of the history and causes of the remittent fever of Bengal, the reader will be prepared to enter on a consideration of its symptoms, pathology, and treatment, as they are presented to us in our day.

SYMPTOMS AND PROGRESS.—In no other morbid state is the physiognomy of disease so marked with general distress as in the severer forms of tropical fevers; for whether we behold the sufferer in the stage of rigor, reaction, or that tending to collapse, the countenance and manner are expressive of anguish, varying

only in degree according to the stages.

The attack of the Bengal Remittent Fever is usually sudden, though occasionally the patient may complain for a day or more of weakness and lassitude, headache, nausea, and general uneasiness, accompanied by mental depression, or subdued anguish. The countenance is pale and shrunk—the conjunctive clear, and of a dull whiteness. During this premonitory stage of malaise the pulse is small, contracted, and frequent, with a slight increase of the temperature of the surface, especially over the forehead and præcordia, and a diminished action of the intestinal and

urinary organs, as well as of the skin. The actual invasion of fever is ushered in by debility of the mind and body, indicative of diminished power of the nervous system, by shivering, or horror, the sensation of cold recurring in paroxysms, not being continuous. The headache now becomes intense and darting, with sense of tension across the forehead as if with a tight-drawn cord, and with pain in the back and loins or on moving the eye-balls. The eyes now look muddy, or are tinged yellow, being void of lustre and expression; or else they have the appearance of deep seated inflammation; occasionally, they are protruded, giving a wild and passionate expression to a countenance already flushed and anxious. At other times the countenance is bloated and torpid, expressive of general congestion, or of what the patient terms inward suffering. There is frequently a distressing giddiness which occasionally proceeds to delirium. When this last symptom appears early, and runs high, we may anticipate a dangerous form of fever.

The tongue is red at the tip and edges, loaded, clammy, and moist; at other times, with a bitter or bad taste, the tongue is but little changed from the healthy appearance. The thirst is usually most distressingly urgent, and the previous nausea is now increased to actual and severe bilious vomiting, accompanied by distressing

sense of oppression and burning pain of the præcordia.

On examining the surface of the body, we find the skin thick, dry, and corrugated, giving to the hand a rough unpleasant sensation of pungent deep-seated and intense heat: occasionally, in severe complicated cases, the skin is of a deep yellow. In the fevers of the rainy season, and especially in persons of a phlegmatic habit and large abdomen, the skin is moist, clammy, and cadaverous, indicative of prostration of the nervous and vascular functions.

The abdomen generally, but more particularly the epigastric region, will be found tender on pressure, tumid and inelastic, accompanied by great anguish and oppression, caused by sanguineous congestion around the nervous centres. The secreting and excreting functions are all seriously disturbed, the bowels being usually constipated; but occasionally there are copious biliary evacuations; at other times, they are acrid and watery, while the urinary excretion is always scanty and depraved, though occasionally it also is copious, limpid, or tinged with bile.

The pulse, previously oppressed, becomes now hard and quick, ranging from 110 to 120 in the minute; and the force and frequency of circulation through the brain, superadded to the already disturbed condition of the nervous functions, gives rise to confusion of ideas, and loss of mental command, amounting occasionally

to actual delirium.

The respiration is hurried and irregular, sometimes interrupted by deep sighing; and with all these accumulated distresses, there is not a point on the entire sentient surface of the body on which the patient can recline for a moment with comfort. No posture affords ease or rest, the sufferer being harassed with a restlessness which nothing can appease, characteristic at once of gastric irritation and oppression, with much general disturbance of the great central nervous functions of life. In a brief space the sufferer goes through a terrible storm of diseased action, his throes of anguish, mental and corporeal, being truly distressing to behold.

After a time, this general distress of body and mind, so difficult, if not impossible to describe, begins to subside; and, with a relaxation of all the symptoms, comes a state more or less approaching to regularity of function, termed remission, more or less complete, according as the fever is mild or severe in its character.

These symptoms rise and fall in daily succeeding paroxysms, until they advance perceptibly into a condition of returning health, or recede and become aggravated and complicated into a state of imminent danger—the paroxysms running imperceptibly into each other, so as hardly to leave the least interval from suffering, during the brief but retrograde process that now remains to be gone through.

It sometimes, although rarely, happens, that a double paroxysm occurs within the twenty-four hours—a circumstance to be care-

fully ascertained, lest the physician's visits and his application of active remedies, and especially of bleeding and purging, should be mistimed; that is, used at the wrong time of remission, instead of

the right time, or that of the accession of fever.

The favourable stage again, or that of remission, is indicated by a gentle perspiration, a reduced temperature, a soft and expanded pulse, diminished in frequency down to about 90 in the minute—a full, free, and regular respiration, and a more abundant action of the secreting and excreting functions. In the rainy season and in dissipated and unhealthy subjects, as well as in the worst of our fevers, the stage of remission is the period of greatest danger—the relaxation proceeding too far, as shown by profuse sweats, coldness of the extremities, great anxiety, and sinking of the vital powers; in short, there is no stage of this fever that does not require a careful observation and watching; for each succeeding event has strict relation to its antecedent, and all of them vary in degree of intensity according to the locality and season—according to the age, constitution, and previous habits of life of the patient.

In the ordinary forms of remittent fever, the first stage, or that of cold, is but short, while that of reaction, or of heat, varies from two to eight hours' duration; and this again is followed by relaxation or remission, which is more or less complete, of longer or shorter duration, proportioned to the severity and duration of the previous stages. In its more concentrated forms, on the other hand, the remission of symptoms is so trifling as to require a close observation for its detection; and unless the progress of disease can be early arrested by art, paroxysm follows upon paroxysm with in-

creasing rapidity and severity, until life is destroyed.

In the milder or less dangerous forms of this fever, under the influence of proper treatment, the daily paroxysms steadily decline in force and frequency; and towards the fifth or seventh day, or earlier, convalescence may be completely established, and a return to health ensured. But however mild the fever may appear, the inexperienced practitioner should be on his guard against evil. In fact, he who has to watch the progress of this fever should be always expecting the unexpected; so suddenly and insidiously may unfavourable symptoms set in. He must be on the watch for the sudden accession of symptoms at once dangerous, unlooked for, and rapid in their progress, if allowed to go on unchecked. Changes also take place suddenly in the constitutional powers of the patient; especially in persons of weakly condition, in such as have suffered from previous disease, or from great bodily fatigue, from privation, or from mental distress.

The previous detail of symptoms will have shown how greatly the nervous and vascular functions have been disturbed, even in the course of one paroxysm, and how much the cerebral and abdominal organs have been oppressed; it is then to these two regions that the attention of the medical observer must be principally directed; for what serious results may he not anticipate from many such paroxysms, if unrestrained and unmitigated! The truth is, that, in respect of changes within the two cavities named, our circumspection cannot be too keen; for congestion, more or less severe, to end perhaps in inflammation and effusion, and consequent destruction of some organ essential to life, may be the result of carelessness, or of inattention. The most serious, rapid, and unexpected changes will be found to occur within the cerebral cavity; but indeed this latter, as well as the abdominal viscera, should be regarded throughout every stage of the fever with the most anxious attention.

Such is a detail of the symptoms and progress of the remittent fever of Bengal. In no instance perhaps will each and all the above-mentioned symptoms be presented, for the fever varies with the locality and season; but so many of the symptoms described will be found in every case as to leave no doubt whatever as to the real nature of the disease.

A consideration, however transient, of the outline here given, must satisfy any one that the disease under consideration is one which involves the safety, or the destruction of the most important organs of the body. It is likewise, varying in degree, the most prevalent form of disease known to the world, and that by which it is supposed that the greater portion of the human race is prematurely destroyed; it therefore claims a paramount attention on the part of the tropical physician, as on his intimate acquaintance with its nature and treatment will mainly depend his character for usefulness.

In placing before the reader a detail of the symptoms of the Bengal remittent fever, I have confined myself to the mention of such as are present in the ordinary and severer forms of that disease, omitting all notice of the minor forms and degrees, as these last may appear at different seasons of the year, in persons of peculiar temperaments, habits of life, or in persons of different ages and sexes.

I have done this with a view to avoid confusion, and the entanglement which results from over minute descriptions of symptoms which vary only in degree; believing too that he who can appreciate the circumstances of danger in the graver forms of fever, so as to overcome them, will be at no loss when called to treat such as are mild, and almost free from danger to life. After having been familiar with the mastering of great dangers, it is easy to bring one's mind down to considerations for the removal of the smaller ills. There is, in short, nothing in the nature or progress of dis-

ease which can disturb or embarrass the man who has been habituated successfully to encounter the storms of tropical fevers.

PATHOLOGY.-A consideration of the phenomena constituting a paroxysm of remittent fever, in a person previously healthy, and the fact that the violence of succeeding paroxysms has during life fallen with augmenting morbid force on the organs within the abdominal and cerebral cavities, will prepare the pathologist to find, that, after death, the results of congestion, or of inflammatory action, are manifested in the peritonæum, stomach, liver, and bowels in the one case, while the membranes, the cavities, or the substance of the brain suffer injury in the other case; and so it is in fact. Another circumstance deserves a preliminary and special notice, namely, the almost universal application to the fevers of the East of the significant term BILIOUS. That it has not been in the olden time, any more than in our own day, an accidental or a misapplied term is evident, for modern statistics fully attest to its pathological accuracy in the most extended and unquestionable manner. We are thus, in India, continually thrown back on the just observation of Clark, that of all the organs of the human body, the liver is the most liable to disease.

The great majority of the complications in the fevers of Bengal are abdominal, whether these consist of mere irritation, of congestion, or of congestion proceeding onwards to inflammation; and this would appear to be the principal cause of the prostration, with tendency to collapse, so common, especially during the rainy season; for, even within a few hours, there occasionally exists here an oppression of the vital functions alarming to the stranger physician.

The depression caused by a violent blow on the abdomen more nearly resembles the febrile collapse than any other morbid condition with which I am acquainted; and both probably depend on a disturbance in the functions of the organic nervous system—the powerful though silent source of many symptoms known to us only by their effects.

It must be held in recollection, too, that what are termed congestions, subacute and acute inflammations, tend, firstly, to destroy life speedily; and secondly, that, in their remoter results, they found organic diseases which have the same eventually fatal effects. In either case they form conditions of the utmost importance; for, in the first instance, the immediate cure is involved, so as to save life—in the second, the prevention of disease more or less remote and deadly.

The oppressive fulness and tenderness of the epigastric region, with nausea, vomiting, or retching, forming so urgent a feature in

the symptoms during life, are followed, in fatal cases, by alterations of structure in the mucous digestive surfaces, more or less extensive and destructive according to the intensity or the duration of

the previous fever.

These lesions will be found in the stomach, duodenum, and mesocolon principally, and they are generally of a congestive or inflammatory nature, as evidenced by a turgid or dirty red condition of the inner surfaces, and by interstitial effusions: in very severe or protracted cases the results of congestions, or of inflammatory actions, as redness, ecchymosis, or even ulceration, will extend to the mucous surfaces of the small and large intestines, while the liver, spleen, omentum, and mesentery exhibit various degrees of vascular engorgement. A severe disturbance of the hepatic function is almost universal in the progress of the remittent fevers of the East, and the evidences of congestion, sometimes of inflammatory actions, in the liver, gall-bladder, and ducts, will very generally be found after death. The spleen is not so frequently the seat of morbid changes; and when they do exist in persons previously healthy, the conditions are those of simple enlargement, or of softening of the organ.

Nor are the evidences of congestion, or of inflammatory action within the cerebral cavity, of a less prominent or declared character; and they consist occasionally of serous effusion into the ventricles or between the membranes, being the results of vascular determination passing onwards to congestion, or inflammation, and their consequences. In general, these results consist in effused serum, events sudden and dangerous in their nature, and such as we should most anxiously guard against by a prompt and effective treatment; for when effusion occurs within the head the fatal termination is too often close at hand. As already stated, however, the greater proportion of the complications, in the remittent fevers of Bengal, are abdominal, and, though hardly of less danger to the patient, they are not so sudden in their invasion, or so immediately dangerous to life, as those which are cerebral. But it is well to remember that against the one or the other, or against both together, the tropical practitioner

must be unceasingly and carefully on his guard.

The evidences that serous effusion is threatened, or taking place within the cerebral cavity will be found in mental incoherence, and a wandering indifference to objects of former interest, ending in stupor more or less complete, if the disease advances. To questions relating to his condition of health, the patient replies that he is "very well." This insensibility to, and unacquaintance with, his actually dangerous state, indicates imminent peril; and the sufferer, when not relieved by timely and proper treatment, first "sinks in bed," under the laws of gravitation, and after-

wards with dilated pupils, flushed cheeks, cold extremities, and

increasing stupor, he sinks in death.

The effusion of serum within the abdominal cavity is accompanied by increased præcordial anxiety, with increased tendency to collapse; the abdomen feels full and inelastic on examination, and sometimes there is tenderness on pressure. The tendency in the remittent fevers of Bengal to found visceral disease within the abdomen is so great that we should be on the careful watch against such complications from the first hour of treatment, until convalescence shall be established and matured.

Death from remittent fever being usually an event of speedy occurrence, there can seldom exist time for emaciation of the body; but the surface generally exhibits a sallow or greenish-yellow appearance. Where jaundice occurs in the course of this fever, and especially where it is associated with cerebral disturbance, such as drowsiness, the danger is again imminent. The collapse here has a double source—a cerebral as well as an abdominal origin; and the surface of the body after death exhibits a deeper tinge of sallowness, with livor of the face and neck.

A presentiment of death is a very unfavourable circumstance in the progress of remittent fever, and one that is too often verified in the result. A soldier will say: "You have been very kind to me, sir, but this time I shall not get over it." There may exist in the poor fellow's case nothing that has the appearance of absolute or immediate danger at this time—yet the man generally dies, and that in the course of a few paroxysms. The fatal impression is instinctive, and is not at first shared in by the medical attendant,

who is seldom able to trace the pathological source.

In July, 1840, the height of the rainy season, when, as stated, the fevers are more than usually severe in type, I passed a gentleman, the head of a mercantile firm, while driving through the streets of Calcutta. He remarked to a friend on my healthy appearance, and on the amount of labour I appeared to undergo with impunity. At 2 P. M. of that day he was seized with the remittent fever; and at 4 P. M., he was informed that, although the younger and stouter of the two, my life was despaired of by my professional friends. His countenance speedily assumed a very grave expression, and after some days he refused to use any means of cure, truly declaring himself to be a dying man. On examination after death a purulent collection was found under the descending colon, a condition not suspected during life.

THE SELECTION AND APPLICATION OF THE MEANS OF CURE.—In every form and variety of fever,—in every climate, and in every season of the year—one of the most important circumstances to observe, is, the nature of the prevailing disease,

whether endemic or epidemic; for not only do fevers differ in their nature in different localities and seasons, but so also do their complications whether congestive or inflammatory, so as to require a manifest and carefully proportionate treatment. It should likewise be impressed on all medical officers that too much attention cannot be paid to the type of every epidemic fever,—to its general

character, as well as to each individual case.

We are told by Sir James Mc'Grigor, in his admirable history of the diseases of the Peninsular army, that not only had fever very different forms in different seasons, and in different quarters of the same seasons, but they required very different, nav opposite kinds of treatment: the knowledge of this, he adds, strongly impresses on us the necessity of becoming acquainted with every attending circumstance, before we venture to censure any particular practice. These circumstances should be for ever present to the mind of the physician. And here I would remark on the serious disadvantages under which the tropical practitioner undertakes the cure of fever, as compared to his more fortunate brethren of temperate climates. The former has, in a large proportion of instances, to treat a fever violent, concentrated, and rapid in its nature and tendencies: -he has but a few days, and sometimes but a few hours—between the invasion of disease and the recovery or death of the sufferer. The means of cure too, while they require the most cool, measured and careful judgment, must be prompt, direct, and powerful to a degree not known, and therefore not required elsewhere. All this involves a most grave responsibility. In Europe, on the other hand, owing to the comparatively subdued and protracted character of the fevers generally, we can afford to watch the slow progress of events, so as cautiously and indirectly to treat symptoms, and thus aid nature in her endeavours to free herself from disease. By him who would conduct his patient with safety through the troubles and distresses of a tropical remittent fever, two preliminary conditions must be settled :- he must first select his means of cure with judgment, and apportion them carefully; and having done this, he must apply them promptly. In fever, above all diseases, we must ever admit with Trophilus that -" he who is able to distinguish what can be done, and what cannot be done, is the true physician;" but how great and difficult his task! He must measure his means to the wants of nature, -doing neither too much nor too little-carefully avoiding what ought not to be done.

Before proceeding to the cure of this formidable disease it is always necessary carefully to observe the duration of the actual stage of the fever;—that is, whether it be of some hours' or some days' duration, and whether the actually existing paroxysm be at its accession, or its decline; for, by the daily recurrence of these paroxysms, changes take place in the relative balance of power in the several functions, nervous and vascular, which render a proportionate modification in our more active means of cure necessary to successful treatment. The means which would be salutary within the first few days cannot be used later to the same effect, or to the same amount; and so again the means which would arrest fever and save life, if applied at the accession of the paroxysm, would induce a dangerous collapse, or even destroy life, if applied at the stage of its decline or towards its termination. After these preliminary cautions the reader will be prepared to enter into the general details of remedies in their usual order of application—into a consideration of their respective physiological actions—into the preferences to be given to some of them, and their application to particular cases.

EMETICS.—Certain authors in various countries have recommended that, in the treatment of fevers, we should begin with the exhibition of an emetic, and each has had his favourite-as sulphate of zinc, antimonial preparation, or ipecacuanha,—the latter being by some persons believed to be the safest. Certainly, when a patient seized with fever is found with a stomach loaded with food and drink, the sooner he is vomited the better; but the coincidence cannot be a frequent one anywhere, and within the tropics the practice of early vomiting in fever has not obtained or held ground. In the treatment of Europeans, suffering from the periodic fevers of India, I have very seldom exhibited emetics; but I am bound to say that, in the treatment of natives of all classes, civil and military, I ever found a combination of compound jalap powder with a few grains of calomel, and one or two grains of emetic tartar, to act in a manner frequently to cut short the progress of fever -the patients very generally requiring neither bark nor arsenic, the free action of both stomach and bowels, having proved curative of the fever.

In the instances in which emetics have proved of real service to European patients, I was disposed to refer the benefit generally to the sudorific and sedative effects, and not to the action of vomiting—a severe and distressing symptom already present in almost every violent case of remittent fever. There exists, in the East at least, a prejudice against the use of emetics; but I apprehend that the just application of this powerful means has not yet been found, through a careful course of observation and experience. All that seems to be really known is, that there are cases of fevers in which emetics may act beneficially, and others in which they prove injurious.

Robert Jackson says that, in periodic fever, particularly in the bilious remittent of the autumnal season, the emetic is often the first remedy prescribed; and, in certain cases, its operation is not unfrequently beneficial, so beneficial indeed in many instances that the disease is arrested, or a condition induced by its action under which it is easily arrested by other applications. This celebrated physician adds, however, that emetics do no good, but even do harm, "where the habit is full, the arterial action high, the pulse hard and tense, or small, deep and concentrated—the skin thick and torpid, or where the functions of important organs, viz., the head, lungs, or liver, are oppressed by sanguineous congestion."

Mr. Wade of Chunar, published a book, in 1791, in which he detailed the results of treatment by means of a solution of Epsom salts and emetic-tartar, with a success "unbroken, except by two cases of unfortunate termination, in the treatment of about four hundred cases of fever and dysentery." That the fevers and dysenteries amongst the old hard-drinking European pensioners forming the garrison of Chunar, in 1791, could not have been of a very formidable character, is sufficiently apparent. Mr. Wade invites "the medical world to draw their own comments;" and the medical world has done so long ago. Meanwhile, this same plan of treating every disease alike, was copied by and attempted to be practised at the General hospital of Calcutta, under the influence of bayonets, by the late Dr. Halliday; but it had no better success this time with "the medical world." As to the patients in this latter instance, no power could induce the Europeans to submit, either to the continued nausea, the frequent vomitings, or the incessant purgings which resulted. There was, in fact, no power at that time in the government of British India sufficient to enforce the emetic-cathartic treatment.

This is the kind of *case-book* that is calculated to do so much harm with the inexperienced and the inattentive, who are averse from the trouble of learning their duties, except by the shortest and easiest of ways.

THE WARM BATH—TEPID AND COLD AFFUSIONS—COLD DRINKS.—Robert Jackson began his treatment of the congestive fevers of the West Indies by a warm bath and active frictions to the whole surface of the body. While thus immersed, with the sensibility of the skin "augmented by the heat of the bath and by friction," and with the fluids powerfully derived to the surface, he abstracted blood from the arm. That such means are powerful and require the hand of a master, no one can doubt;—but, in point of fact, neither the warm bath nor the tepid and cold affusions have had, any more than the emetics, any systematic or sustained support from the practitioners of the East. Perhaps the great tendency to collapse in the fevers of Bengal may have at all times deterred us from the very liberal use of means which tend powerfully to affect the balance of circulation.

I remember the instance of an European who was employed on the deadly island of Sagor, and whose case interested me greatly at the time. His fever was in its seventh day when I saw him, and the native boatmen, to prevent his leaping overboard, had tied him in his delirium to the benches, in which state he was exposed, bareheaded and half naked, to the direct rays of a March sun during the two days of his passage up the river to Calcutta. The whole surface of the body was of a lurid lead-and-brick hue, as were the eyes. His delirium was subsiding into stupor, no coherent reply being obtained from him; his restlessness was of the most distressing kind, for he tossed and changed his posture violently in every few seconds of time; his tongue was black and dry, and his pulse was extinguished by the weight of my fore-finger:the case looked hopeless. I seized, however, on what appeared to me the most urgent feature of this poor fellow's case, namely, the most excessively pungent and unpleasant heat of skin I had ever felt by means of the touch; and I ordered him at once, and under my inspection, to be dashed with cold water over head and trunk, from a few feet of height. Some six or eight earthen pitchers of water were thus poured on the patient, after which he was well rubbed and put in bed. The fearful restlessness at once subsided, and in half an hour he was in a gentle perspiration and sound sleep. I have never seen such a case, or such a result before or since. A few mercurial purgatives concluded this man's cure, and I did not see him until two years subsequently. But it now became apparent how severe had been his fever, for the faculties of the mind never regained their former power, and he was eventually sent to his friends in England. The unusual nature and amount of heat in this case induced me at once to apply the cold bath; and I was of opinion at the time that, had I adopted any other course of treatment, the patient must have died. But it is only fair to state that Robert Jackson, the originator of this plan of treating fevers (and not Dr. Currie, of Liverpool), and who had had more experience of its effects than any other person since his time, declares that the salutary action of cold affusion, in febrile diseases, is not necessarily and indispensably connected with the presence of increased heat. His caution that cold affusion is "ineffectual or dangerous where deep congestion or strong inflammation exists in any of the interior organs," may constitute another reason why this means has not been more generally had recourse to in the fevers of the East Indies. Measures difficult of application and sometimes doubtful of result—excepting when used with an extraordinary tact—like Robert Jackson's, will not be generally adopted or employed by ordinary medical practitioners.

The topical use of cold and tepid water is much practised in India, and that with signal benefit when applied to the scalp, fore-

head, and temples, where cerebral determinations occur with severity, as in the ardent fever of the hot season.

Warm baths have been limited, in my own practice, to the latter stages of fever, dysentery, and hepatitis, and when convalescence was approaching. They will then be found both grateful and beneficial.

Nature prompts the free use of cold drinks in all forms of fever, and the extinction of thirst is everywhere esteemed a powerful means of cure; so much so that certain mild forms of fever are curable by cold water alone. The abundant introduction of American ice into the three capitals of India has proved of great service in the treatment of the fevers of those cities. The ice is useful not only as a local and general refrigerant, but as a powerful means also for allaying the distressing irritability of stomach so common to our fevers.

BLOOD-LETTING.—I come now to consider blood-letting, the most powerful remedy, the most difficult of just application, but the most needful of any we possess, if we would cure the remittent fever of Bengal, and save the patient from subsequent visceral disease. The question of blood-letting in disease is almost the most important which can engage the attention of the practical physician. I am aware that a kind of fashion is at present moving against the practice, almost to the desire for its total exclusion; but those who are acquainted with the history of medicine, and who have seen and understood the nature of acute tropical disease, will take fashion here at its proper value, and at no more. "It may be very well," says a Dublin critic, "to refrain from blood-letting in the treatment of feeble and unsound patients in large cities, but what will our provincial friends say to the wholesale interdiction of the lancet? That we do not now bleed as we formerly did is obvious enough; but do we err in the opposite extreme? We have ourselves 'a theory,' that it might be better at once to diminish the quantity of the circulating fluid than to render it unfit to support life or repair injury, by slops and physic. By the way, if it is so bad a practice to bleed, how does it happen that we have so much leeching and cupping, especially in private practice? These questions are worthy of consideration, both in a pathological and medico-ethical point of view." Certainly, these questions are worthy of consideration; and, to borrow their own language, how long will medical men persist in thinking that if any given proceeding be wrong, its directly opposite must necessarily be right; -that if profuse and indiscriminate bleeding, for instance, be wrong, the not bleeding in any case must be right?

One very important circumstance should be impressed on the attention of all young naval and military surgeons, viz., that when,

upon careful consideration, general blood-letting is deemed necessary in the severer forms of tropical fevers,—when there is violent tumult of arterial action, or an oppressive venous congestion,—conditions alike dangerous to life—the said general blood-letting, to be effective, should be practised as early as possible, and be sufficient in amount to make an impression on the febrile movement. A small bleeding, or repeated small bleedings, may moderate a

mild, but can never arrest a violent tropical fever.

In the practice of Robert Jackson, in the West Indies, the largest bleedings, amounting in some cases to "ten pounds in twenty-four hours," were made chiefly in the ardent and inflammatory cerebral fevers, and occasionally in the congestive abdominal complications common to the West as to the East Indies. Of the complications referable to the cavities of the abdomen and cerebrum, the latter is the more ardent, and the more immediately dangerous to life. It is but proper to add, that "this apparently revolting practice," as Jackson terms it, was of such effect in his hands "that the greater number of persons who were treated in this manner returned to their duty within a fortnight." As here stated, the rule of depletion that applies to an acute or ardent fever, applies with equal force to the same means of cure in the more difficult form of congestive fever; and there is not in the whole range of practice of medicine any question of a more nice discrimination than that of the measure here referred to; and next to that important question comes that of the proper time of using blood-letting. To render this important operation "the more certain, and the more perfect," Jackson urges:-

1. "That it be made within six hours from the invasion of continued fever, or before the paroxysm of the periodic fever of vio-

lent excitement has attained its acme.

2. "The simple act of abstracting blood from the circulating system is often decisively effectual in arresting the course of fever, where it is resorted to at an early period, and where the process is

properly conducted.

- 3. "It is still salutary, but of less decisive effects, where the course is more advanced, that is, beyond the third day. It is not of dependence, but is not prohibited, and it is occasionally useful even at late periods. It is thus not safe to carry it to the extent of effecting precipitate arrest; it is safe, as well managed, to alleviate impending dangers, and often to facilitate the development of regular crisis:" thus he conceives that guarded blood-letting favours the tendency of fevers to subside under the law of periodic action.
- 4. "If employed thus it obviates or removes congestion, and thereby prevents effusion into internal cavities, and into the substance of internal organs.

5. "Abstraction of blood is, moreover, safe, and its effect is

often important, at the first moments of relapse.

6. "Prescribed with consideration, and applied with management in execution, blood-letting is both a safe and powerful remedy, either decisive of cure by its own power, or preparatory of the curative powers of others. It is the main engine of successful practice; it is not the sole remedy.

7. "But though I say this in truth, I do not say that bleeding in a large, even in any quantity, is uniformly proper, or uniformly

safe.

8. "Though the abstraction of blood is generally beneficial, it is only so conditionally, and under management. If the effect produced by it be not seconded by well-considered means of stimulation, or conservative of auxiliary power, general or local, adapted to the circumstances of the case, the chances are that the harm will be greater than the good. But while blood-letting is a remedy of nice management in cases of relapse, it is to be avoided, as well as other remedies of strong operation, where there exist appearances of approaching a favourable crisis. On the contrary, where the critical power labours, and the critical effect is marred through internal impediment, the abstraction of blood in a given quantity is often followed by signal benefit; it gives facility to the course of the salutary process artificially obstructed.

9. "If there be no prohibitory circumstance in the case, one

bleeding is preferable to repeated small bleedings.

10. "But though I regard the abstraction of blood as a remedy of the first importance, for the abrupt and successful cure of fever, either primarily or secondarily, I am yet free to own that its good effect depends principally on the manner of adjusting the subtraction to the condition.

11. "It often fails, and even sometimes does harm where, employed as principal, its real place is only that of auxiliary; and, per contra, where employed as auxiliary, its real place is that of

principal."

Not to speak of the practical importance of the rules here laid down by our greatest military physician, and numbered by me the more to mark their separate import, they are of extreme interest in the history of tropical fevers, and of their treatment; and it is principally on this latter account, and not as absolute rules of practice in India, that they are here presented to the reader's consideration. Such histories I conceive to be of great importance, as from their attentive consideration and comparison we learn, generally after a lapse of years, to deduce from them a just balance in the use of all the known means of cure, old as well as new. In the principles involved in the rules above quoted from the writings of Robert Jackson, he will, I think, be found just, although the

degree and extent to which he carried blood-letting will not apply to the fevers of our day, in the East Indies at least. The rational and the timely employment of blood-letting are the two great points for consideration and determination by the tropical

physician.

It has been well observed that the most distinguished physicians of all ages, who have treated of fever, coincide entirely in the feeling that with regard to this important class of disease it is impossible, in the short life allotted to the most aged, to do anything more than add a little knowledge to the common stock. If this be true, and who that knows anything of medicine can doubt it, we should study with care the history of fever, and learn from it not to boast of our success, or blame those who have gone before us; for in truth we are too often only able to see the defects of the older writers by the lights which they have placed before us. The accurate observer Tennant, speaking of the treatment of the fevers of his time in Bengal, 1796, says that it was "simple and decisive, and more efficacious than that of any sort of diseases

equally malignant in Britain."

Jackson designed to produce immediate curative result by cutting short the progress of fever, and so did the older medical officers in our tropical possessions by "throwing in the bark." In the circle of varying opinion, doctrine, and practice, we come again, after the round of a century, to the same exclusive treatment last mentioned, in heroic doses of quinine. This plan of arresting fever has been of late had recourse to in the continued types of Europe, and in the formidable paroxysmal malarious fevers of tropical regions, but as yet without any determinate results. The treatment by Peruvian bark, and by bark chiefly, was long and faithfully practised in former times, and an enormous amount of visceral disease, the infarctions of the older writers, was the result; and it is to be feared that, in hot climates at least, such will again be the result, whether we employ, after an exclusive manner, the disulphate of quina, or the more galenical preparations of the bark. All experience points, in fact, to the union of the depletory and refrigerant with the antiperiodic means in the cure of tropical fevers, remittent and intermittent, thus blending all the safe and effective means of cure, not limiting ourselves to one exclusively.

In the treatment of fevers, as of other acute diseases, we must be content with what can be done with safety and efficiency. As matter of theory, or even of reasonable endeavour, it seems sound in principle to cut short a fever in its career, however difficult the execution; for, as in military affairs, what can be done most speedily is done most effectively, and we should bring into action all our forces as soon as possible. As yet, however, we

have nowhere succeeded in absolutely cutting fever short by coupde-main, either by blood-letting or by the Peruvian bark; yet the two means constitute beyond doubt our heroic remedies. Robert Jackson was supposed to have come nearer the desired result than any physician with whose practice we are acquainted; but it is admitted also that, in hands less skilled than his, the practice of large and repeated bleedings was not always safe. In the East Indies they have never been used, nor would they there be justifiable as a general rule. No tropical practitioner can peruse the works of Robert Jackson, however, without great benefit; but it is to those practising in the West Indies that, with a certain reservation, they

will prove of the most special and signal use.

All circumstances considered, I give a very decided preference to general over local blood-letting in all cases of acute and aggravated tropical disease, whether it be the remittent fever, Dysentery, or Hepatitis. When the medical attendant bleeds from the arm he sees and knows what he is doing; he measures his means and adapts them to the circumstances of the case, and he sees carefully, and at once, that it is not carried too far; whereas, by the application of leeches, no one knows the exact amount of blood abstracted in the first instance, and the amount of subsequent oozing, in the absence of the medical attendant, is even more uncertain, and in remittent fever even more dangerous when, as occasionally happens, this oozing takes place during the night, and in the declining stage of remission. This much is certain, however, that so much blood does occasionally ooze from leech-bites as to cause an immediate prostration of the powers of life amounting to dangerous collapse, with a subsequent anæmia very difficult of cure. Of these facts various emphatic examples are daily presented to me in the instances of invalids returning from India, and the proofs will be furnished in a subsequent portion of this work. From the recorded histories of the cases it is often evident that, where the medical attendant in India intended to abstract the amount of twenty ounces, sixty ounces have been lost, or three times as much as was intended. Now, no medical officer of ordinary education, however young, could look on venesection, and see his patient approach to a sunken condition without immediately tying up the arm; but where leeches have been applied I see very lamentable results from neglect and oozing, the patients being at the time under the general care of very able and experienced medical officers. The application of leeches is too often left to relatives and to servants, a contingency which cannot in the nature of things happen in venesection. When I ordered the application of leeches in the fevers of Bengal, I invariably sent a native dresser to apply the requisite number, with directions to dress the leech-bites so as to preclude any

oozing. I would strongly recommend, therefore, that whenever it is proposed to abstract by leeches so much blood as twelve ounces or more, the operation be performed by the lancet instead.

Fevers, idiopathic and symptomatic, are traced by Dr. Billing " to a loss of the functions of the nervous centres, and subsequently of the organs depending on them." Fever, he says, results " from injury to the centres of the nervous system, which arises either from peripheral injury propagated to them, or through lesion by miasm, which, by the route of the circulation, directly poisons them-most probably by chemical combination or alteration-instantaneously lowering their power or energy. I have shown throughout, that the immediate effect of the lowering of the power and energy of the nerves or the nervous system is inflammation, or congestion of the capillaries, the first degree of inflammation." Referring to the treatment, he says :- "If we can relieve his debility of body and mind, the patient will recover from his fever; that debility we have shown to be caused by an overloaded state of the nervous centres. We uniformly see that the only successful means of relieving them consist in diminishing the injecting force when the pulse is strong, increasing at the same time the contracting action of the capillaries by antimony, mercury, salines, bark, &c. &c., or even when the pulse is not strong, we find bleeding sometimes necessary to diminish the actual quantity of the load, as the constringents alone may not be sufficient to produce contraction, so that, with reference to the indication of bleeding, we have much more to consider the state of plethora of the internal vessels than the state of the pulse or vis a tergo. And though emetic substances have an influence similar to that of bleeding in lowering the vis a tergo they are, nevertheless, of most essential advantage when the pulse is even almost gone, by their immediate constringent effect on the internal capillaries: hence it is evident that the pulse, which was so long considered as the indication for the use of bleeding or sedative medicines, is often no guide at all, in which cases the necessary practice of sedatives, from having been hitherto unexplained, has always been called indirect practice. This distinguished physician adds, "I do not admit this term, and never practise indirectly; my indications are always founded upon physiology, as I have explained them up to this point." Principles of Medicine. The doctrines here propounded appear to have anticipated that of Virchow, who refers the primary phenomena of fever to "loss of power in the nerves;" and likewise that of Traube, who ascribes many of the symptoms of fever to "a weakened, more or less paralytic condition of the vagus."-Brit. and For. M. C. Review, for April, 1855.

PURGATIVES.—When we reflect on the intimate pathological

connection which exists between the stomach, small intestines, and liver, and on the facility with which morbid irritations of the former organs are conveyed to the latter, when to these circumstances we add the frequency with which the liver is congested or inflamed in the course of the febrile movement, we shall be at no loss to perceive how it is that purgatives have at all times been held of high repute in the treatment of the remittent fevers of Bengal. When we regard the intimate nervous and vascular inter-communications of every organ within the abdomen, the one with the other, and that each and all of them are disturbed in their functions during this fever, while some of the organs are more liable than others to become congested or inflamed during its progress, we shall be prepared to find that purgatives rank as powerful means in aid of the cure, and such they have always

very justly been held to be.

Given in their uncombined forms, and with reference to their special and peculiar actions, they remove retained or vitiated matters from the bowels, thus carrying out of the system sources of contamination, while they promote a more free arterial and venous circulation; they excite the secreting and excreting functions of the bowels, and by these several actions they produce a sedative effect, while they derive from the head so as to relieve that cavity, as well as the abdomen, from an oppressive load of congestion. Combined with mercurials they act as cholagogues, and thus relieve the portal circulation from stagnation, and the liver from consequent disease. They rouse the dormant sensibilities of the bowels, and aid in restoring the balance of the general circulation, especially when blood-letting has preceded their administration so as to remove congestions, and thus afford a renewed activity to the functions of secretion. When we consider the certain action of purgative medicines, and that their influence is exerted on the most extensive secreting surface of the human body, we shall at once perceive how powerful must be their effects. By previous depletion the system is, in fact, prepared for the more free action of this class of remedies, whether exhibited in their simple or combined forms. The compound extract of colocynth, with calomel and James's powder, followed in a few hours by infusion of senna with sulphate of magnesia, or by the compound powder of jalap, is a favourite mode of purgation in this fever. It is usual also to add the sulphate of quina to purging mixtures, so that the tonic and antiperiodic may thus be applied to the whole mucous digestive surface. I found this last an excellent remedy, both in hospital and private practice. Where, in the latter stages of fever, the bowels are torpid and the hypochondria inflated, Robert Jackson recommends a tineture composed of aloes and myrrh, and also the oil of turpentine. He

is partial to the use of the tineture; but the oil of turpentine, he says, while purgative, stimulates the whole alimentary system to the proper exercise of its functions.

DIAPHORETICS.—If the main cure of fever, as stated by Robert Jackson, consists "in turning the tide of circulation to the exterior, in maintaining it there, and thereby disembarrassing the interior organs," the value of diaphoretics must be great. In the very milder forms of fever a few doses of purgative medicine, with or without mercurials, as may be, and followed by sudorifics, given so that their action shall be maintained, and by diluents, will cure the disease; but in the more grave cases here under consideration, blood-letting and purging must precede and give effect to the sedative action of diaphoretics by their relaxing effects on the system, and by their reduction of its unnatural heat. As in the instance of purgatives so with sudorifics—the system must be prepared for their full effects by other and more powerful means.

Minute and repeated doses of tartar emetic have always appeared to me to constitute the best diaphoretics, given simply; but when mercury in any form was added, I then used James's

powder in preference.

"All the metallic salts," says Dr. Billing, "have, more or less, an astringent effect on the capillaries; and to this influence I attribute the universal efficacy of antimony as an antiphlogistic remedy, it being doubly valuable in acute cases from its sedative effects on the heart and pulse, combined with its locally tonic or astringent effects on the capillaries of inflamed or congested parts, as well as on those of all the secreting structures. Hence, too, its efficacy, in small repeated doses, in cases where there is great depression of the system, by its relieving the relaxation of the capillaries by which the depression is caused. Nay more, we can manage to insure its full antiphlogistic effects without the inconvenience of nausea, by combining it with a little opiate and aromatic."

MERCURY.—Second generally in time of application, and only second in point of importance, comes the action of calomel. When blood-letting shall have produced its results on the circulation, the febrile movement, though subdued, being still in serious progress, we have recourse to this mineral. It is at once the best aid to the abstraction of blood, and the best preparation for the effective action of antiperiodics, such as quinine, arsenic, &c., through its active operation on all the depurative functions. Robert Jackson, though no favourer of mercurial treatment, admits that in fevers, complicated with abdominal congestion, calo-

mel is a remedy of the first importance. He says that it repairs mischiefs which no other means with which we are acquainted are capable of touching. This is the simple truth, and it constitutes the reason why this remedy has at all times held so high a

place with the practitioners in our worst climates.

As it is better to repeat than to leave an important subject in any doubt, I here beg leave to transcribe from my article on Hepatitis a summary of the actions of mercury, and especially of those of calomel: -It is for the very reason that calomel assists powerfully, both in "drawing off" accumulations, and in promoting "increased secretion" that it proves of such value in aid of blood-letting. It is, in fact, by this very double action of purging and increasing secretion at the same time, that calomel relieves the loaded and inactive vessels of the diseased gland; not to speak of the other acknowledged physiological influences of this mineral-such as its increase of all the secretions and excretions of the body-its influence on the capillary circulation -its febrifuge effect-the peculiar specific power ascribed to it by Gooch and others as an antagonist to inflammations, whether general or local—its stimulant power over the absorbent function -its power of unloading, at the same time that it gives a new impulse to the vascular system—its peculiar power in removing viscid and tenacious intestinal secretions-its alterative, solvent, and antiplastic effects on the blood;—these are the actions and uses ascribed to mercury by the ablest of our physicians and surgeons, and they are such as place this remedy second only in order and in importance to blood-letting, in all the more acute hepatic affections of India. That mercury enters into intimate union with the elements of the blood is now an ascertained fact. It must, therefore, "modify its plasticity, and influence all the organic functions to which it is subservient." I have only to add here that the above summary is quoted from the former edition of this work, with commendation, by Dr. Copland, in his article on Diseases of the Liver.

Dr. Billing, speaking of the treatment of disease by "calomel and opium," says justly, that though powerful allies to the anti-phlogistic treatment, they have often failed from being employed without being supported by bleeding, or purgative, or emetic, or diaphoretic medicines. This is equally true of tropical diseases, and I have seen enormous quantities of calomel given with nothing but evil effect, because its action was not solicited and supported by previous depletory means.

Robert Jackson states, that the preparation of the subject by previous abstraction of blood is indispensable to the success of mercury, in the more concentrated fevers and more complicated

dysenteries of tropical climates.

It is thus, by regarding each means of cure as constituting a

part only of a whole, that we shall best avoid the errors and dangers of relying upon exclusive means in the treatment of dangerous and complicated diseases, and that we shall learn to

apportion to each remedy its just value.

BARK.—Sir James Annesley observes, that visceral disease is the necessary result of the neglect of depletory means, including mercurial purgatives; and that congestions often lead to inflammatory action, especially if bark or arsenic have been freely administered during the stage of congestion, and of disturbed and impeded secretion; and this is in accordance with general experience. Here again we perceive that quina, like other powerful remedies, may not only prove of no effect, but may be administered so as actually to be injurious. It may do signal good, or even prove hurtful, according as the patient may or may not have been prepared for the actions of the bark. To apportion the dose and determine the right time for exhibiting it, so as to secure the full tonic and antiperiodic influences of the quina, we must appreciate accurately the circumstances in each individual case, the stage of the fever, and the effect of the previous antiphlogistic means. The random exhibition of quina by timid, careless, or inexperienced practitioners does infinite harm; and, in former times, when bark alone was relied upon for the cure of

all tropical fevers, the results were horrible.

Where, with a state of general plethora, visceral congestions remain unsubdued, or only partially removed, the secretions being scanty and depraved, with the pulse full and hard, and the skin constricted, dry, and hot, the time for the exhibition of bark has not yet arrived. Robert Jackson says, that bark is not to be relied on for the precipitate arrest of remittent fever, either in the West Indies or in other countries. Bark and the disulphate of quina are efficacious in arresting the remittent fever of Bengal when venous congestions have been overcome by previous depletory means, when the pulse has been reduced in force and frequency, when the secretions are in free action, and the skin relaxed; when these preparatory results have been obtained, we may then be sure of establishing the antiperiodic influence of quina, with the best effect, and without risk of producing any injury. It then becomes a sovereign remedy; and the nearer the patient can be brought to the state of remission, the surer will be the operation of the antiperiodic. It must now be given in full doses, so as speedily to establish its influence. The amount of dose, and the frequency with which it is to be repeated, will depend on the nature of the fever; -where paroxysms are violent, or where the sufferer is in a malarious locality, the doses should be large, and often repeated. It is better to exhibit the quinine in five grain doses, often repeated, than to give scruple doses as recommended by some writers; for by the former plan we have the advantage

of observing the effects as we proceed, and we can withhold excessive doses.

"Quinine," says Dr. Bence Jones, "acts in weakening the pulse in a similar way to arsenic, hydrocyanic acid, opium, and other narcotics, antimony, and nitre."

Quoting the general conclusions of M. Briquet, he adds:-

"1. That the maximum diminution of the pulse is rarely twenty to twenty-five pulsations a minute, even in typhoid fever.

"2. That the diminution is always in direct relation to the pre-

vious frequency of the pulse.

"3. That the reduction is never below forty beats a minute.

"4. That much fibrin in the blood, or active inflammation, pre-

vents the depression.

"5. That large doses of quinine produce so serious a perturbation of the economy that they should not be given unless the illness, as regards length, seriousness, and accidents, is sufficiently important."

With regard to the pathological action of quinine, M. Briquet

considers :-

"1. That it does not act directly on the marsh poison.

"2. That it does not act on the general state of the organs, or on the blood; but it has especial action on the nervous system—C'est la médicine du quitte au double, celle que les militaires emploient quand pour couper la fièvre ils avalent une double ration d'eau de vie chaude mélé de poudre à canon.

" 3. It does not act by increasing the vital forces or by sustain-

ing them.

"4. It is not by a tonic action, or an astringent action, or a stimulant action, but by a sedative stupifiant action that it acts on the marsh poison.

"Lastly. The absorption of quinine by a sound skin is very

doubtful."

ARSENIC.—This mineral acts as a powerful tonic and antiperiodic, ranking in these qualities next to the cinchona. Arsenic has been used from time immemorial in Hindustan by the native doctors; and I saw at the Native Hospital of Calcutta daily examples of permanent injury to the circulating system, and to the mucous digestive organs from the abuse of this mineral by the Bazar practitioners, in their endeavours to cure fevers, rheumatisms, and venereal diseases. "Arsenic," says Dr. Billing, "often acts as equivalent to a union of bark and mercury; for arsenic, besides its tonic effect on the nervous system, increases the secretion of bile, and otherwise acts on the liver; it possesses also the power, like mercury, of curing chronic inflammations; and even further resembles it in occasionally producing the inconvenience of salivation."

We should commence the use of this mineral in small doses, and not persist many days in its exhibition; in fever not beyond eight or ten days. If, besides its tonic and antiperiodic effects, it acts "on the liver as much as mercury," according to the lastnamed authority, we should hold this remedy in more regard in India than we now do. It is one that may safely be administered during the paroxysm of fever; and the number of days stated are sufficient to test its virtues. In the practice of the medical officers in the East Indies, in so far as the fevers of the country are concerned, the exhibition of the arsenical preparations is almost exclusively confined by them to the intermittent forms; and, even in these, arsenic is far from being in general use.

WINE.—In the treatment of the ordinary remittent fevers of Bengal wine has little to do; for unless a tendency to collapse occur, with profuse sweats, an anxious countenance, a laborious respiration, a cold damp skin, and sunken pulse, we have not to exhibit spirituous or vinous stimulants. On the occurrence of collapse, however, diffusible stimulants are invaluable, and they are always resorted to. In cases where the cure is delayed for many days, either from neglect of early treatment, or from constitutional peculiarity, wine is occasionally requisite. But whether in recent or in protracted cases, the tendency to sinking is the event to be carefully watched; and then the wine should be given, with or without farinaceous food, according to circumstances, and with quina in the intervals. Wine and spirits must be regarded as simple stimulants; they do not change the course of events in periodic fevers. This is so much the case that, after the use of wine, we may in the next coming paroxysms have to use antiphlogistic means:-such are some of the difficulties to be encountered in the management of this formidable disease.

In some forms of continued fever, and in certain malignant periodic fevers, Robert Jackson admits of the use of wine; but he says that the application must be nicely adjusted. Champagne he considers a good wine for the occasion; and porter, he adds, is relished by the British soldier.

OPIUM.—Opium in stimulating doses I have seen prove of great value, under two circumstances not uncommon in the progress of remittent fever. The first is, the disposition to faintness or to profuse perspiration after blood-letting. Here half a grain or more of opium with calomel, or tincture of opium, with chloric ether, given with the calomel, will relieve from vascular and nervous depression, and rouse and restore the patient. The other occasion is when, after the use of depletory means, the hard-living patient, or the man habituated to the free use of various stimulants, begins to feel the depressing effects of fever, and of active

treatment, and of the absence of the accustomed exciting beverages;—then it is that symptoms of nervous exhaustion come to be complicated with the existing fever, so as frequently to constitute a masked or subdued delirium tremens. There is here great danger, in the hands of the inexperienced or the inattentive. Depletory measures would now destroy the patient; whereas opium, applied in a moderate dose, at the right time, calms while it restores the sunken nervous functions, and the patient is saved; under ignorance or neglect he is lost. Given in the manner proposed, and for the occasion only, conjoined with calomel, the opium imparts its influence to the nervous system, without locking up the secretions—an important reserve in the case before us.

## APPLICATION OF REMEDIAL MEANS TO THE INDIVIDUAL CASE.

After the general view of the various therapeutic agents employed in the treatment of fever, and of their effects, we come to consider their joint and several applications to the individual patient, suffering from the remittent fever of Bengal; and here we find, on experience, that the first and most immediate object is, to reduce the force and frequency of arterial action during the paroxysm, which if allowed to go on unrestrained, would injure

or destroy some organ essential to life.

If the patient be seen in the forenoon, on the accession of the first, second, or third paroxysm of ordinary remittent fever, if he is of a sound constitution and not beyond middle life, blood-letting from the arm, while the patient is in the recumbent posture, should be practised to the extent of relieving the sufferer from præcordial oppression, from visceral fulness and congestion, or from the intensity of headache, whichever may predominate. If along with reduced force and frequency of the pulse, and reduction of the morbid temperature, we obtain from the operation a gentle relaxation of the skin, we have the best evidence of relief from visceral congestion, whether the operation be performed for the cure of fever, dysentery, or hepatitis; the quantity of blood abstracted being regulated by the effect, and not by an arbitrary measure in ounces. It will sometimes happen, however, from peculiarity of habit, or other cause, and notwithstanding the utmost circumspection, that the relaxation of the skin will proceed to sweating, with symptoms of depression of the vital powers; then from half a grain to a grain of opium, or from fifteen to twenty minims of laudanum, with as many of chloric ether, should be administered. This will impart tone to the heart's action, and soothe the nervous excitement, while it will allay gastric and intestinal irritation. It is only in cases of depression, however, such as this, that opium is to be recommended in the very early treatment of fever; but, when requisite, it will be found to calm both mind and body. Let the physiology of the disease, and the habit and condition of the patient guide the application of remedies, and we shall approach as nearly to the correct measure of means of

cure as human endeavour can compass.

Unless the fever assume a severe form, one general bloodletting, practised as stated, in the recumbent posture, and under the studied observation of effect will, on the average, be found sufficient to relieve the patient from abdominal or cerebral oppression. Blood-letting, used as here directed, will be found to simplify the application of all the subsequent means of cure. Within an hour after the bleeding a dose of calomel with compound extract of colocynth and James's powder should be exhibited, followed in two hours by a powerful cathartic, such as infusion of senna with sulphate of magnesia. After the free action of these remedies we shall obtain in the afternoon some degree of remission; and the patient should be directed to take at bedtime from six to ten grains of calomel, with four of James's powder, if the skin be dry. Here, within eight or ten hours of first seeing the patient, we perceive that of the means above generally described, blood-letting, mercury, a sudorific, and a purgative, have been used, while the patient has been allowed free recourse to cooling diluents.

On the early morning visit of the following day the patient will probably be found in a more complete state of remission, when the sulphate of quina alone, or in combination with the purging mixture, should be freely and repeatedly administered;—given in this latter manner the quina applies itself to the whole extent of the mucous digestive surface, so as to give full effect to its tonic and antiperiodic influences; and a larger dose of it can thus be borne in the early stage of fever than if given without the purgative. This is an important consideration; for the quina rightly administered will arrest the progress of a mild fever, and save the life of the patient in a severe one, by mitigating the

symptoms.

By the forenoon the paroxysm may recur in a milder degree, though to such an extent as to demand the application of leeches to the epigastric region, if any oppression or fulness exist there, or behind the ears if there be headache; while a mixture composed of antimonial wine with the acetate or nitrate of potash should be given every two hours, so as to soften the skin and determine increased action of the kidneys. It is usual in the remittent fevers of Bengal to give from five to ten grains of calomel, with or without antimonial powder, at bed-time, followed in the

early morning by an active purge, and when the last is under operation quinine in solution is given freely. This is a good practice, and a sufficient means in itself to cure some of the milder forms of the fever under consideration;—preceded by an adequate abstraction of blood, this practice will even cure a severe fever. Thus, under the favourable circumstances contemplated, we witness the daily decline of the disease, and the daily diminishing occasion, therefore, for the use of active measures of cure, until towards the third, fifth, seventh, or ninth day convalescence is

completely established.

If remittent fever has existed unrestrained, however, for several days, and the patient is not seen till the accession of the third or fourth paroxysm, or even later, provided the general powers of the constitution remain uninjured, a general blood-letting is still the principal means to save life, followed by calomel, purgatives, and quinine, in the manner previously indicated. But if, on the other hand, the duration of the disease being as above, the paroxysms have become indistinct, running into each other with but brief or ill-defined intervals, while abdominal or cerebral complications arise, as indicated by epigastric fulness with anguish, or by approaching stupor or delirium, congestion now wearing the aspect of inflammation; then the time and manner of applying our more active means demand the nicest care to insure, not only their just effects, but their safety. Blood-letting may even now constitute the principal means to save life, but the blood must be guardedly abstracted, whether generally or locally; and calomel becomes indeed a remedy of necessity, as Robert Jackson would rather refuse to call it, in the treatment of the fevers of the West. To save the patient from impending dangers, in the case of unrestrained fever here contemplated, the medical attendant must be neither rash nor supine: sinking would result from the rash application of means, and effusion into one or both cavities involved in those fevers would be the result of timidity or indecision. Generally speaking the blood must here be abstracted by leeches at the accession of the paroxysm; antimonials must be usedcold must be applied to the shaved head-sinapisms and blisters must be applied; but on the influence of calomel chief reliance must be placed. It must be given every three or four hours. with an occasional mild aperient in the intervals, until the dangerous symptoms shall have yielded; and this favourable state is often observed to be coincident with the mercurial influence, as evidenced in the odour of the breath, or on the gums. Dangerous symptoms such as are here described will sometimes arise suddenly, without any loss of time, or without any neglect in the treatment; and when such conditions are associated with yellowness of the skin, in persons broken in health, or of a feeble constitution, or of dissipated habits of life, or who may have undergone much mental distress, the chances of a fatal termination are imminent.

In the remittent as in all forms of periodic fever, the stages of danger are those of the accession and of the decline of the paroxysm: in the first, the violence of arterial action may proceed at once, unless restrained by treatment, to destroy life by serous effusion; while, in the second, so great a prostration of the vital powers may succeed to the previous tumult of vascular action, as to terminate in a feeble and irregular pulse, a damp coldness of the surface and of the extremities, despair, and death. I have seen one paroxysm of Bengal remittent fever, and of jungle fevers. cause death in both ways, within twelve hours; and the first instance, attended with very painful circumstances, occurred within a few months of my arrival in India. A young officer was seized with fever at 11 A. M.; but nevertheless, he went into the China bazar of Calcutta. He returned to his quarters in Fort William at 3 P. M., and placed himself under the care of a staff surgeon there. He was bled at 4, and he was dead at 7 P.M. The same operation, practised at noon, would probably have saved his life.

When the spleen is affected with enlargement, either of the acute or chronic nature, mercury in all forms had better be avoided in the treatment of the fever; for there is in the splenic complication a dissolved condition of the blood, with general cachexia, that should preclude the use of this powerful mineral. Bloodletting, general or local, when found necessary, should be cautiously used, and carefully regulated in these cases; and the oozing from leech-bites should be carefully and promptly arrested.

It is necessary to be on our guard against irregularities in the paroxysms of remittent fever, as they occasionally vary not only in their time of accession, but there may arise a double or anticipating night paroxysm, in addition to that of the forenoon. This will require a double attention, especially in respect of the time of

using remedies.

A close observation is indeed necessary as to every event and circumstance which may arise in the course of these fevers; for upon the amount and accuracy of such knowledge will depend not only the selection and application of our means of cure, but

the affixing of the proper time of using them—often a vital question.

The result of each paroxysm on the cerebral and abdominal organs should also be carefully noted, for by this knowledge we regulate both the force of our remedial means, and the frequency with which their application may be needed. Through such observations also we come to estimate aright the powers and capabilities of the patient; for the greater have been the numbers of paroxysms, the more the viscera are likely to be oppressed and

congested, and the less power consequently do we find in the patient either to sustain him under further invasions of fever, or to bear him well through the operations of the necessary remedies. In such a case there will be a tendency to exhaustion, and, while we act with decision and calmness, we must be doubly watchful of effect. I have preferred a simple narrative of the treatment of remittent fever, such as is here presented, to a detail of hospital or other cases; for I have seen that, in the treatment of fevers and other acute diseases, cases are too often seized upon and followed out to the letter, by young medical officers, to the disregard of the differences of circumstances, and to the exclusion of reason and reflection on the cases before them. Cases thus become a kind of pattern in routine, and the young naval and military surgeon is injured where it was intended to afford him help. He is thrown off his reasoning powers by an array of cases, and enticed into a groove of routine, out of which it will require much exertion and firmness on his own part to extricate him. A case-book thus, while it purports to be a guide for the treatment of dangerous, varying, and violently acute diseases, carries with it, when used in the manner stated, sources of weakness, even to those whom it intends to benefit. It is otherwise in respect of chronic lesions, the results of acute tropical diseases, in which the constitution of the patient, and the symptoms are at once more subdued, brought more into a common level, rendered thus less variable, and for the management of which ample time is given to the inexperienced for consideration of means, for reflection on their effects. and for consultation with the elders of the profession.

In chronic disease, again, no sudden changes occur, as the result of youth, constitution, influence of season, or of epidemic conditions—all which exercise powerful, immediate, and varying

effects in acute disease.

We cannot defer till to-morrow the treatment of a case of fever, dysentery, hepatitis, or cholera; duty requires that we should act on the instant; and to do so with justice to the patient requires at least a knowledge of pathology and of the principles of medicine; we should not have to rely on a guide or case-book.

MANAGEMENT OF CONVALESCENCE.—I cannot conclude these observations without adverting to the importance of the management of convalescence from fevers, remittent and intermittent—not the least serious of the duties imposed on the naval and military surgeon. In all cases of recovery from fever, but especially in those wherein the complications have been severe, or where important organs have been affected in the course of the fever, or as a sequel to it, it is impossible to be too careful in the diet, and in attention to the activity of the secreting

Calcutta will declare that you have killed the Chief Justice." I persisted, nevertheless; and the removal to the river, which was about 500 yards distant, was accomplished without accident. But, in the act of being removed from the boat into a 74-gun ship at Khidgeree, notwithstanding my warnings to the contrary, the head and shoulders were raised to near the sitting posture, when the patient fell back in a fainting condition which for some minutes looked like death. But from this sunken state he recovered rapidly in the fresh breeze; and in ten days, he reached Madras, whence he was reported to be eating mutton chops. I have always been of opinion that but for the timely removal from malarious influences this distinguished gentleman must have died.

Next to the benefits to be derived from early and effective treatment, come those which secure and maintain convalescence; and this truth will be found to apply whether we refer to organized bodies of men, or to individuals. It is to the mismanagement of convalescence principally that we must refer the numerous and fatal relapses in the fevers and dysenteries of our seamen and soldiers, not to speak of ill health contracted through the sequelæ to those diseases, in the form of visceral affections. In India I have continually seen that almost all the relapses in fever, so often fatal to the soldier, were caused by mismanagement of the stage of convalescence, and by his too early discharge from hospital.

The mean age of a British regiment may be taken to be twentysix years—a stage of life when the rallying powers of the constitution may be asumed to be at their acme, and when, therefore, the tendency of nature towards convalescence is at its height. But in the individual cases of older men, and with such of the young men as have been exhausted by the violence of disease and the activity of depletory measures of cure, wine, porter, and bitter tonics will be requisite, together with more time, and a longer exemption from duty, for the purpose of maturing health. This latter consideration is of great importance.

It is true that the authority of the naval and military surgeon does not extend beyond the hospital; but better days than those of Robert Jackson are coming, both for the soldier and the surgeon. Humanity is now at length pretty sure to have a powerful press on its side; and no governor or commander, by sea or land, can any longer venture to do, or leave undone, the things which I

have seen done, and left undone.

When the disease is endemic, whether it be fever, dysentery, or hepatitis, the first means towards maturing convalescence, and obtaining the complete restoration of health is—a removal from the local and endemic influences.

This measure, coupled with great attention to diet, clothing,

exercise, and bathing, will fill the ranks with healthy soldiers; while leaving them in crowded or otherwise unhealthy hospitals is too often but condemning them to wait for death. In all our intertropical Possessions we should establish Convalescent Depôts. conveniently situated; and where practicable they should be in some near mountain range, or other elevated ground, or in well selected positions on the sea, which latter are, I think, always preferable. I know that in the East Indies one or other of such positions may generally be found by the intelligent medical topographer; and, for the rest, everything will depend on the authorities, civil and military. Nor are the advantages of such positions to be confined to convalescence; for where the locality is pestilential, or where barracks, camps, or hospitals are badly situated, the sick should at once be removed to the elevated or other selected site. Such early removal will be found to modify the progress of disease, aiding thus the medical treatment, and saving the patient from those congestions which are so liable to occur in the progress of fevers, especially in malarious situations.

GESTATION IN THE OPEN AIR.—Gestation in the open air presents itself at once as one of the most refreshing restorative means for maturing convalescence, and it is seldom difficult of application under willing authorities. It should never be neglected in the late stages of fevers; for it gives tone and vigour to the flagging functions, and conduces to refreshing sleep where medicine has failed of securing rest. Of gestation in the open air, Robert Jackson observes that it is neither useful nor safe in the early stages of fevers, while plethora exists; but that in the late stages of concentrated endemic fevers, it constitutes the "last anchor of hope" to the sinking invalid. He complains justly that this powerful remedy has not been applied generally and systematically; for he had frequent opportunities of directing gestation in the open air, and of observing its beneficial results, in the first American war, in the West Indies, and in Holland. He says that "moving the body rapidly in an open cart or carriage through woods, or on the green turf; or in defect of woods, shaded, if in the day time, with boughs of trees, exposed, if in the night time, to all the freshness of the air, and all the dews of heaven, has appeared to do what no other means were capable of doing." Jackson considers that airings for an hour or two in carriages, where the subject is defended from the weather, or but partially touched by the salutary influence of the air, constitutes but a feeble substitute for journies of six or eight hours in open conveyances, even when the patients are exposed to wind and rain, to heat and cold, and other apparent disadvantages.

Where, as in the retreat through Holland, in 1795, this measure

was adopted from necessity, he says that "it often succeeds; for the operation is continued till the effect is confirmed." Under all the disadvantages of this retreat, Jackson declares that the benefits to the sick where demonstrably felt. "Where due attention was bestowed in disposing the sick in wagons, the travelling was agreeable during its continuance even in cold and rainy weather; the good effect was strongly manifested at the end of the journey; even the most enfeebled acquired an evident accession of strength. Where the diseased action has actually ceased, but where the commencement of healthy movement is slow, and the healthy action imperfect, more benefit is visibly derived, and more strength visibly gained, by travelling for six hours in an open carriage, exposed to all the chances of weather, than by the best treatment that can be devised in a crowded hospital for a space of six days." The beneficial effects, he adds, are confirmed and rendered permanent by a continued application of the means.

This great authority expresses his regret that gestation in the open air—a means that has the smallest chance of any other medical appliance of doing harm—is not numbered among the regular means of physicians in the cure of fevers; and this regret must be shared by all who have seen and served in the sickly camp. The old 71st Regiment, or Fraser Highlanders, with which Jackson served during the first American war, was forced to retire before a large body of the enemy, carrying along with them a heavy sick-list of men suffering from remittent fever. "The sick, during the march, had little opportunity of taking medicine; yet no one had died; some had got entirely well; and in others, indeed in all, where the disease had not yet ceased, the form was changed to that of dis-

Those few survivors, who, like the writer of this work, marched with the miserable remnant of the European army originally sent to Rangoon, during the first Burmese war, will never forget the drooping attenuated form and haggard eye of the British soldier on the day we broke ground from the last mentioned town, and the amended condition of the same man on entering Prome two months afterwards. Yet here the soldiers were moved under every conceivable disadvantage, excepting in the matter of diet; for, as we advanced into the interior of the country, we procured some fresh meat, and occasionally some fruits and vegetables.\*

It may now prove both useful and interesting to take a glance

<sup>\*</sup> There perished in the first year of the first Burmese war, at Rangoon alone, of the British portion of the force, 48½ per cent., of whom 3½ per cent were killed in action, or died of their wounds, being within one-half per cent. of the annual casualties from wounds in the Peninsular war, during 41 months of the most active service. The total European loss within the Burmese Territory, during the two years of the war, was 720 per thousand of the strength—being the greatest mortality of which as yet there is any record.

at the prominent parts of the treatment of tropical fevers by the following authors, all British but the first-named, in the order of their respective dates:—

1629.—Bontius:—Bleeding, general and topical, and repeated as occasion required—purgatives—opiates—extract of saffron.

1751. Gleghorn.—Bleeding, repeated according to occasion—cathartics—bark.

1757. Dr. Bogue, Bengal.—Bleeding—emetics—purgatives—mercury—bark—camphor in the cold stage.

1757. Dr. Huxham.—Bleeding—purgatives—diluents.

1760. Dr. Huck.—Blood-letting, repeated according to occasion—ipecacuanha and tartar emetic, so as to vomit and purge—bark during the remission.

1762. Dr. James Lind.—"Putrid and remitting marsh fever"—cautious blood-letting—vomits of tepid water, occasionally of emetic tartar and ipecacuanha—saline purgatives—cooling acidulated drinks—bark, speedily and freely administered.

1768. Pringle.—Bleeding, repeated according to occasion—active purgatives—antimonials—bark, as an occasional means.

1790. Dr. Balfour.—Vomits—calomel and purgatives frequently

repeated-bark and opium.

1791-98, 1820. Robert Jackson.—Blood-letting, copious and repeated, according to the severity of symptoms—the operation always performed in the recumbent posture, while the head and shoulders are dashed with cold water, and heat applied to the feet; purgatives, followed by moderate doses of calomel, antimonial powder and cathartic extract, so as to keep up "effective evacuation,"—bark—blisters, largely and freely applied when local congestions require them—gestation in the open air—change of air. In fevers with congestion of the cerebral and abdominal viscera, calomel as a remedy "of the first importance," one which "repairs mischiefs which no other means with which we are acquainted are capable of touching."

1795. Dr. Chisholm.—Calomel and opium to salivation.

1796. Dr. J. Hunter.—A strong saline purgative, followed by James's powder till remission took place—then bark freely administered throughout the disease, and during each remission—blisters—cordials—opiates—change of air.

1797. Dr. John Clark.—Mercurial purgatives—bark—anodynes. 1799. Blane.—Bleeding—vomits and purges—sudorifics—bark

-anodynes.

1799. Dr. Lempriere.—Blood-letting—active purging by mercurials and cathartics perseveringly used—diaphoretics—blisters—pediluvia—saline draughts along with the bark, to allay nervous symptoms—ether, camphor, and wine; in doubtful and dangerous cases, mercury to affect the system.

1804. James McGrigor.—Emetics occasionally—blood-letting—cold affusion—calomel with purgatives—mercury to affect the system—blisters—nitric acid—opiates—bark.

1807. Mr. Curtis.—Evacuants and diluents, in the first stage-

calomel, ipecacuanha, and purgatives in protracted cases.

1808. William Fergusson.—Early and copious blood-letting—calomel to affect the system—sudorifics—purgatives.

1811. Dr. Bancroft,-Bleeding-cold affusion-calomel-pur-

gatives-bark.

1813. James Johnson.—Blood-letting, general and local—mercurial purgatives—diaphoretics—calomel, according to severity of symptoms, to affect the system—bark—change of air.

1816. William Burnet.—Bleeding both general and local—

purgatives.

1818. Ballingall.—Blood-letting, general and local—purgatives—cold affusion—moderate use of calomel—occasional emetics—bark.

1819. Dr. Dickinson.-Vomits-blood-letting-active purging

-cold ablution-diluents.

1827. Mr. Geddes.—Blood-letting, mercurial and other purgatives—diaphoretics—diluents—opiates before the paroxysm—quinine.

1828. Annesley.—Blood-letting, general and local—emetics—full doses of calomel—purgatives—diaphoretics—cold affusion

and cold lotions to the head—bark during the remissions.

1832. Mr. Twining.—Blood-letting, local and general, repeated as occasion required—calomel, followed by purgatives—calomel so as to effect the system in severe cases—quinine in the remission.

1833. Dr. Joseph Brown, Cyclopædia of Practical Medicine.—Bleeding, general and local, aided by warm baths—mercurial purgatives—cold affusion—cold to the head—cold acidulated drinks—in the advanced stages, opium—change of air.

1835. Copland, Dictionary of Practical Medicine.—An emetic—blood-letting, general and local—"full doses of calomel followed by purgatives,"—evaporating lotions to the head—cooling diapho-

retics—quinine during the remissions.

From the coup d'œil here presented we learn that to the cure of fever there is neither a short path nor a royal road, and that a disease so varying in its nature, so general and complicated in its influence on the system, is not to be justly treated by one remedy. Bark and calomel, each a remedy of great power, will nevertheless not succeed in the cure of fever, if used exclusively, and so it is with the most powerful of all means, blood-letting. We must, therefore, give to each remedy its proper place in the treatment; we shall thus be as nearly right as is possible. No one can be found now-a-days to follow Dr. Clark or Dr. Chisholm in his

exclusiveness. Men of science cannot be induced to believe that because one practice is wrong its very opposite must be right, thus meeting one extreme by another extreme. The truth is that exclusive doctrines and practices never take any general or permanent hold on the mind of the medical profession; and even the great names of Brown and Broussais have secured them but few followers in Europe.

GENERAL REMARKS IN CONCLUSION.—If tropical fever and dysentery were always simple morbid actions, or mere inflammatory states of the system, or of particular organs, no doubt bleeding and purging, as recommended by some writers, might often prove of themselves sufficient to the cure; but unfortunately, in both instances, we very seldom find this unmixed condition of disease in actual observation in Bengal, where, besides the morbid condition of the entire system, including a violent disturbance of the nervous and vascular systems in particular, we have in our fevers continually to combat dangerous abdominal complications, congestive and inflammatory, with the addition, in the fevers of the hot season, of the cerebro-spinal, all demanding a more or less complex and careful treatment, a speedy unlocking of all the secretions and excretions of the body, which the most ample experience proves that bleeding and purging alone will not effect. Tropical fever, dysentery, and hepatitis are produced by the action of specific causes, and such diseases are nowhere cured by the exclusive use of blood-letting and purging, as in the instance of common or of mere inflammations. Yet here, as in dysentery and hepatitis, bleeding is the standard remedy. subject to age, constitution, and length of residence in India. Bleeding, whether general or local, should be practised to the extent of reducing the force and frequency of the pulse, and of relieving the loaded and oppressed abdominal viscera. It precedes all other means of cure, both in order of time and in importance. The fevers of Bengal have for a length of time been treated according to indications grounded on explanation of the symptoms, on the physiology, and not on any exclusive plan. Accuracy of diagnosis and simplification of treatment, the true aims of the practical physician, have very generally been held in view. Blood-letting has been had recourse to, to moderate the force and frequency of arterial action, and to relieve congestions; free purgation, by means of mercurials conjoined with cathartics, to remove accumulations or vitiated secretions, and to aid in correcting the latter; calomel with sudorifies to act on all the secretory and excretory functions; bark or quinine during the remissions, so as to arrest the coming paroxysm; cold affusion in the ardent continued fevers of the hot season—these have long been in use with us.

Subject only to the limitations already stated, bleeding-early bleeding-whether general or local, and always practised at the very onset of the stage of reaction, is very generally necessary in the severer forms of Bengal remittent fever; then come full doses of calomel and sudorifies, short of producing salivation, with saline purgatives, and quinine in the intervals. If under such treatment the disease does not speedily yield, but on the contrary, if the secretions become of a watery nature, or if they be suppressed, and the paroxysms recur at shorter intervals, or with increasing severity, leaving but imperfect remissions, then there is imminent danger, and inflammation or congestion, more or less acute, in some important abdominal or cerebral organ may be more than suspected. For the cure of this state, in addition to topical bleeding and cold applications to the head, when it proves the seat of disease, calomel in small and repeated doses, with antimonials, must be exhibited so as mildly to affect the system; its judicious use, and that of quinine, constitute the only known means of saving the patient by anticipating the destruction of some organ essential to life. I have seldom had occasion to urge mercury to the extent of salivation, during the whole period of my service in India. Where there is much irritability of the nervous system, or tendency to sinking of the vital powers, the union of camphor with the calomel, as recommended by the Indian medical officers of old, will be found useful. In such conditions it will be found more safe than opium, especially in the hands of the inexperienced, for it does not favour congestion, or lock up the secretions as opium is too apt to do. But opium is necessary in some cases, and given with care and judgment it will then save life. In treating seamen and soldiers suffering from tropical fevers and other acute diseases, we should very carefully inquire into the previous habits of the patient, for when these have been dissipated, including the free use of ardent spirits and tobacco, we must be on our guard against the complication of delirium tremens more or less masked by the fever. This is a delicate point to determine for the young surgeon; but, looking to the history of the man's life, and to the predominance of nervous over vascular disturbance, he will soon perceive that an opiate, and not the lancet, is the remedy; the latter would indeed be fatal. Here the nervous system has been rendered feeble and irritable, while the blood has been depraved, by habitual excess.

The tendency to sinking of the vital powers is another most important consideration, never to be overlooked in treating the remittent fever of Bengal. This impending event is the reason why all our active antiphlogistic measures of cure, as blood-letting, purgatives, and sudorifics, must be so cautiously regulated as to

the time of using them; for what was a saving means at the commencement of the paroxysm, is as surely destructive at the end of it. On the subject of blood-letting, Robert Jackson's works abound in valuable injunctions, such as ought to be present to the recollection of all who undertake the onerous duty of treating tropical fevers. After the exposition of admirable rules as to the "just point of time" for the application of this most powerful means, he concludes by declaring that "the same remedy, after the delay of a few hours, not only ceases to be useful, but the application of it even sometimes becomes unsafe:" further on he adds, with equal truth, that "it requires much discernment in many cases to discover the cause, a very correct judgment to measure the means, and even no small degree of knowledge to be able to ascertain that the end is attained."

That our fevers in Bengal alter their types, nay that instances both endemic and epidemic occur in which blood-letting is not only unnecessary but injurious, the pages of this work sufficiently attest, and death even from its ill-timed use has more than once come under the author's observation; yet the fact remains untouched that blood-letting, practised with the care and thoughtfulness which should mark every step we take in this disease, at the proper time, and apportioned to circumstances of constitution, age, sex, season, and length of residence in India, forms the principal means of cure in our endemic fevers, and of preventing those organic lesions which otherwise so frequently follow them. Leeches are a ready and powerful means in the treatment of tropical fevers of every type; but I would caution the inexperienced against this mode of abstracting blood as a general rule, or as a substitute for venesection and I would likewise place him on his guard against the ill effects of prolonged oozing from leech-bites, especially in the instances of persons at the two extremes of life-the very young and the aged. I have seen the most serious results from carelessness in this respect.

In regard to the next most powerful remedy in paroxysmal fevers, namely, the peruvian bark;—where the remissions are becoming sufficiently well marked, quinine should be given in full doses without waiting for a perfectly favourable condition of the system:—in bad fevers we should seize the very first dawn of remission to exhibit quinine. Some practitioners recommend that before this drug is used we obtain a clean tongue, natural secretions, and the absence of all heat of skin, and of local complication. I believe this to be a very dangerous practice: if we are to wait for everything favourable we shall often have to wait too long, and till it has become too late. It is in acute diseases as in critical circumstances political and commercial—everything is for him who can wait; but, unhappily, in the formidable remittents of the tropics we can wait for nothing, for there is not

a moment to lose. I have always given quinine in the more favourable cases now under consideration, in disregard of certain abdominal complications (those of the cerebral cavity should in general exclude its use), believing that if I arrested the paroxysms, the progress towards curing the disease as a whole, greatly outweighed any harm which the quinine could possibly do to the local affection, the treatment of which, by local depletion, by mercurials, or by counter-irritants, is not interfered with by the means in question. Again, all tenderness on pressure, or local pain, does not necessarily constitute inflammation; and even where such is actually present, its character is often so modified by malarious influences, and anamic states of the general habit, as to constitute a state very different from idiopathic and uncomplicated inflammations, such as may demand only the kind of treatment here immediately referred to. In respect to quinine, I have often combined it with sudorifies, and I think with advantage: where there existed depression of the vital powers I have added small doses of laudanum. It is also usual with us in Bengal to combine this powerful antiperiodic with common purging mixture. a mode whereby it applies itself to the entire and extensive surface of the mucous digestive organs, so as materially to add to its influences.

It is well known in the history of the treatment of the fevers of India—a history so instructive that it ought to be known to all of our profession who may serve there—that at different times bark, calomel, and blood-letting have, each in its turn, been in almost exclusive favour with some individual practitioners; but such partial plans of treating so formidable and complicated a disease has never, for any length of time, held influence on the more thinking portion of the profession. I would guard and caution the younger medical officer alike against the adoption of exclusive measures of cure, and against the over-treatment of fever, both being equally mischievous. In truth, it is only by a rational and well-adjusted plan, which shall call into operation all the aids suggested by science and by experience, that such a disease as fever can anywhere be justly treated.

Finally; it cannot be too much or too often impressed on the Indian surgeon that it is on his careful attention to the phenomena and treatment of fever that nine-tenths of his usefulness depends. I have here presented the reader with a summary of the nature and treatment of the Bengal remittent fever: it will be found to correspond in most of its details with the history of endemic fevers generally, with the bilious remittents of the Indies, whether East or West:—they all have a pathological community, they are all fevers of locality, and do not by any means differ so much as medical writers of partial views and partial experiences would have us believe; their supposed differences, or nasological divi-

sions, are more frequently the work of man than of nature:—they may and do differ in degree of intensity; but their essential phenomena, and the organs affected in their progress so as to endanger or ultimately to destroy life, are the same, and so like-

wise are the essential parts of their treatment.

I have now concluded this the most important article in this work; and I trust that, provided I have treated the subject worthily, the reader will not consider that I have devoted too much space to the discussion of it; for he who can successfully combat the remittent fevers of tropical climates need not despair of acting with a like success in the treatment of the dysenteries and hepatic diseases of those regions. The principles here inculcated will, if I mistake not, guide the medical officer to such favourable issues, under every variety of type and complication of remittent fever as may reasonably be attainable, in the present state of our knowledge.

For the following Table, exhibiting the comparative frequency and the intensity, as shown by the relative mortality of remittent fever, throughout the wide extent of climate occupied by British soldiers, I am indebted to my friend Colonel Tulloch:—

STATIONS.	Period of Observation.	Aggregate Strength.	Remittent Fever.			
			Attacked.	Died.	Proportion of deaths to admissions	
Windward and Leeward }	20 years	86661	17799	1966	1 in 9	
Jamaica	20 "	51567	38393	5114	1 in 8	
Gibraltar	19 "	60269 {	YF.* 314 1522	28 423	1 in 11 1 in 3#	
Malta Ionian Islands	20 "	40826 70293	384 6934	16 623	1 in 24 1 in 11	
Bermudas	20 "	11721 {	YF.* 19 277	6 101	1 in 3 1 in 24	
Nova Scotia and New !		10110	15		1 in 15	
Brunswick	20 "	46442 64280	294	18	1 in 16	
Western Africa	20	1843	1601	739	1 in 2	
Cape of Good Hope	19 "	22714	15	1	1 in 15	
St. Helena	9 "	8973	25	1	1 in 25	
Mauritius	19 "	30515	6	1	1 in 6	
Ceylon	20 ,,	42978	4643	868	1 in 51	
Tenasserim Provinces	10 ,,	6818	594	22	1 in 27	
Madras	5 ,,	31627	1139	54	1 in 21	
Bengal	5 ,,	38136	1311	89	1 in 144	
Bombay	5 ,,	17612	2854	114	1 in 25	

<sup>\*</sup> The prevalence of Epidemic Yellow Fever is here indicated.

### INTERMITTENT FEVER

THE proportionate amount of sickness from intermittent fever, in any given communities, will depend on the relative amount of local sanitary improvement, or of the neglect of it, in the districts, stations, and cantonments occupied by them throughout our intertropical possessions. The geographical distribution of the class of fevers here treated of will be found to be very much the same with that of remittent fevers; where the one form of fever prevails, there we generally find the other likewise; and whether an European is to be affected by a remittent or by an intermittent fever will depend on length of residence in a tropical climate, on age, constitution, and previous habits of life, on season, and the intensity of the exciting causes.

The usual history given by Europeans who have resided in malarious districts or stations is, that they had first been affected by remittent (jungle) fever, and that then followed the intermit-

tent form.

Dr. Bryson, in his Statistical Reports on the health of the navy serving in the East Indies, says of the fevers of China:—"A large proportion of or nearly all the cases of intermittent were the sequelæ of fevers which had first appeared in the continued or remitting forms; still there were a few which seem to have been intermittent from the commencement, although it is probable they were connected with preceding attacks which were not observed,

or had been forgotten."

In Calcutta, through the hitherto gradual improvements in its ill-chosen site, intermittent fever has become, for many years past, a mild and infrequent disease comparatively, especially among the better classes of society, and amongst Europeans of better habits of life. This simple fact strongly illustrates the beneficial influences resulting from local sanitary improvements, while it constitutes a powerful inducement to the government and the public of the city in question, to proceed in the same course of amelioration which has already secured to its inhabitants so great an exemption from disease.

At home, and in our own capital, we find that the sweating sickness, scurvy, dysentery, malignant ague, and other formidable diseases, have yielded to the slow progress of sanitary improvement in modern London; so, in our Eastern capital, we have no longer "the obstinate putrid intermitting fevers," described by the older writers, with their cold stage of "twelve hours," and their long list of sequelæ in the form of tumid livers and spleens,

diarrhœas, dropsies, &c. &c., a condition of public health which we now regard, even in the capital of British India, with horror.

Dr. Bogue, who practised in Calcutta in 1757, speaks of this form of fever "as the most fatal" of that time. It began with the rainy season, and continued with increasing violence during its continuance, and "for some time after," the paroxysm recurring daily, so that "the patients had not above four or five hours respite from it." Thus he says, "we had sick at the same time, in this place, one half of the men of the squadron under the command of Admirals Watson and Pocock."

The fevers here described would seem to have possessed the malignant character of the *febres intermittentes algidæ* of Torti, in which the power of generating heat was so impaired that the patient died in the cold stage at the end of two or three accessions.

Immediately, our modern intermittents are in general not dangerous to life, but, consecutively, the dangers arising from visceral diseases, in after years, are considerable. Of this latter fact I have had presented to me in England very numerous instances, and one very remarkable will be found under the head of Chronic Diarrhœa. In certain districts of the East Indies, as in others of our colonies, the type of intermittent fever assumes a malignant form, and then the dangers of a rapid dissolution, or of visceral disease and a broken constitution, are very great.

I have said that ague is an "infrequent" disease in Calcutta, and strangers will read with surprise that I do not think I saw above a dozen cases in a year, on the average; and these occurred in persons who went into the neighbouring jungles on hog-hunting and other such excursions. But this disease is of frequent occurrence in the marshy and jungly portions of Bengal Proper,

as well as of all such countries throughout India.

I remember attending a mercantile gentleman who contracted his ague in the Sunderbunds, about the middle of October. On his return to Calcutta, about the 26th of the month, he experienced a good deal of malaise, and took medicine from a chemist with partial relief. Towards the end of the month the native servants who accompanied him fell ill of ague one after the other, and on the 20th of November their master was seized with a violent paroxysm of the same disease. Here we perceive the superior power of resistance to the morbific influence exhibited by the European over the natives, as shown by the difference in the period of latency or of the abeyance of the fever in each, being more than thirty days in the former, as compared to ten and fifteen days in the latter. This gentleman was temperate in his habits, and inured to the climate; and both master and servants entered the Sunderbunds in good health, and neither had been previously affected with ague.

CAUSES.—That the emanations from marshes produce agues is the universal belief of mankind in all ages and countries; and the accordance of the medical profession in this opinion has

always been quite as general.

Many authors believe with Cullen that, in the climate of England, this fever can alone be produced by exposure to marsh exhalations; and Sir John Forbes has shown in a very able memoir on the medical topography of Land's End in Cornwall, "that neither impure air simply, nor wet, nor the alternations of cold and heat, nor all these combined, can give rise to fevers of this type." This I believe to be quite true in respect to a first seizure; but after that, and when the disposition to relapse is once established in the system, such a combination of influences will certainly in tropical climates prove an efficient cause; and even Cullen admits "the concurrence of other existing fevers," when the malaria is not "strong enough to produce disease." In support of this latter view I quote the following important passage

from Sir James McGrigor:-

"After the effluvia from marshes or the exhalations raised by a powerful sun acting on a humid or luxuriant soil, we found that in those who were convalescent or lately recovered from agues, the causes next in power to reproduce the disease were exposure to a shower of rain or wetting the feet, exposure to the direct solar rays, or to cold, with intemperance and irregularity, or great fatigue. Many other causes would excite the disease in the predisposed, but these never failed to do it. In marching troops in a country where the disease is endemic, particularly if they have been lately discharged from hospitals, the above causes should by all means be avoided, since the whole of our experience in the Peninsula showed that relapsed cases seldom or never get completely well in the country in which they were contracted, under all the circumstances of a soldier's life. In making calculations of efficient force this description of men could not be depended on for operations long continued in the field." Pringle—the father of British military medicine-writes to much the same effect:- "After the frosts of November the intermittents never appeared unless upon catching cold, and even then such only as had been ill of them in autumn were seized in that manner."

The important observation is continually repeated also by this great author, as to the excessive liability to attack of those who occupied the lower floors of houses and barracks, and this

happened all over Flanders.

SYMPTOMS AND PROGRESS.—The symptoms of intermittent fevers are well known to consist, like those of remittent fevers, in paroxysms of three stages—a cold, a hot, and a sweating

and the disease is protracted, the patient incurs much risk of disease of the abdominal viscera.

PATHOLOGY.—All the organs and functions which are disturbed or oppressed during the paroxysm of remittent, are likewise more or less affected in that of the intermittent fever. During the cold stage of the latter the brain is occasionally oppressed by congestion, while all the abdominal organs are similarly affected; and in the succeeding hot stage the same organs and functions are more or less seriously disturbed in the tumult of vascular reaction. There is flushing of the face, headache, and sometimes delirium, the patient being at the same time distressed by general uneasiness, restlessness, and epigastric anguish. It is no wonder that morbid actions and alternations such as these, repeated during weeks, months, or even years, should result in organic diseases of a formidable character.

Agues which have become habitual by repetition will also impart a character of periodicity to other ailments, such as neuralgic affections. A few years ago I became subject to facial neuralgia, which came on regularly at eleven A. M.—the very hour at which ague used to commence with me in India more than twenty years previously. Further, when the intermittent fever has been severe, or long-continued, the disposition to recurrence of the same disease seems to last for life. After a residence of ten years in Europe I happened to pass three nights at the best hotel in Strasburg, at a time when ague prevailed in the garrison amongst the French soldiers who had served in Algeria; and two days after quitting that town I was seized at the accustomed hour of eleven A.M. with ague, and I was the only person of the party who was so affected.

We have in the East not only every variety of type of intermittent fever, but every degree of intensity also—certain districts and provinces being noted for the malignity and fatality of their agues; and even in Bengal Proper we have often to combat severe and obstinate visceral diseases, the sequelæ to these intermittents. Under neglect of early and effective treatment such cases terminate in permanently broken health, through disorganisations of the liver, spleen, or mesentery, or of all three. In the malignant intermittent fevers of Aracan, and Gondwana, and of provinces of a like unhealthiness, we perceive such remote ravages of disease as to prove at once conclusive both of the original intensity and danger of the disease, and of the ultimate serious consequences, even after the lapse of years. One such case I will adduce by way of illustration:—

Captain — contracted the intermittent fever of Aracan in 1837, which was speedily followed by great enlargement of both liver and spleen. At my recommendation he was sent to

England for the recovery of health, whence he returned to India in 1839. His health continuing very indifferent under various changes of climate, he was constrained to resign the service in 1847. But even in England he was always an ailing person; and in October, 1852, I was consulted on the state of his health, which was then desperate. He died two months afterwards in a

state of universal dropsy.

For the following post mortem record I am indebted to Mr. Pollock of St. George's Hospital :- " The peritoneum was covered throughout its free surface with an opaque white false membrane. This membrane was thin on the surface of the abdominal wall and small intestines, but was very thick elsewhere, as on the surface of the stomach, transverse colon and liver, from which viscera it could be peeled off. From the peculiar manner in which it was spread over and attached to these organs, when the abdomen was opened, the small intestines were the only viscera that could be observed. The large intestine, stomach, liver, and spleen were hid from view by the false membrane glueing them together; and the only indication of their situation was an irregular nodulated mass covered with this membrane, and situated at the upper part of the abdominal cavity. The membrane on the right side was continuous from the abdominal wall over the ascending colon to the spine, and so bound down the ascending colon that, until the membrane was removed, the situation of the bowel was not ascertained. From the surface of the bowel the membrane passed to and attached itself to the mesentery of the small intestine over the vertebræ, in some parts thicker than others, so that it might almost be said to consist of bands, and, from their attachments, they must have much compressed the ascending colon. On removing the membrane from the irregular mass, the stomach, transverse colon, omentum, and spleen, with the liver, were brought into sight; but all these viscera had been compressed as it were into the mass by the false membrane spread over themthe membrane being strongest in this situation, and equal to three or four sheets of paper in thickness. The membrane here was readily removed. The liver was enlarged, but chiefly on its left side; and between the longitudinal fissure and the extremity of the left lobe and its anterior margin, there was a whitish yellow mass of hardened deposit which dipped some two inches into the substance of the liver, and it was some three inches broad. When cut into, it was firm and consistent, and appeared to consist of fibrous tissue with portions of fatty matter, somewhat resembling the character of scirrhus, but less hard. The spleen was healthy, but larger than usual. The kidneys were healthy. The other viscera were not examined."

The reader will now be able to appreciate the original violence

of the intermittent fever in this instance, and the necessity therewas for the most active antiphlogistic means, including bloodletting, to overcome inflammatory complications. Such means were not applied;—or, if applied, they were not of sufficient power; -and hence, in all probability, the unfortunate issue.

In all the severer intermittent fevers of India, such as that by which the European and Native force under General Morrison was destroyed in Aracan, during the first Burmese war, we find the sequelæ of diarrhœas and dysenteries to be very frequent also; \* and it is justly observed by Dr. Schulhoff, in his Notes on the Diseases of Turkey, that where there is ague dysentery is not far off. Congestive and inflammatory affections of the spleen and liver, diarrheas and dysenteries, the results of intermittent fevers, are most frequent and most severe during the cold season in Bengal.

The influence, real or supposed, of splenic disease in keeping up the morbid train of actions of the original fever, and in producing relapses, is a subject well deserving the careful attention

of the tropical practitioner.

M. Piorry, on his attentive examination of the state of the spleen in more than five hundred cases of ague, has come to the conclusion that that organ is invariably enlarged during the progress of the fever; and he has ascertained with equal accuracy the fact that, by the use of quinine the spleen is diminished in size. M. Piorry states :-

. The devotion of the medical officers attached to this portion of our invading army was most notable. Mr. Grierson, of the Bengal Medical department, says :- " Of the first five deaths that occurred among the European officers, four were in the medical branch of the service. This blank was severely felt; but twice the above number were soon on the sick list; and at length only eight medical officers remained for the whole duties of the division, the sick of which at that time amounted to near 5,500 men, besides the sick of the various public establishments, and the camp-followers." All who were not dead were in hospital. From this simple statement, the hopeless nature of the duties devolving on the few surviving surgeons of the army of Aracan may be inferred; and it will be matter of no surprise that in a few months, three-fourths of the entire force, European and Native, perished on the spot; while the suffering survivors were ruined

Mr. Lewis Grant, of the Bengal Medical Department (brother to the gallant Sir Colquhoun Grant), was the principal medical officer. It is related of him that "he was attacked by the fever, and recovered. When told that to save his life he ought to go, he replied, 'No, Sir, this is our post, and I will not set the example of leaving it.' He soon afterwards had a relapse, which proved fatal." The Native officers of Gardiner's Horse tried to cause the authorities to interfere with their surgeon to take care of himself, and be less in hospital. "The death of a dozen of us does not matter," they said; "but if he

dies, the whole corps will perish."

Mr. T. C. Robertson, the humane and most able Commissioner on the spot, says, in his "Political Incidents of the First Burmese War:"-" The devotion of all members of the medical staff to their duties was most exemplary. Not one of them, however ill, withdrew from the Province, where many of the number died. I have myself seen some of them, in the intervals of their own agues, wrapped up in cloaks and busied in visiting the sufferers under their charge." The Commissioner, who subsequently rose to be Governor of Agra, might well designate the conduct of Mr. Grant as "a fine example."

organs referred to, as well as to the state whether congestive or inflammatory of the cerebral and abdominal organs generally. It is easy in the average of cases to cure a simple uncomplicated ague; but it is not so when the type of the fever is malignant, or adynamic, or where the complications are severe. In every case of ague there will be more or less of simple turgescence of both liver and spleen during the cold stage, especially if this last prove of long duration; but this turgescence disappears in mild cases with the sweating stage of the fever. It is different, however, under concentrated endemic influences, and where the ague is consequently pernicious; for here the spleen and liver, if they do not become immediately affected, are sure to be so eventually. Repeated over a succession of months or years, even simple turgescence may terminate in organic disease, in defect of early or proper treatment. In truth, the difficulties encountered in the treatment of agues consist in the presence of the original malarious influence, and in the existence of the complications above mentioned. Wanting these two, or, in other words, under change to a purer air, and in the absence of cerebral or abdominal complication, agues are very

Emetics would seem to be indicated in the cold stage of this fever, with the view to restore the balance of circulation and relieve oppression resulting from congestion of the viscera; but here, as in the remittents of Bengal, emetics are not, and never have been, in favour with the practitioners there. Warm drinks, ammonia, ether, camphor, and other diffusible stimuli, and warmth externally applied, are preferred by the great majority.

When the hot stage has come on, a full dose of calomel with James's powder should be given at once, and in three hours this should be followed by a brisk cathartic—diluent drinks being freely used meanwhile, along with some cooling diuretic: the tartarised antimony, with nitrate of potash, answer the double purpose of determining the fluids to the skin and the kidneys. On the following morning, the intermission being complete, the sulphate of quinine is to be administered, at intervals of three hours during the day, the patient being kept in bed, and supplied with farinaceous food only. The treatment here briefly indicated will speedily cure a mild and uncomplicated ague. In the simple cases here contemplated it may not be necessary to administer mercurials more than once or twice; but active purgatives are always beneficial in relieving the full and congested state of the abdomen generally during the continuance of intermittent fever.

But where the disease is severe, and accompanied by præcordial oppression, pain, or fulness of the spleen or liver, or of both, or where there is severe headache, or headache with giddiness, or an oppressive fulness of the chest, a general or local blood-letting,

or even both together, will be necessary, as a means of present cure, and as a preventive of future ills. Measures of depletion are as necessary here as in remittent fever, and repeated applications of leeches, with several doses of calomel and strong purges are required, in these severe cases, to prepare the system

for the antiperiodic power of bark, quinine, or arsenic.

"Depletions," says Copland, " are almost indispensable preliminaries to the quinine or bark, especially in the complicated and congestive forms: for, without them, it will either not be retained on the stomach; or, if retained, will convert congestions, or slight forms of inflammatory irritation, to active inflammation, or to structural change." He adds that it is chiefly to a neglect of this practice that complications and unfavourable consequences so often follow the use of bark, quinine, or of arsenic; for these often interrupt secretion, and over-excite and inflame loaded, obstructed, or congested organs. As in the milder instances, mercurials with sudorifies, occasional diuretics, followed by powerful catharties, and quina, are here necessary; and on their proper application the efficiency and the speed of the cure will depend. Blood should be drawn here, as in the remittent fevers, at the very onset of the hot stage, or that of reaction; and it should be regulated by the constitution, age, and habit of the patient, as already explained. By such timely depletion congestions and inflammatory conditions will be removed, and much future disease and suffering will be prevented.

Where, on the other hand, the intermittent fever assumes a low adynamic form, or where the subject is in a state of anæmia, with or without enlargement of the spleen, mercurials should be most carefully avoided in the treatment, and reliance placed on change of air, quina, chalybeates, and improved diet. Under the condition of the general habit now referred to, mercury must not be administered for the cure even of hepatic enlargement:—the nitro-muriatic acid must be used instead, both internally and externally. Mercury would here be at once ineffective and injurious—indeed dangerous—while the mineral acids are most powerful to good.

In the low form of ague, in the anæmic habit, and where there is chronic enlargement of the spleen, with the splenic cachexia, blood-letting, general or local, should also be avoided; but even in weakly persons who are not anæmic, local blood-letting by means of leeches will be found extremely beneficial, under the cautions as to subsequent oozing already pointed out. In removing epigastric oppression or tenderness, or similar states of either or both hypochondria, fomentations followed by leeches will always, in a tolerably sound state of the constitution, be safe and proper, as will occasional blisters. Warm-baths, used at bed-time, are likewise very beneficial.

The sulphate of quinine is exhibited variously by various practitioners-some, as Maillot, giving very large doses of twenty and thirty grains four hours before the expected paroxysm-while others begin to administer the quinine on the subsiding of the paroxysm, and during the sweating stage. But the most rational plan, so far as my experience extends, would appear to be that of exhibiting the quina every three or four hours during the apyrexy, and in such doses as the urgency of the symptoms may demand. In this manner a sufficient quantity of the antiperiodic will be best borne, while the risk of injury by overdosing will be avoidedseeing that during the intervals of exhibition we can watch and measure the effects. I have always exhibited the sulphate of quinine in this manner, I believe, with an average amount of success; and it can hardly be necessary to observe here that I have seen, over a long range of years, and in various countries of the East, very severe types of intermittents. To cure the patient by means of the quina is all that is wanted; and to exhibit more of that medicine than is needful to this object, or to protract the use of it beyond the occasion, is to injure him. I always administered the quinine in solution, and with the addition of a small quantity of dilute sulphuric acid.

In the instances of plethoric subjects, I often conjoined the quinine with antimonials; and where the patient was feeble and irritable, with symptoms of exhaustion, I added a few drops of tincture of opium to the antiperiodic. Several very able medical officers are in the habit of giving quinine in the intermittents of the East Indies, in doses of twenty-five to thirty grains "as soon as the patient began to perspire freely after the hot stage." They declare that this plan has proved in their hands more effective, and

more economical of the medicine.

Arsenic has for ages been known to and largely exhibited by the native doctors throughout India, in every form of fever, continued as well as paroxysmal; and they are in the habit of exhibiting the mineral in very large doses, and for a long continuance—frequently with permanent injury to the health, as mentioned at page 167. When arsenic is preferred in the treatment of ague, it should be given in the College formula of solution, not in bitter infusions, as recommended by some writers, but in some aromatic water, with or without a few drops of laudanum to each dose of the arsenical solution, as may be deemed necessary. From six to eight drops given in this manner, every three hours during the apyrexy, and continued only for a few days, prove very effective in some cases.

After the force of the fever has been broken by the use of quinine, or where relapses in lesser degrees of violence still harass the patient, the arsenical solution will be found a valuable aid and substitute; and this becomes an important consideration in the

field of active service, where quinine may be scarce.

The Swietenia Febrifuga, and the kernel of the Cæsalpinia Bonducella, and the Chiretta, are much used by the native practitioners of Bengal, as antiperiodics and tonics.

Where the intermittent fever has become chronic, or when as secondary results there is organic disease of the liver or spleen, or of both, change of climate becomes a measure of necessity; and it is one that should never be neglected in such cases. I trust I need not add to what has been said in the last article on this subject.

As, notwithstanding its scientific claims, fashion unhappily still holds some influence in medicine, and as bleeding in the cold stage of intermittent fever may again come under professional review, in the round of medical opinions and practices, I think it right to notice the subject. My own personal experience will not allow of my speaking in favourable terms of this practice; neither will a reference to authority prove more encouraging, as the following comparative quotations will show:—

#### DR. MACKINTOSH.

"The practice prevents debility in a direct manner by saving the vital fluid."

#### MR. TWINING.

"I may say that in all regular intermittents, with cold, hot, and sweating stages, and tolerable uniformity in the hours of accession, the practice of bleeding in the commencement of the cold stage, has proved always safe, and generally more successful than any other remedy."

#### DR. MACKINTOSH.

"I believe bleeding in the cold stage, conjoined with the occasional use of sulphate of quinine and laxatives, to be as certain a mode of treating intermittents as any other set of medicines can be said to be certain in the treatment of any other class of diseases."

#### DR. MACKINTOSH.

"The practice may be adopted in the first stage of all fevers."

#### DR. ELLIOTSON.

"It is not at all right to take away blood, and thus impair the power of the patient;" when, as he adds, he "never saw a case which he could not cure by the sulphate of quinine."

#### DR. WATSON.

"I object to bleeding because it appears to me quite unnecessary; because it is not such as the nature of the symptoms would suggest; because it tends to produce subsequent debility, which we should not needlessly inflict; and because the experience of other sober minded men, who have given the method a fair trial, does not bear out the statement made by Dr. Mackintosh in respect to its usefulness. If in this country bleeding be requisite at all, it is in the hot stage."

#### DR. STOKES.

"Bled in the cold stage, and found it useless and injurious; and, after all, was compelled to give quinine sooner or later."

#### DR. ELLIOTSON.

"The quinine which cures it best interferes with no other measures."

### THE CELEBRATED DR. GREGORY.

"Blood-letting, which at another period of the disease might have been proper, if employed in the *first stage* never fails to be attended with the most dangerous consequences; or it is, to use the words of Celsus, hominem jugulare."

#### DR. COPLAND.

"In a case where I directed bloodletting before reaction supervened, the loss of three or four ounces caused profound and prolonged syncope, yet within four hours, when reaction had come on, fifty ounces were taken before any effect was produced upon the pulse; and before the sun of the same day had gone down, forty more were abstracted at one time, in all ninety-four ounces within twelve hours."

# DR. DENMARK, PHYSICIAN TO THE FLEET.

"In these instances venesection was never had recourse to till the vascular system had fairly emerged from the depressed state incident to that stage of the fever, and reaction had clearly manifested itself by the returning glow of the skin, the filling of the previously shrunk and dejected features, and the firm though frequently oppressed beat of the pulse. Former experience not only taught me that an earlier abstraction of blood was never borne to an extent productive of ultimate benefit, but, on the contrary, seemed to be injurious by tending to protract the first stage of the paroxysm."

#### DR. WATSON.

"Drs. Townsend and Law, of Dublin, found it fail in the majority of cases. In Dr. Stokes's hands the most usual effect of blood-letting was to check the shivering; and, next to this, to mitigate the severity without abridging its duration. In most instances no modification was produced of the hot and of the sweating stages. In Dr. Kelly's experience the general effect was to shorten the cold stage, and to render the hot one milder; but in some cases it seemed to aggravate the symptoms. Mr. Gill found that, although the bloodletting might cut short the cold stage, it appeared to lengthen the period of febrile disturbance."

#### DR. STOKES.

"Having now described the effects of the practice on the paroxysm, and on the local symptoms, I must next mention some very untoward circumstances which appeared to follow the bleeding in the

#### MR. TWINING.

"The recent practice of bleeding in intermittents, as recommended by Dr. Mackintosh, not only accords with the acknowledged pathology of that class of fevers, but seems to bring our system of therapentics, as applied to them, within the limits of those established principles adapted to the treatment of other fevers, from which, formerly, intermittents were almost excluded."

#### MR. TWINING.

"The safety and efficacy of this practice have been so far established by Dr. Mackintosh that I have not hesitated to adopt it." . . . "The great advantage of V. S. in the cold stage of ague appears to me to depend on the prompt and decided relief it affords, guarding the patient against the ulterior results of repeated congestions of internal organs." . . . "It appears to me that V. S. in the cold stage of intermittent fevers deserves to be classed among the best remedies which we possess for the effectual cure of many of the most obstinate cases of these diseases in Europeans."

#### DR. MACKINTOSH.

"Bark has been long in use, and although I never denied that it had virtues, yet when given in substance in the large doses which are admitted to be necessary, I have so frequently seen it do mischief that the question has often suggested itself to me, whether it was not more injurious than beneficial? It seems to be injurious in many cases by overloading the stomach and bowels with indigestible ligneous fibre, and I have seen it cause serious intestinal irritation, as displayed by griping pains in the bowels, diarrhea, and painful tenesmus. On examining the stools in these cases, they seemed chiefly to consist of bark, with a considerable quantity of mucus occasionally tinged with a little blood.

"The preparation of bark, which is known by the name of the sulphate of quinine, is the greatest improvement in modern pharmacy, and the knowledge of its beneficial effects in simple intermittents affords sufficient proof of the virtues of the substance from which it is extracted; yet this remedy, all-powerful as it is, is useless in the cold stage, and must also fail in cases complicated with organic disease. . . . My youthful readers may rest assured that the same observations are generally applicable to the sulphate of quinine; yet they will most probably meet with many practitioners, who will assure them that they have never seen a case in which bark, exhibited in substance, or in any other form, has failed in their hands. When they hear such statements they may be satisfied that such practitioners never met with a severe case. or that there is some subterfuge. Some medical men, it is but charitable to suppose, are in the habit of deceiving themselves. In the instances which fell under my own observation, and to which I have already alluded, fever and violent cerebral symptoms succeeded, and, in two or three instances, local inflammations."

#### MR. TWINING.

"In the early stages of intermittent fever, or to speak with more precision, within two or three weeks of their commencement, in persons of robust habit, there is very often disorder of the functions and secretions of the digestive organs, and particularly of the stomach, co-existent with congestion of the brain, and attended in some cases with tolerably

cold stage-these were the occurrence of new local inflammatory symptoms, and the supervention of a low irritative fever. From the examination of the cases, I apprehend that an impression will be received against the indiscriminate or even frequent use of bleeding in the cold stage of ague. It may be remarked that in the great majority quinine had to be administered before the disease was eradicated; that many of them had an extremely slow and dangerous convalescence; that in several instances the disease, so far from being relieved, appeared exasperated by the practice; that local inflammatory affections occurred several times after the operation; and, lastly, that the bleeding appears to have a tendency to convert intermittent into continued fever. In one case, that of Caseley, death from pneumonia and softening of the brain occurred. In none of my cases did any bad effects from sinking of the powers of life follow the practice immediately; but I am informed that, in the practice of a highly respectable individual, there occurred two cases in which the patients did not recover from the collapse produced by bleeding in the cold stage. These facts should make us very cautious how we interfere with nature by means of the lancet in simple intermittent, where we have so certain, and as far as I have seen, so infallible a remedy as the sulphate of quinine. I do not deny that cases may often occur where venesection may be proper, such as intermittent complicated with severe internal inflammation; but shall only remark, as these cases have not come under my own immediate observation, that to offer my opinion on a purely practical point connected with them would be wholly useless. I may mention that I have been informed by my friends, Dr. Townsend and Law, of this city, that they have given the practice a trial, and have found it to fail in the majority of cases."

### DR. JAMES JOHNSON.

"What kind of inflammation must that be which explodes, as it were, the moment the clock strikes a particular hour, and this for days and weeks together? What kind of inflammation is that which, every second day, terminates in profuse perspiration from head to foot, and yet is renewed after an interval of forty-eight hours with the symptoms as before, and

distinct evidences of inflammatory condition either in the cerebral membranes or in the brain itself." \* so on? Do we see real and unequivocal inflammations pursue this course? Never. Are the causes of these intermittent phleymasiæ (if such an expression be not a solecism in medical language) of a periodical or intermittent nature? No. They do not accommodate themselves to any particular theory."

Thus it appears, at least in Europe, that the treatment of intermittent fevers by blood-letting in the cold stage, whilst it has the show of being prompt and energetic, proves in effect haphazard, systemless, operose, and tedious; and, from all that I have seen and heard in the East, the result there has not been more favourable. Modern science, indeed, is steadily though slowly achieving a signal victory, in the treatment of fevers especially—namely, that over exclusive systems. It is no longer with us as with the character in Moliere:—"A dead man is but a dead man, but rules are everything."

To conclude:—The rule of practice laid down by Pringle and Cleghorn has received little or no addition in more recent times. Where general blood-letting is had recourse to, in the treatment of intermittent fevers, whether simple or complicated, it should, as in the case of all other fevers, be performed at the very onset of the stage of reaction. Practised at this period of the febrile movement, it will diminish the force and frequency of arterial action, relieve venous congestion, usher in the sweating stage, and thereby pave the way for the administration of purgatives, sudorifies, diuretics, and quinine, on the just application of all which the prevention of recurrence must depend.

In the milder forms of ague, in feeble habits, or in the cases of persons who may have resided long in India, local depletion, by means of leeches, will answer every purpose.

<sup>\*</sup> The italics in the two last quotations are mine, and I have added them with a view to mark the vagueness and want of precision in both writers when they speak of inflammation, whether as affording an explanation of the physiology of this disease, as some writers would have it, or as applicable to mere occasional complication. It is to the former notion that Dr. James Johnson's observations apply.

# THE CONGESTIVE CONTINUED FEVER OF THE COLD SEASON IN BENGAL.

This fever, like all the diseases of the cold season in Bengal, is dangerous from its insidiousness. It is a frequent form of disease; and none are exempt from its seizure—not even the prudent of the better classes of Europeans. It attacks persons of all ages, and of both sexes; but men are, from their habits of life, far more exposed to it, and that in its graver forms. Its approach, unlike the fevers of the hot season and rains, is very gradual, being generally but little noticed either by the patient or his friends; or, if noticed,

it passes for catarrh, or dyspepsia.

At this stage, the functions of circulation and secretion are but little affected; yet, there is a harsh dryness of the skin, especially of that covering the abdomen, coupled with a sense of fulness and oppression at the epigastrium, which will not fail to attract the notice of the careful medical observer. In the course of a few days, provided these symptoms receive no professional attention and treatment, the circulation and respiration become hurried; headache is present, together with lassitude and anxiety, loss of appetite and rest, the tongue indicating disorder of the digestive functions. The loss of more time brings with it serious complications in the form of abdominal and cerebral congestions, as indicated by increased fulness and oppression of the præcordiatension of the whole abdomen-intense headache, with occasional delirium-and sometimes with yellowness of the whole surface of the body, jaundiced eye, and a surcharged state of the urine; in short, there are present all those morbid conditions which we might, a priori, anticipate as likely to arise from neglect, under the influences described at pages 49 and 50, and which cannot be too deeply impressed on the recollection of all who have to encounter the cold season in Bengal; for the same neglect may, according to circumstances, induce the fever here described, dysentery or the still more dangerous, deep-seated and insidious hepatic inflammation common to the cold season.

It is believed that in animals whose pulmonary system is comparatively less perfect, there is a greatly increased quantity of blood transmitted through the liver. In hot climates then, where respiration is less perfectly carried on than in such as are cold, owing to the greater rarefaction of the air in warm regions, according to Tiedmann and Gmelin, a vicarious decarbonization of the blood is

established by an increased flow of bile; and hence it is that the function of the liver, now weakened and rendered torpid, in proportion as its function has been unnaturally excited during the previous hot and rainy seasons, becomes disposed to congestion and inflammation of its parenchyma during the cold season, and thus are produced the dangerous states of disease noticed.

An irritable or inflamed state of the mucous digestive surface is a frequent complication; and these combined circumstances constitute the great dangers of our congestive fever of the cold season in Bengal, as well as those of the autumnal fevers of the more un-

healthy countries of Europe.

The form of fever is generally continued; but when protracted, it may and does frequently assume the remittent character, particularly where the patient has recently been exposed to malarious influences, as during the sporting excursions made during the cold season into the districts around Calcutta. In cases where we have to lament the neglect of early treatment, cerebral congestion with delirium prove occasionally long-continued and troublesome, especially when to these states are added retention of urine—all demanding much care and attention: but serious as these cerebral affections appear, I have generally seen that the patient recovered, provided they were not associated with abdominal complication of a grave nature. But when we have the attendants of yellow suffusion, parched and black tongue, jactitation, and a generally typhous condition, we must perceive that the condition of the sufferer is one of imminent peril.

The appearances on dissection are such as the nature of the disease and its complications might lead us to expect; various degrees of acute congestion, or of inflammation, with their results, being very generally found within the abdominal and cerebral cavities. In neglected cases we find hepatic abscess, and sometimes ulceration of the mucous digestive surface. This latter condition I found to be very prevalent amongst the labouring classes of natives, whom I had continually to treat at the Native Hospital of Calcutta on account of neglected fevers of from fifteen to twenty days' duration; and yet a large proportion of them recovered.

The treatment of the continued fever of the cold season in Bengal, amongst Europeans, is conducted on the same general principles which guide us in our management of those of the other seasons. We have to diminish arterial action, and relieve local congestions, by blood-letting, suited to the age, sex, and length of residence in India of the patient, and to the urgency of the symptoms. Having effected these primary objects, we proceed, without loss of time, to promote the secretions by a full dose of calomel with James's powder, followed in a few hours by active saline cathartics. Where no complications exist, one or two moderate

bleedings, followed by brisk purgatives, and tartarized antimony with nitrate of potash, will suffice for the cure, when these have been accompanied and followed by total abstinence from food, and by cooling diluent drinks. But where the brain, or liver, or both. have become the seats of diseased action, then our remedial means increase in vigour; the general and local blood-letting must be larger in amount, and more frequently repeated—the calomel with antimony, and the purgatives, must be more largely and more frequently exhibited-until relief from the fever, and from all local congestion, shall be obtained, together with a restoration of natural

secretion; for these are the only true signs of recovery.

It frequently happens in the course of this fever, that, with or without cerebral oppression, we have a sluggish condition of the bowels, with defective hepatic and intestinal secretions. In order to elicit these secretions, I was in the habit of combining plain scammony and the James's powder with the calomel, exhibiting them thus united over night, and following them up in the morning by a brisk draught of infusion of senna with simple jalap powder; or, where viscid mucus impacted the bowels, by a terebinthinate draught. In the fevers of children especially, we often find that the most alarming cerebral symptoms, with a doughy inelastic state of the abdomen, and torpid bowels, are removed at once by the night exhibition of medicine here mentioned, followed in the morning by the rectified oil of turpentine.

When the patient is seen after having been many days ill, without previous medical aid, congestion having proceeded unrestrained within the cerebral or abdominal cavity, or within both, blood-letting, general and local, calomel, sudorifies with diuretics, and strong purgatives, must still constitute our means of reliance; and we shall frequently have to exhibit the calomel so as gently to affect the system. These means, powerful as they are, will not, under the circumstances of neglect here contemplated, prove so prompt in their results as when applied earlier; but, nevertheless, they must be used, powerfully and repeatedly in bad cases, other-

wise the patient will be destroyed by the force of disease.

Where, on the other hand, we have evidence of irritation of the mucous digestive surface, we must determine to the skin freely by mild diaphoretics in the form of antimony or ipecacuanha, together with warm baths, relieving the bowels when necessary by

the most unirritating aperients.

Natives of Bengal were daily brought into the Native Hospital, in the most advanced and neglected stages of this form of fever; with tumid abdomens, confirmed stupor, and tongues black and dry as charcoal; and it was surprising how many of these poor people recovered under the following plan of treatment which, on a large amount of experience, became at length the settled management in such cases. On admission into hospital, the patient was put into a warm bath, while the entire surface of the body was purified by the free use of brush and soap—the head being shaved at the same time, and cold applied to it. A large sinapism was then applied to the abdomen; and where the state of the pulse admitted of it, a few leeches were applied to the epigastric region, or behind the ears, or to both regions, followed in a day or two by blisters to those parts. A very mild aperient draught was always administered on admission, aiding its operation by enemata.

A course of treatment was then commenced, of alkali, mild aperient and sudorific, all conjoined. The formula which I ordered was generally as follows:

This medicine was exhibited three or four times in the twentyfour hours, according to the severity of congestion, while the diet consisted only of iced barley water; and, as I have already stated, the result from such simple means was such as to surprise medical officers habituated only to the treatment of Europeans. The progressive recovery of sense, and gradual restoration of health, of persons who a week or ten days before had appeared in a hopeless state of disease, was very remarkable: but the natives of India will recover from wounds, and from advanced states of diseases resulting from fevers, which would destroy any European. The cases of fever here referred to were those of persons of the labouring class of natives, in whom fever had gone on unrestrained by any treatment for fifteen or twenty days, and in which, consequently, the period for more active measures of cure had long passed away. We were therefore content to work on gradually with nature—to see the patients improving; as any heroic endeavour to hasten the operation of cure would only have endangered their safety—so reduced were the powers of life.

To conclude:—it is scarcely necessary to urge that where this fever assumes the remittent character, whether in the instance of an European or native, quinine in proper doses, and administered during the remissions, will prove of signal advantage.

# THE ARDENT CONTINUED FEVER OF THE HOT SEASON IN BENGAL.

I have placed my notice of this fever last in order, because though very dangerous to persons of irregular habits of life, it is yet by no means so frequent of occurrence as the forms of fever previously noticed. A consideration of what has been said on the general influence of a high range of temperature at pages 37 and 38, will have prepared the reader for an unusually severe and dangerous disturbance of the nervous and vascular functions, under the united influences of direct, and of long continued solar exposure, especially where there has been intemperance of habit; and so we find it in the European soldier.

While examining the records at the Director General's Office, to which the kindness of Sir James Macgrigor gave me access, the clerk who assisted me in the examination said:—" Here, Sir, are twenty-two admissions into hospital at Berhampore in one season from apoplexy, and twenty-one deaths." They were examples of the most violent cerebral fever, amounting in some cases to actual coup-de-soleil:—"Where the line was to be drawn," reports Dr. Henderson, "between apoplexy and remittent fever, is here hard to say." The history of these fatal cases, as given by Doctors Henderson and Mouat, the able officers then in medical charge of Her Majesty's 13th Light Infantry, will afford a correct idea of this fever when in its worst form.

The soldiers consisted of older men who had recently returned from the first Rangoon campaigns, and recruits but recently arrived from England. The corps was marched in two detachments, cruelly, because unnecessarily, during the hot season, from Nuddea to Berhampore—a distance of sixty miles. The recruits had been drilled in the sun thrice a day before quitting Calcutta, so as greatly to injure their healths; in short, both classes composing the corps were in but indifferent health on commencing their unfortunate march; and though it was chiefly conducted during the night, with every care within reach of the medical officers, the effects were fatal in a remarkable degree, while "none of the natives were taken ill during the trip." Such of the Europeans as could not reach the ground of encampment by 9 A. M., "were seen to drop down and instantly expire: others less severely attacked were saved by timely and copious bleeding." But presently Dr. Henderson, the medical officer in charge, became seriously ill; and the day closed with a sick-list of sixty-three, and eighteen deaths, out of the right wing of the regiment alone; both the sickness and the deaths being

principally confined to the recruits.

The symptoms as described by Dr. Henderson, were:—flushing and swelling of the face, amounting in some cases to lividness—a parched and burning skin—a full and frequent pulse—difficult and oppressed breathing—giddiness and sense of fulness in the head, amounting to acute pain in some instances, with burning heat of the eyes, succeeded by loss of sense and motion, and faltering of the tongue—dilated pupils—twitching of the muscles of the face—subsultus tendinum, and involuntary evacuations. There was no stertor or paralysis, the symptoms being referred to a violent excitement of the nervous and vascular functions, with congestion of the cerebral vessels, the results of excessive heat acting on the irritable frames of hard-drinking and unseasoned European soldiers.

Such is an example of the severer form of ardent fever peculiar to the hot season in Bengal; and from a very interesting report by Dr. Milligan of the 63rd Foot, we find that like symptoms resulted from the exposure of his corps to the sun during a military funeral at Madras. "The greater number of the men," he says, "were in the prime of life; but there were amongst them some old soldiers who had served twenty years and upwards, some of it in the West Indies, and were much broken down by service and intemperate habits. The entire corps had just arrived from the Australian colonies, "where spirituous liquors can be had on easy terms." The regiment landed at Madras in the month of May; and from the date of the "untoward circumstance" of the funeral, "the hospital became filled with cases of fever." Two men dropped down and died on the very day of the funeral, and for several days afterwards the fever cases augmented considerably. "I have reason to believe," adds Dr. Milligan, "that the effects of this exposure to the rays of a vertical sun did not rest here, but laid the foundation of future mischief in assisting to originate fever, hepatitis, and dysentery, from which the regiment afterwards suffered much." Examination of the body after death was unfortunately impracticable in both the instances cited, as the sickness of most of the medical officers, and the harassing duties imposed upon all, left no time for the customary investigations.

In the Bengal instance the medical officers directed frequent bathing of the head and hands as a precautionary and preventive means, the march being conducted slowly, and the troops halted every hour; during the halts the men were urged to refrain as much as possible from drinking the indifferent water to be found along the line of march. When the actual disease set in, with "sinking of the vital powers, or a kind of collapse, a grain of opium was given to produce reaction, after the occurrence of which they were largely bled; and, with these precautions, though thirty men were admitted into hospital after the first march, none died."

Of Dr. Mouat's detachment sixty-three men were taken ill during the march, of whom fifty were recruits. "The disease," he says, "was a remittent fever of the severest kind." Subsequently Dr. M. describes "a cerebral disease arising from exposure to fatigue under the influence of a high temperature." In this latter instance blood-letting to the extent of fifty, sixty, and even a hundred ounces, was had recourse to; but "the remedy was worse than the disease; for though the first attack might be relieved, vet the subsequent symptoms required the same profuse evacuation for their relief; and from the previous loss of blood the patient was not in a state to bear such depletion, in fact, two men became convulsed, and died shortly after they were bled, and it was found after death that, though the heart was empty, the vessels of the head were loaded with blood." Here the brain would seem to have been overpowered by congestion beyond relief. Cold affusion often repeated, and continued cold to the head, were subsequently premised in the treatment; and after being put to bed, when reaction took place, the soldiers were bled to thirty or forty ounces, followed by full doses of calomel, purgatives, and leeches to the head. This latter treatment is proved to be more successful.

To the reports above quoted we have appended, in the Madras Medical Journal, important observations by the late distinguished Inspector-general of Her Majesty's hospitals in India, Dr. Murray. He "apprehends that there is an essential pathological difference between the morbid affections arising from exposure of the bare head to the direct rays of the sun, or from over exertion under great heat, and such as are induced by the influence of a high temperature in the shade. In the former case, strong excitement is at once produced in the organ affected, by the direct stimulus of the cause. Thus, in the case of exposure of the bare head to the sun, the scalp being protected by the hair, and peculiarly constructed, does not blister as the skin of other parts of the body would do if exposed to a strong solar influence; but its temperature becomes exalted, and the caloric, traversing it and the skull, gives immediate rise to active hyperæmia, and inflammation of the contents of the cranium. In the latter case, the application of the morbific agent being upon the general system, the cerebral affection which results appears to be induced in a different manner; at first there is nervous depression or collapse, and sanguineous congestion, upon which reaction follows." Then follow remarks of great value from this able officer, on the seat, symptoms, predisposing causes, diagnosis, necroscopy, cure and prevention of the cerebral disease; and it is with regret that I find myself precluded from doing more than quoting the summary of treatment: "The remedial means consist in blood-letting, general and local, cold applications to the head, cathartics, clysters, antimonials, counter-irritants, cordials, darkness, rest, quiet, and abstinence."

Dr. Murray urges that, excepting in cases of extreme necessity, the patient be not subjected to coercion such as that of the strait waistcoat, as the cerebral excitement would thereby be increased in the patient's efforts at release. All articles of clothing capable of obstructing circulation in any part of the body, but more especially about the neck, should be removed, and the posture should be such as to favour the free return of blood from the head.

During the fatigue and severe exposure of the Bengal detachments marching to Berhampore, it was remarked that while the Europeans suffered so fatally, "none of the natives were taken ill;" and such I dare say will generally be the case. I recollect one instance, however, where it proved the reverse of this, in the Governor-General's Body Guard of Cavalry, with which I served during the two campaigns of the first Burmese war. We mounted at 11 A. M. of the 12th of May, and, with one short halt only, we made a forced march of forty miles under as powerful a sun as ever shone in Burmah. The heat oppressed us almost beyond endurance, and many very powerful men, native officers as well as troopers, fell off their horses, vomiting, convulsed, cold, and bathed in a clammy sweat. Where a tree could be found I had the men placed under its shade, and they were dashed over with water. The detachment then pushed on, leaving several men thus affected on the road; but not one European officer, whether commissioned or non-commissioned, fell sick either during the march or after it. But we were all young, healthy, temperate in our habits, and well seasoned to the climate; and such men, it is well known, will bear with comparative impunity fatigue and exposure almost beyond belief.

By seasoning no more is here meant than the accommodation of the constitution, through the gradual operation of physiological changes, to extraordinary ranges of temperature, and which we everywhere observe to occur during the first few years of residence within the tropics. This, and no more, is meant by the term acclimation; in short, as stated in the article on climate, it is the ill habits of life of the European soldiery that give such fearful activity of power to tropical heat, and, but for these habits, its influence would be less felt by them, comparatively. In the two wings of the 13th Light Infantry, for instance, we hear of no officer's having fallen sick, excepting Dr. Henderson; and his fatigue of mind and body will alone account for his illness.

But I must now refer briefly to the fever of the hot season in Bengal, as it appears amongst Europeans in civil life. It cannot be necessary to enter into all the minute details of symptoms in the milder forms of this fever, seeing that its graver type has already been described; for it is always easy to bring the mind to the comprehension of simple and uncomplicated diseases, when once we have been rendered familiar with their most concentrated

and complicated states.

The type of this fever is usually continued, but sometimes remittent; it seizes with suddenness and violence, the heat, thirst, throbbing and frequent pulse, restlessness and racking headache coming in rapid succession. The complications are generally cerebral, with occasionally severe forms of gastric disturbance, as indicated by pain, oppression of the præcordia, and vomiting; sometimes the liver is involved. We have not here the tendency to collapse so characteristic of the true Bengal remittent fever; but the ardent rapid progress of the disease, together with the cerebro-spinal and gastric complications, constitute this fever of the hot season one of very considerable danger.

The appearances on dissection are such as might be anticipated from the symptoms during life; they consist of inflammatory and congestive states of the cerebro-spinal organs; and where the stomach, lesser bowel, and liver have been implicated, we perceive the morbid appearances common to our other fevers of the climate. Serous effusions are frequently found in the cerebral and abdo-

minal cavities, more rarely within the thoracic cavity.

The treatment consists in active blood-letting, practised to the extent of allaying the vascular disturbance, relieving the pain of the head, and reducing the temperature of the body; these are the

tests and the proof that the operation has been successful.

Leeches applied behind the ears, cold affusions, continued cold to the shaved head, and calomel and antimony, followed by very active purging, will generally confirm the relief of the patient, and complete his cure. In this fever, as in that of the cold season, the skin will generally be found harsh to the touch, compacted, hot and dry, so as to require a more free use of antimonial diaphoreties, united with cooling diuretics, than we find to be necessary in the fevers of the rainy season. When remissions take place, provided the head be relieved, quinine must be had recourse to for the purpose of preventing the recurrence of paroxysms; and should symptoms of exhausted nervous power, or of delirium tremens, make their appearance, from half a grain to a grain of opium, united with calomel and antimony, will soothe and procure sleep. In this, as in the ordinary remittents of Bengal, where an anæmic irritable condition of the brain has become manifest, symptoms occur which prove very embarrassing to the young surgeon; and the treatment of them by camphor, or by camphor

and opium, becomes an important practical question.

"For the congested state of the nervous centres in fever," says Dr. Billing, "syrup of poppy, or a few drops of laudanum, will produce an effect equal to that of a larger dose of opiate medicine in other states of disease." The same authority observes that narcotics are frequently useful during fever, nay necessary, as may be understood by referring to the essence of fever, and knowing that sleep is nature's restorer of deficient nervous influence. The obstinate wakefulness which takes place in some cases of fever must have a deleterious tendency to produce collapse, hence the benefit of gentle opiates.

Drs. Corrigan and Gordon, of Dublin, have for some time past exhibited chloroform internally to procure sleep in fever, to relieve pain in painter's colic, and to overcome nervous agitation

and restlessness in delirium tremens.

"The affections of the cerebral functions," says Dr. Gordon, "frequently assume great importance in fever; their derangement is sometimes indicative of disease in the respiratory apparatus, or in the digestive canal; more commonly, however, it is owing to the organic nervous influence produced by the fever on the brain and nervous tissue itself. One of the most prominent characteristics of this lesion is insomnolency, causing, or, at all events, followed by delirium, subsultus, &c." It is in such highly dangerous states of disease that chloroform internally administered produces the happiest effects.

Dr. Corrigan states truly that "the loss of sleep, if continued, is of itself sufficient to kill;" and he adds that, "if even the shortest sleep is procured some advantage is gained." His formula, of which one ounce is the ordinary dose, is as follows:—

Chloroform .		*	3v.
Pulv. Glycyrrhzæ	.*		3v
Mist Camphore		- 10	Tixes.

To afford sleep in fever, to relieve pain in painter's colic, the minimum dose exhibited is half a drachm of chloroform repeated every two or three hours. In delirium tremens the dose at first is a drachm, and is often increased to two drachms. It may with safety be given in larger doses, and unlike opiates it does not lock up the secretions.

Speaking of the use of wine in fever, Dr. Billing says, that the person who best understands the nature of delirium tremens will be the quickest to discern the propriety and necessity for the administration of wine and opiates in fever, crysipelas, &c. Even in inflammations occurring in debauched and debilitated

constitutions, it is absolutely necessary to give more or less

stimulants at a very early stage.

Where wakefulness threatens injury during the course of, or in the more advanced stages of tropical fevers, or where symptoms indicative of delirium tremens arise, opiates are not only safe but necessary; but the inexperienced medical officer must be on his guard against error in diagnosis. He should feel his way with small doses at first, given at intervals; for, as remarked by Dr. Latham, it is a fearful thing to strike a heavy blow in the dark,

where the alternative is of such magnitude.

It is in this fever that cold affusion exerts its greatest power in abstracting heat and subduing nervous and vascular excitement. I remember the case of an officer who had been much exposed to the direct rays of the sun, and who was not subjected to treatment until the advanced stage of his fever. His sufferings and distress were beyond description, and he appeared to be rapidly sinking. His reduced state rendered blood-letting in any form inadmissible; but, the skin being of a pungent dry heat, I had him soused with cold water, and the relief was, excepting in one other case already narrated, the most surprising of any that I have ever witnessed. This person to all appearance lost, was, through cold affusion chiefly, speedily restored to health.

## NOTE ON THE FEVERS OF THE NATIVES OF INDIA.

The subject of the diseases of the natives of India deserves a separate treatise, and I trust that the time is not distant when we may hope to possess one worthy of so important a question. But the scope, purport, and very title of this work do not admit of any further discussion on the subject than the most cursory. A knowledge of the diseases of Asiatics is intrinsically of great importance, while, contingently, it will present an interesting and extensive field of comparison with the diseases of Europeans, and their causes.

British India, divided into four great portions—that of the Ganges to the east, of the Indus to the west, of the Dekhan to the south, and of that bounded by the great Himalayan range to the north—is inhabited by nations differing in religion but descendants of the same great family; they are distributed over the vast extent of what is termed the Peninsula of Hindustan—the widest extent of empire with the smallest and most easily defensible line of frontier of any existing on the globe.

There is reason to conclude, from the notices by the Greek authors, that no change whatever has taken place in the institu-

tions, character, or manners of the Hindus, from the age of Alexander to the time when they were discovered by the nations

of modern Europe.

The extreme length of Hindustan from south to north is 1,880, while its breadth from east to west is 1,620 miles, or 27 degrees of latitude and 25 of longitude, comprising 1,280,000 square miles;—all its features being on the most magnificent scale. Washed by the Bay of Bengal to the south-east, and by the Indian Ocean to the south-west, it is bounded to the north and east by mountains of prodigious elevation, from whence the two great rivers of India have their sources, and flow to opposite points of the continent.

Varying in physiology of constitution as they may be divided by geographical distinctions of latitude, quality and elevation of soil, or by differences in diet, the natives are separated, Hindu from Hindu, and Hindu from Mahomedan, by impassable barriers of caste and of religion; yet, under British rule, this wonderful aggregation of one hundred and fifty millions of subjects has attained to a degree and kind of union, such as no antecedent rulers, whether ancient indigenous Hindu, or more modern conquering Mahomedan, ever could conceive, far less accomplish.\*

Out of this enormous mass, every description of person enters eagerly into the various public employments presented by the British government, whether civil or military, and natives of every caste and class are thus brought into familiar and friendly communication with their European superiors. Not only is this the case, but natives, both Hindu and Mahomedan, undergo laborious and protracted courses of education, the better to qualify themselves for the various public duties offered to them; and here, to the honour of the medical profession, it has taken the lead, and accomplished a noble triumph of education. Always prominent in every endeavour to advance the moral and intellectual condition of the people, youths may now be seen by the hundred, trained and instructed by our medical officers at the three Presidencies, who but a few years back, had they touched a dead body, would have become outcasts. Now these youths are seen to go through years of the dissecting room, and to leave their colleges able physicians and expert surgeons.

The relative proportions of the followers of the two dominant religions of India, the Hindu and Mahomedan, may be taken as about eight of the former to one of the latter, for the whole of Hindustan; Pegu and the Island of Ceylon being inhabited by Budhists. It is estimated that, throughout India generally, there are 3,400 Natives to one European; but insignificant as this proportion appears, the presence of the European offers to the Hindu

<sup>\*</sup> The Mahomedan conquest of India was completed A. D. 1215.

and Mahomedan the only hope of ever attaining to truth in religion or in science.

The complexion, stature, bulk, and weight of the body vary in the several nations, as they may inhabit the south or the north of India, as they may occupy countries near to or far from the Equator; the advantages in all the above respects being generally much in favour of the northern nations. But whether of high or low stature the trunk of the Asiatic, whether Hindu, Mahomedan, or Budhist, will be found slender, as compared to that of the European. The Siah-Posh, an undoubted branch of Hindu race, and who inhabit the cold mountain region of Kohistan have, according to Sir Alexander Burns, arched eyebrows, a fair complexion, blue eyes, and Grecian features, while, on the contrary, the descendants of fair Persians and Affghans have, in the course even of a few centuries, become of darker and darker shades as we trace them into the farther south of India. Generally speaking, the Mahomedan, whether of Persian, Affghan, or Baluchi descent, will everywhere, but in the north especially, prove the fairer as compared to the Hindu, but this distinction will be found to have gradually disappeared under a foreign climate, and the fairer Mussulman will have become more and more dusky with advancing generations.

The various natives inhabiting Hindustan appear to differ from Europeans less in feature than in the more trifling circumstances of colour, size, and form. They are nevertheless moulded by a great variety of climates, localities, habits of life, diet, occupation, &c., so as to constitute in reality a people varying exceedingly in moral and intellectual qualities, in physical powers and appearances. The medical officer will therefore have peculiarities to investigate in almost every state and province throughout the whole extent of our Eastern possessions. Placed under few of the rigorous necessities which, in northern latitudes, impart energy to the mind and body of the inhabitants, the people of the east are handed over by the richness of their soil, the power of their sun, and by climate and institutions of religion, to a state of indolence and inactivity as compared to western nations.

Of Caucasian origin, but exposed during countless generations to the same succession of external influences of high temperature, and corresponding habits of life and diet, receiving the same reiterated impressions peculiar to climate and religion, an acquired temperament is formed, which constitutes the Hindu and Mahomedan, men differing widely, morally and physically, from Europeans, while from each other they differ principally in religion and in diet. These general causes, together with the premature development of the generative function, produce an excitability of the nervous system, diminished volume, enervation, and relaxation of the muscular system as compared to Europeans, all which dispose the natives of India generally, and those of Bengal Proper in particular, to tetanic affections under wounds, surgical operations, and the impressions of cold and damp.

Generally speaking, the agriculture and the mechanical arts are in but a rude state with the Hindus, both being supposed to have remained stationary for two thousand years or more. "India," says Dr. Royle, "vast in extent and diversified in surface, is remarkable as the cradle of one at least of the nations who earliest practised the arts and cultivated the sciences which characterize civilisation, and from whence these travelled to the west, and perhaps also to the east. Its present inhabitants continue to venerate sciences which they know only by name, and to practise arts of which they know not the principles, and this with a skill not only remarkable for the early period at which it attained perfection, but also for the manner in which it has remained stationary for so many ages. This can be explained by the fact that the son was unable to add to the manual dexterity of the father, and could not improve an art which he knew only as a routine process. But when commerce was in its infancy, or dealt only in the most precious commodities, these arts could not have been practised unless India had contained within itself all the raw materials which art could convert into useful articles or elegant ornaments. Without cotton the so-called 'webs of woven air' could have had no existence. Without numerous woods, barks, and flowers, dveing could not have been practised, and calico-printing would probably not have been invented. If an Indigofera had not been indigenous, indigo would never have derived its name from India, nor have afforded us the proof, in the stripe of mummy-cloth, of the early commercial intercourse between its native country and Egypt. Neither would sugar have been arranged by the Greeks with honeys, nor the Indians described as those who 'bibunt tenera dulces ab arundine succos,' unless they had had the cane-like Saccharum as a plant of their country. Neither in Persia would the proverb of 'giving an Indian answer' have been considered equivalent to a cut with an Indian sword, unless the Hindus had possessed the ore which enabled them to manufacture the far-famed wootz steel; and gunpowder is likely to have been invented at an early age only in a country where 'villanous saltpetre' is abundant." Another ancient saying amongst the natives, repeated to this day with glee by the Sepoys of Bengal, exhibits alike the warlike character of the people, and their readiness at all times for mercenary service :-

<sup>&</sup>quot;Jis ka Deg, oos ka Teg."

This popular and admired proverb is translated by Brigadier Hodgson of the Bengal army, as follows:—

"He who can keep the pot boiling, Will never want for trenchant blades."

The mention of an Indian answer, wootz steel, trenchant blades, and gunpowder, brings us naturally to consider the native

army of India, the proper subject in the present article.

The native soldier of whom alone I can pretend to personal knowledge and acquaintance is-the Sepoy of Bengal. He is brave, obedient, cheerful and patient under labour, fatigue, and privation, abstemious by religion and habit, honest and amiable in all his dealings, and possessing a very high sense of military honour; attached to his European officer, and grateful for kindness shown him, he never abandons his colours; and when kindly and justly treated, and properly commanded, he has never disappointed the reasonable expectations of his commanders. He is cheerfully resigned in sickness, never exhibiting any unmanly qualities under suffering; he thus stands forth an object deserving the special and high consideration of his European medical officer. A fine example of the kindly feeling of the Sepoy towards his surgeon was afforded after the battle of Mahidpore, when the Commander-in-chief of the Madras army, Sir Thomas Hislop, on visiting the hospitals was addressed by the wounded Sepoys, who "made a personal request of his Excellency that he would now take care of the surgeons who had been so kind and attentive to them." Colonel Blacker in his history of this war, and referring to the "unanimous petition" of the Sepoys, adds, that "unassuming merit could not have a more gratifying and appropriate reward, intrinsically more estimable than any formal commendations flowing from authority."\* That the qualities here recorded, of the men forming the native army of Bengal, are derived from the domestic circle and habit of civil life, shared by the Sepoy with his brothers, is as certain as that all the qualities of a great military commander are civil qualities. A soldier is a citizen and something more; the addition being but the result of military discipline and military habit. To form a just estimate of the soldier we must everywhere trace him back to the recruit.

The natives of India generally are amiable and respectful in their intercourse with Europeans; and towards no class of their rulers do they show more real kindness of feeling and gratitude than to such of our medical officers as may prove deserving.

<sup>\*</sup> Colonel Blacker is quite philosophical and even kind on the subject of handing over the surgeons of his army for rewards to the bare contemplation of their own merits; but he does not say that, in his own person, he would prefer such mode of recompense to the companionship of the military order of the Bath.

Personally, I must say that I shall ever look back to my early connection with the Sepoy of Bengal, and to my later friendly communications with the native inhabitants of Calcutta, with

sentiments of sincere regard.

Medical officers in the East Indies, holding as they do military commissions, are principally employed in the duties of the native armies; but a certain small proportion is allotted, or "lent," to the civil department of the state, for conducting the medical duties of civil stations. But the great body of the medical corps, numbering for the three Presidencies of India about eight hundred officers, are under the direct and immediate orders of the Commander-in-chief, and are thus placed much more in contact with

the native soldiery than with the civil population at large.

Nor is the social position of the medical officer less befitting a man of science. Entering the Indian army with rank corresponding to that of lieutenant, his position is at once established as that of an officer and gentleman; and it depends on the conduct and character of the individual how he shall uphold such privileges. To secure the confidence and respect of his brother officers of the army, and to become distinguished in his profession, he must possess knowledge, indefatigable industry, judgment, energy, resolution, address, and an untiring patience; and to all these qualities he must add a courage not fed by excitement of the heat of conflict, or buoyed by any hope-a courage that does not look to rewards, a courage of the highest order: in truth "the two o'clock in the morning courage," which Napoleon prized so highly, is quite as necessary to the surgeon as to the captain. Towards natives of all classes his conduct should be just, kind, and liberal; while in his intercourse with superiors and equals of his own country, his conduct should be that of a modest unpretending man of science. He must learn that familiarity in the mess-room or elsewhere is not friendship, and that such conduct is never supported by real friendship. And let not these observations be considered irrelevant, for they have in truth much to do with moral as with professional effect.

Excellent as are the native qualities of the Sepoy, his usefulness to the British government depends mainly on good discipline, aided by a just and kind treatment on the part of his European officer. Taken from the middle class—the pith and marrow of British society, whether for peace or for war—the officer of the Indian army is at once a gentleman and a soldier by profession; but he is neither an amateur nor a dilettante soldier. He does not enter the army to make it an amusement rather than a business; he is not a monied idler; and wherever he is placed in command, he is not found wanting. Freed from the weaknesses of class or caste, which for a long time have damaged certain branches of

the British army-freed also from the democratic infusions of the armies of continental Europe; he stands in a middle position between the two extremes; and if at any future time he is to exhibit well-intentioned weakness and incapacity, it will be from causes which are not now in operation. After fifty years' service in the British Army, Robert Jackson declares that "for any great purpose of peace or of war, the middle classes of the empire are the saviours of their country." From the earliest formation of the Indian army, its officers have remembered in a proportion which does them honour, that in reading, and the command of one's reading, consists education, and that no officer can deserve the name without such mental discipline. On the staff too, we seldom find the offensive peculiarities against which Shakespeare has warned us in your "dog in office." In short, the officers of the Indian army do not regard military life as a mere dashing and casual pastime, but as "a serious and permanent profession." It follows from what has just been stated, that the medical officer who has the good fortune to serve in the Indian army will find the criticisms of its officers on his general and professional conduct, matters not to be disre-

But with all these advantages, there still remains a source of affliction and of debility, to the medical officer of Bengal especially. The medical department is there ruled, as in the other presidencies, by the three oldest men in the service. To attempt the defence of such an institution were to build its tomb: to endeavour to instruct members counting forty years' service under tropical influences, or to make them understand the true purpose and object of their official calling; or even to comprehend their own true interests, were about as hopeful as to send forth missionaries to convert the Pope to Protestantism, or the Emperor of Russia to representative government.

The strength of the Native armies varies with the e

The strength of the Native armies varies with the exigencies of the state; that of Bengal, in 1825, mustered 152,843 men of the several arms, infantry, cavalry, and artillery; while, in 1832, the same army was reduced to 78,346 men

The following is stated by Colonel Sykes to be the distribution of castes in the Bengal army:—

Christians.	Mahomedans.	Brahmins.	Rajpoots.	Inferior Castes.	
1.076	12,411	24,849	27,993	Hindus. 13,920.	
Grand Total 80 249					

The average admissions into hospital of this army during twenty years was for each soldier one admission in two years; while the average mortality for the same period was 1.79 per cent., and the invaliding 1.5 per cent. per annum.

The strength of the Madras army varied from 71,488, in 1826, to 48,571 in 1837. The average mortality in this army during twenty years, was 2.09, and the invaliding, 1.96 per annum.

In the Presidency of Bombay, the strength of the Native Army varied from 49,873 in 1844, to 25,782 in 1833. Six-eighths of this army are natives of Hindustan, there being in the proportion of eleven Hindus to one Mahomedan; and low easte men constitute about one-eighth. The mean percentage of deaths during twenty years gives 1.29—a remarkable degree of health—while the invaliding during the same period was 3.31 per cent. per annum.

The mean mortality during twenty years for the whole Native Army of India, as given by Colonel Sykes, is 1.80 per cent., per annum, and the invaliding as 1.93—exhibiting a remarkable condition of health even in troops serving in their native climate.

It is with Native as with European troops in the East, fevers and bowel complaints constituting by far the greatest amount of sickness. The maximum of admissions into the hospital of Bengal troops occurred in 1842, when 98,936 men were treated medically out of a strength of 113,020; the minimum occurring in 1827, when only 30,903 were treated medically out of a strength of 130,303 men. During twenty years the average admission into hospital was for each soldier, one admission in two years.

In the three Presidencies of India we have the following mortality amongst the British and Native soldiery per cent:—

Bengal. Madras. Bombay
European Troops . 7:38 . 3:846 . 5:07
Native Troops . . 1:79 . 2:095 . 1.29

Thus a European regiment is renewed in Bengal in ten years, in Bombay in twelve years, and in Madras in seventeen years.

To account for the circumstance that in the Madras Presidency the Native mortality is greatest while the European mortality is the least, Colonel Sykes states that the constituents of the Madras Native Troops are the reverse of the other two Presidencies. In the Madras Cavalry there are six to seven Musulmans to one Hindu; and in the infantry there is one Musulman to  $1\frac{1}{2}$  or  $1\frac{3}{4}$  Hindus; but amongst the latter there is a considerable number of low castes, without prejudices of caste; therefore the majority of the Native troops of the Madras Army can eat and drink like Europeans, and the mortality returns show us that they suffer from cholera as much as Europeans, and that the mean mortality from all causes is 2.095 per cent., or more than  $\frac{3}{4}$  per cent. beyond that of the Bombay army for twenty years.

"I never followed," says Colonel Sykes, "a farinaceous or vegetable regimen myself in India, nor do I recommend it to others; but I ate moderately and drank little, and I have a strong conviction that much of European disease in India is attributable to over-stimulus, and that the mortality among the European troops will not be lessened until the European soldier is improved in his habits, until he is made to understand that temperance is for the benefit of his body, libraries for the benefit of his mind, exercise for the benefit of his health, and savings' banks for the benefit of his purse. The climate of India is less to blame than individuals; for in the case of foreigners, the people of a country being healthy, they should, to a certain extent, conform to the habits of

the natives to be healthy also."

The same authority adds:-"The Natives of India are generally considered to be very temperate in their habits; but it is quite a mistake to suppose that they all live upon farinaceous or vegetable matters, and do not drink fermented liquors; it is equally a mistake to suppose that the general food of the people is rice, which is only very much the case in low lands subject to inundation, and along the coasts. In the interior rice is generally so much dearer than the bread grains, of which there are many (wheats, millets, the genera holcus, panicum, paspalum, &c. &c.,) that rice is rarely consumed, at least in Hindustan and the Dekhan. The Hindustance soldier lives almost exclusively upon unleavened cakes of wheaten flour, daily baked upon an iron dish, and washed down with water. On the other hand, all Mahomedans, and all low caste Hindus are consumers of animal food, spirituous liquors, opium, ganja, or hemp-water; and many castes of the Sudras, the Mahrattas, for instance, eat mutton and fish, when they can afford to do so; but meat is not essentially necessary to health and strength. Liebig says, that only those substances can possibly be called nutritious which are capable of conversion into blood, and that farinaceous food has also this nutritious principle in high degree. The truth of this profound assertion of Liebig is established by the food of the great majority of the Native soldiers of the Bombay and Bengal armies. I have shown that six-eighths of the Bombay army consist of Hindus, and considerably more than half of the whole army are Hindustanees. These men never taste meat, fish, or spirituous liquors, but live, I may from personal observation venture to say, almost exclusively upon unleavened cakes of wheat, or other cerealea, baked upon an iron dish, and eaten as soon as cooked."

The natives of India are very subject to the influences of the various climates contained within the British empire in the East, and none more so than the Sepoys of the Bengal army, recruited as they are almost exclusively from the plains of Upper Hindustan; but wherever circumstances admit of the use of artificial means, both Hindus and Mahomedans, though born and bred on the soil, take greater precautions against both heat and cold than the stranger European. "To brave the climate" forms no part of native disposition or habit.

The climate of Bengal proper is, for instance, most prejudicial to the health of the "up-country" Sepoy; for, although according to Colonel Henderson, but one-fourth of the Bengal army is stationed there, the deaths of that fourth are more than a moiety of the whole mortality reported. Much of this mortality is doubtless due to the excessive humidity and to the malarious character of the climate; but much is also due to the great change of diet from wheaten bread to rice, which latter the Sepoy will use, not because he prefers it, but because it is by so much the more cheap. The native dealers in grain, taking advantage of the Sepoy's frugality, offer cheaper and worse qualities of rice, or mixtures of rice and wheat flour; so that within two or three years' residence in the lower provinces, what with the injuries from climate and the inferior and imperfect nutriment—the robust Sepoy wastes, and falls a sacrifice to the fevers, dysenteries, and diarrheas of an unnatural climate; and thus it is in many of the low malarious countries throughout India.

As with the European soldier, so with the sepoy of Hindustan Proper when transferred to the malarious low regions; both shared a common fate when, through the feebleness and incapacity of General Morrison of Her Majesty's army, the united force under his command was planted by him in the marshes of Aracan, in the first Burmese war. Those native soldiers who did not perish on the spot sank soon afterwards from general cachexia, and enormous enlargements of the spleen, followed by dropsical effusions, colliquative white flux, or lientery:—in fact, all perished alike—

In Ava again, in the same war of 1824-25, where I was witness of what happened, and where the climate generally was good, the Sepoy, by a proper diet and clothing, was preserved in comparative health through two severe campaigns; but the British soldier, the instrument of the power and glory of the Government, was suffered by its criminal negligence to perish miserably through

the native and the European.

Dr. Finch, of the Bengal army, in an able report on the effect of climate on the health of the native army of that Presidency, gives the following comparative view of the influence of the climate of Hindustan, and of Bengal Proper:—

	Years.	Average Strength.	Average Sick.	Average per cent. Sick to Strength.
Hindustan	${1835 \brace 1836 \brace 1837}$	. 753 .	854 .	. 46
Bengal Proper	\begin{pmatrix} 1838 \\ 1839 \\ 1840 \end{pmatrix}	. 899 .	649 .	. 79

"During three years at Mynpuree," he says, "the corps lost by death but twenty-six men; while during its stay in Bengal

Proper, the casualties amounted to not less than 203!"

The manner of locating native troops, with reference to the country of their birth, the period of service in climates which are unnatural to them, and the proper time for relieving corps and divisions, so as to preserve the soldier's health, are matters that as yet have not received from the authorities the attentive consideration due to them.

As with the European, so with the native soldier, the season of greatest amount of disease is, from June to January—cholera being most prevalent in April, May, and June. During the rainy season fevers, remittent and intermittent, prevail the most; and, towards their decline, bowel complaints which, in low malarious tracts prove very intractable, become frequently associated with splenic cachexia. Again—as with the European, so with the native soldier—fevers remittent and intermittent generally constitute the first inroad on his health; after which successions of recurrences ensue, followed sooner or later by enlargement of the spleen and general cachexia—the disease which closes life being dysentery or diarrhea.

Amongst the natives in civil life, the Bazar preparations of arsenic and mercury, so freely administered by the Bazar empiries, prove a source of much broken health, causing severe rheumatisms, diarrhœas, and dysenteries, especially during the cold season. Of all this I observed much in the Native Hospital of Calcutta, of which institution I was surgeon during ten years.

The fevers of the natives, whether remittent or intermittent, present the same events in the same succession as in the European, differing only in degree; and in the Indian army the patient, like his western comrade, has the advantage of coming under the care of his surgeon on the instant he falls sick; and within the first twenty-four hours much may be done to place him in a position of security, and to prepare him for convalescence.

I generally began the treatment by an emetic-cathartic, composed of from one to two grains of emetic tartar, from three to four grains of calomel, added to two scruples or a drachm of compound jalap powder, according to the age and strength of the patient. This powder placed dry on the tongue, and washed down with water, produced a speedy and free action of both stomach and bowels, followed by copious perspiration, and a more or less complete remission of all the symptoms. Generally speaking the patient was next day in a condition to use bark, quinine, or some of the native bitter tonics to complete the cure. Such of my servants as had served me long, and been with me when in the army, always asked for what they called "the fever powder"

whenever seized with the disease, and the other servants soon learned to follow their example; and my European friends, whether civil or military, when going into camp, or on a long march, always carried with them a supply of the same medicine for the use of their native establishment. It continually happened that, in the milder cases of fever, no tonics of any kind were necessary, the triple action of the emetic-cathartic and sudorific proving sufficient for the cure; and this was especially the case in the sudden and acute febrile seizures of the hot season. I may here add that mercury in any form, beyond the first moderate dose, I seldom if ever administered. When heat at stomach, with urgent thirst, are complained of, the greatest relief will be obtained from the use of saline effervescing draughts.

But in malarious countries adynamic forms of fevers, both remittent and intermittent, will occur, accompanied very generally by abdominal congestions with enlargement of the spleen. Here, in addition to bark and other tonics, warm aromatic aperients, chalybeates, improved diet, and change of air must be had recourse to. The Sepoy of Upper India is bound by warm ties to his family, to which he never fails to remit his pecuniary savings. In sickness he eagerly looks to rejoining his relatives; and, where the exigencies of the service admit of it, such indulgence should always be recommended by the medical officer, whatever the colonel

may think.

In treating the fevers of the natives we must not expect to find in them the sustaining powers of constitution of the European;—acute disease subsides in the former much earlier than in the latter, requiring, therefore, antiphlogistic measures of much less energy in the case of the Asiatic, while a much earlier recourse may be had to quina and other tonics in conducting his cure. Whenever we feel satisfied also that acute disease has been overcome, an improved diet may be allowed at an early period—the native constitution not bearing a severe and protracted abstinence; and condiments may be allowed, especially during the rainy season.

Finally:—Let the surgeon be careful that the Sepoy is not discharged from the hospital too soon, out of a small, narrow, or mistaken notion of military efficiency, or of what an unreflecting ing adjutant or crack colonel may designate, "keeping the effective list complete." The effectiveness of convalescents amounts only to deceptive numbers borne on the muster-roll; and such sham—efficiency will bring disgrace on the medical officer, stamping him at once with want of knowledge, want of firmness, and want of humanity.

## ACUTE DYSENTERY OF BENGAL.

This is at once a dangerous and a frequent disease throughout our intertropical possessions;—so much so that no medical officer should be permitted to enter on the duties of the public services without a careful previous study of its nature and treatment. Dysentery has at all times proved one of the most severe scourges of our fleets on foreign stations, and of our armies in the field, even when campaigning in temperate regions. It is truly designated by William Fergusson an army disease. "The soldier," he says, " when in the field may escape fever, but never dysentery, if he lie on the ground." It is the disease of the famished garrisons of besieged towns, of barren encampments, and of fleets navigating tropical seas, where fruits and vegetables cannot be procured. Dysentery has occasionally been found to assume a malignant character in low damp situations, so as to prove very deadly; but on the sufferers being removed to elevated sites, and a better air, the disease has been rendered amenable to treatment.

In tropical climates dysentery, though most prevalent during the hot and rainy season, will be found to exist at all periods of the year; but in European countries it is almost exclusively a disease of summer and autumn:-it stands thus associated with a

high range of temperature, and atmospheric changes.

Colonel Tulloch states, that out of an aggregate force of 25,433 men of his Majesty's army serving in periods of eight and ten years respectively, between 1823 and 1836, in the stations of Calcutta, Chinsurah, and Berhampore, all in Bengal Proper, there occurred 8,499 cases of dysentery and diarrhœa; and though the years within this return were subsequent to the first Burmese war, still the proportion of dysenteric cases is excessive, making every proper allowance for such unfavourable influence. The climate of Lower Bengal has always been very unfavourable to European health, as compared to Upper India; but making allowance for all circumstances, including the ill habits of life of the soldiery, the amount of sickness from dysentery and diarrhœa here exhibited is enormous.

In the Presidency of Madras again, out of an aggregate British force of 82,342 men, serving there from 1842 to 1848, there occurred 10,531 cases of dysentery, and 9,189 cases of diarrhœa, making a total of 19,720 cases of bowel diseases, exclusively of cholera. It thus appears that, next to the malarious fevers of India, bowel complaints are the most prevalent diseases, while the dangers to health and to life from these last are even greater than from fevers. There were admitted into the European General Hospital of Calcutta, according to Dr. Macpherson, in the years 1830 to 1850, 2,044 cases of dysentery, of whom there died 457, or 22.3 per cent. The extremes of mortality were 14.8 in 1833, and 34 in 1845.

Robert Jackson regards dysentery as "one of the most important of the maladies that occur among troops, particularly in the West Indies, where, in some of the islands it amounts to onehalf, even to more than half of all the forms of acute disease which appear in the hospital return of sick." This great authority adds: -" It is dangerous in itself; -more fatal, in fact, among the military in the West Indies, either primarily or secondarily, than any other, the concentrated fever, as incident to strangers, excepted." The losses of the French army in Egypt were greater by dysentery than by the plague—the deaths by the latter disease being 1,689, while by the former they amounted to 2,468. During two years and a-half only of the time occupied in the war in Portugal and Spain, the British army, according to Sir James M'Grigor, lost 4,717 men by dysentery. Of the diseases which destroyed the European portion of the expeditionary force in Ava, during the first Burmese war, dysentery-scorbutic dysentery -was by far the most prevalent and fatal disease.

PREDISPOSING CAUSES.—Whatever causes predispose to fever will exercise a like influence in producing dysentery; but more especially the following:—The very duties of the fleet and of the camp, which necessarily expose seamen and soldiers to wet and cold, suppressing the secretions—to high ranges and to sudden and violent changes of temperature, causing checked perspiration especially;—the use of crude, ill-prepared, indigestible, or otherwise unwholesome diet—excesses in the use of wine, spirituous liquors, or tobacco—the use of impure water—retained excretions—fatigue and privation—endemic, epidemic, and malarious influences—previous fevers, of the remittent and intermittent types especially—diseases of the liver or spleen;—all these circumstances will, severally or individually as may be, dispose the subject to the invasion of dysentery. The causes of scorbutic dysentery are:—Deprivation of fresh meat, vegetables

and fruits—sameness in diet—crowding—mental depression—hardship and fatigue—exposure to damp and foul air—the use of impure water.

SYMPTOMS AND PROGRESS.—The immediate and urgent symptoms in tropical dysentery are—sense of constriction of the skin, followed by irregularly developed febrile movements, more or less marked according to the season of the year, the length of residence in India, the constitution and previous habits of the patient. The amount of attendant fever is very variable;—in some cases it hardly excites attention, while in others it amounts to heat and dryness of skin, flushed face, furred tongue, and hard frequent pulse. Fever is most considerable in the hot season; while in that of the rains it is less declared, the tendency being here as in remittent fever, to congestion and collapse. In some forms of malarious dysentery there is considerable fever, with but little hepatic complication comparatively;—at least it proved so in China. But here, as in all the diseases of the stomach and bowels, the power of the pulse rapidly fails as the disease advances.

Soon the patient experiences, during from twelve to twentyfour hours the uncomfortable feelings attending upon frequent loose discharges from the bowels; when at length irregular pains, at first of a griping nature only, along the course of the large intestine, becoming gradually more severe, shooting, or "cutting," as the soldiers designate them, with sense of heat ascending from the rectum, and pain extending sometimes to the hypogastrium, until the whole abdomen becomes involved in soreness and feeling of distress, accompanied by frequent purging, griping, and straining, all aggravated during the night and early morning, and leaving behind them the wearing sensation that there has always remained in the bowel something which has yet to be discharged. At this time the discharges are generally scanty, and consist of mucus and blood, or bloody slime as the soldier calls it; and this is a matter of the utmost importance to investigate, as from the character of the discharge we derive important information as to the extent and nature of the disease.

Towards a right estimate of the condition of the tubular, as of the associated solid viscera, we may also receive much aid from a careful exploration of the cæcum and colon, including its sigmoid flexure; and a careful examination of these regions, as of those of the liver and spleen, should never be neglected; for pain, fulness, or thickening of the bowel may always be detected, unless the subject be inordinately stout. When pain and fulness are absent from the regions named, and there is, nevertheless, much local distress, such as urgent and constant tormina and tenesmus, with strangury, the seat of inflammation and thickening will be found to be centered in greater degree along the rectum; but this circumstance does not render the case less severe, or less difficult of cure.

The matters voided in the dysentery of India consist at first of liquid fœces, generally of unnatural colours, streaked with blood or mixed with blood and mucus. If the disease advances in the absence of treatment, or in defect or insufficiency of it, then we shall find all the symptoms, constitutional and local, much aggravated, while the intestinal discharges contain more of blood and mucus, together with shreds or larger sloughs of mucous membrane, so as often to assume the appearance of putrid blood, or of the washings of raw meat. Scybala are rarely seen in the dysenteries of the East. When to symptoms such as these are superadded, a pungent heat and dryness of the skin, with furred tongue, an urgent thirst, high-coloured and scanty urine, an increased frequency of the pulse, and an increased urgency and frequency of call to stool-all augmented towards night-they indicate considerable danger. When, again, patients suffering from the tropical dysentery present the further retrogression of a cold clammy surface, a cadaverous smell from the body and intestinal discharges, an anxious and oppressed countenance, and a sinking pulse with hiccup, and motions voided involuntarily, the case is generally hopeless.

Such are briefly the symptoms of acute dysentery, which will be found to vary in degree according to the severity of the inflammation; and of this last the medical officer will best judge by an attentive observation of the constitutional symptoms, together with a careful daily examination of the nature of the discharges from the bowels:—this, indeed, cannot be done too often, or too minutely.

The duration of dysentery varies with the force of the predisposing and exciting causes, the age, constitution, length of residence in hot climates, the season and other contingent circumstances. Some cases will go through their course even unto death within three to five days; while others will be received into hospital having suffered during periods from five to twenty days, with appearances the most unfavourable, and yet they recover.

The progress of the disease will, in fact, depend on the condition and circumstances of the individual case, past and present. The scorbutic form of dysentery may be recognised at once by the dusky hue and dull melancholy expression of the countenance—the livid complexion of the surface generally, but especially of the extremities, which are cold and spotted over with petechiæ, which the least scratch will convert into an ulcer. The tongue is dark in common with the gums, which latter are spongy,

and bleed on the slightest touch, or blood oozes spontaneous from them. There is a disrelish for the previously uniform disand especially for the use of salted meats, while there is an urge craving for vegetables and subacid fruits of every description. The pulse is feeble and frequent;—in short, all the symptom general and local, indicate an extreme and aggravated cachexia a dissolved, diseased condition of the blood. The purgit soon exhibits the true scorbutic character, the discharges being composed principally of grumous putrid bloody matter, and even the urine has a bloody hue.

The progress to recovery from dysentery implies a healing ulcerated surfaces. Dr. Parkes accords with the German ana mist Sebastian, and others, as to the fact of the reproduction intestinal mucous membrane; while Dr. Bleeker regards cical zation as the normal termination of dysentery; but the numl and extent of the cicatrices "may withal produce a condition the colon which renders it unfit properly to perform its function Cicatrices, be they ever so completely formed, cannot make go the loss of the mucous membrane with its absorbent and secretion organs."

COMPLICATIONS.—Dysentery is found in all climates complicate readily with the prevailing fevers, and within tropics it is frequently associated with fevers remittent and int mittent:—it is also occasionally complicated with scurvy. In East Indies we find that, either from the beginning, or during progress of the dysentery, diseases both functional and structurare apt to arise in the liver.

Amongst the European patients in the General Hospital Calcutta, Dr. Macpherson found that, out of an hundred and six acute cases, the liver was altered in eighty-four cases—containing abscess in twenty-one cases—being enlarged in forty cases, & while in fifty-five cases of chronic dysentery the proportion hepatic diseases was nearly the same. This able officer stathat, on a comparison of acute with chronic dysentery, the liver more frequently altered in the latter, that abscess is about equal frequent in either form, that in acute dysentery the liver is formently enlarged and soft, while in the chronic it is more general small and indurated. He adds that the stomach and small testines suffer also more frequently in the chronic form, and the mesenteric glands are more frequently altered in it.

In the Madras Presidency, out of fifty-one cases of acute dysetery there were twenty-six hepatic abscesses; and in Bombs out of thirty cases of acute dysentery, twelve cases of hepa abscess occurred. "It is worthy of remark," says Dr. Macph son, "that the liver has been found, in the General Hospital

Calcutta, to have been altered in 118 out of 215 cases; in the Medical College Hospital in thirteen out of thirty cases, while Sir James M'Grigor found it in India altered sixteen times in twenty-two cases, and in Egypt, as in India, he found it diseased."

The hepatic complication can in some cases be perceived from the commencement of the dysentery, while in others it probably arises in the course of the principal disease. The daily exploration of the hepatic region, and of the course of the large intestine, will therefore be necessary in every case, in order to the just appreciation of all its attendant circumstances. It sometimes happens that enlargement of the liver is not manifested till after the disappearance of the dysentery; and we observe also that hepatic abscess is discoverable only after the cessation of the primary bowel disease.

The association of hepatic diseases with dysentery would seem to be most frequent in the climate of the East Indies, and in such as have a common influence. Dr. Morehead, of the Bombay army, found that out of thirty fatal cases of dysentery, twelve were attended with hepatic abscess; and out of twenty-five dissections of dysenteric patients at Moulmein, in Burmah, Dr. Parkes found hepatic abscess in seven cases, while the same proportion of com-

plication was observed in Ceylon by Henry Marshall.

"As functional disease," says Dr. Parkes, "implies a proportionate certain amount of change in the molecular structure of organs, either primary in the organ itself, or secondary to changes in the blood or vessels, the most delicate test of the condition of an organ will in time be the chemical constitution of its secretion, when this knowledge is obtainable. We already see this test beginning to apply itself in the case of the kidneys and urine. Judging from this test, the liver is found to be more or less dis-

eased in every case of dysentery."

In the province of Oran, Algeria, the French surgeons state that the hepatitis and consequent abscess were frequently coincident with dysentery. Of 157 deaths in two-thirds of which diarrhea and dysentery had prevailed during life, "there were in sixty-five cases notable marks of diseases of the liver, including twenty examples of abscess." The tendency to hepatic complication was found in Algeria to increase with age, and with the length of service in that country. The coincidence of disease in the cocum and liver was noticed by me in the wards of the General Hospital of Calcutta; and, reporting on the diseases of Bengal in 1839, I find the following statement:—"I have seen many cases in which morbid action was co-existent in the liver and cocum, and I would beg to call attention to the subject." This official record referred to cases which terminated in recovery; and the "calling

attention to the subject" had reference to the verification of the fact in dissection.\*

Severe illness obliged me to quit Bengal before this opportunity had been afforded me; but the fact of the morbid coincidence in question has since been established both at home and abroad. It is now established that disease is frequently "co-existent in the liver and cœcum"-that an intimate connection exists between ulceration of the mucous membrane of this portion of the intestine and hepatic abscess, owing to suppurative phlebitis, to purulent absorption, and to contamination of the portal blood, in chronic cases. In diseases so complicated as dysentery and hepatitis it is often difficult, with uncertain histories, to determine with accuracy where morbid action begins; but the extension and coincidence of serious evil, in all the associated organs, is early discernible to the careful observer of symptoms in both diseases during life, and we can readily trace the results after death. But here, as in many other instances, we arrive at an ultimate fact, and can proceed no further with certainty:-we can see and examine the co-relation, but we do not, therefore, understand which is the cause, and which the effect. We know not, in too many instances. which of the diseases is the primary—that in the bowel, or the liver. But there are cases, and they are numerous, in which we can plainly discern hepatic disease and hepatic abscess as primary, concomitant, and secondary affections in the time of their occurrence; but whether they are to be regarded as a cause, a concurrence, and a consequence of dysentery, it is not so easy to determine; but, practically, a prompt and accurate diagnosis is here of much importance; for, by a speedy removal of hepatic disease, in the first and second instances cited, we may obviate suppuration of the liver while we cure the dysentery; and by the early healing of ulceration of the bowel, in the third instance cited, we may cure the dysentery and perhaps prevent the occurrence of hepatic abscess, or purulent deposit of any kind. I would direct the special attention of pathologists to the state of the lining membrane of the ducts and gall-bladder, as affording, perhaps, some clue to the pathology of hepatic abscess in association with dysentery.

In a review of this subject, published in the "Lancet," the

writer draws the following conclusions :-

1. "That dysentery, in a great number of cases, commences and runs its course uncomplicated by hepatic disease.

<sup>\*</sup> It is worthy of remark that in the only dysenteric case of a Hindu, in which "the cocum with its appendages was totally destroyed by a sloughing ulcer perforating in some places all its coats—similar ulcers being formed in the colon and its ascending portion, and sigmoid flexure, as well as in the rectum," there was found, by Dr. Parkes, in the liver, "throughout its substance, numerous small abscesses filled with viscid pus."

2. That hepatic disease may in some cases be the predisposing or exciting cause of dysentery.

3. That a large majority of the fatal cases of dysentery are

complicated with hepatic abscess.

4. That in a much larger majority of these cases, ulceration of the intestine is the primary disease and the source of the hepatic abscess."

Diarrhœa, easy of cure, occasionally results from vitiation or redundancy of bile; but not so dysentery, as suggested by Annesley. It is an every-day occurrence, on the other hand, to find that diseases of the liver, and consequent suppression of the biliary

function, will cause both diarrhœa and dysentery.

The whole question, however, is still an open one. "The connexion between abscess of the liver," says Dr. Ballard, "and dysentery, as a clinical fact is indisputable. Dr. Henoch, however, refuses his assent to the explanation of it, supported by Budd, who views it as a result of phlebitis, the pus from which, carried by the portal blood, is arrested in the liver, and becomes the focus for the formation of an abscess. Apart from the debated question of the capability of the inner membrane of veins undergoing inflammation at all, Dr. Henoch considers, as opposed to this explanation, the fact that Dr. Parkes, on the most careful examination of such cases, never found the slightest trace of inflammation in the small veins of the intestine, while no direct proof has been advanced of the mediation of the portal blood in the process." Dr. Henoch concludes:- "I believe we must give the preference to that view which regards the two diseased processes, dysentery and abscess of the liver, as without mutual relation, but as running their course together, dependent upon one and the same cause; in favour of which view is the circumstance, that in hot climates abscess of the liver also very frequently occurs associated with remittent fevers, or consecutive to them, without dissection exhibiting any ulceration of the mucous membrane of the intestine."—British and Foreign M. C. Review, July, 1854.

When dysentery follows upon, or is associated with intermittent fever, the spleen will frequently become enlarged. We then perceive a general anæmia, or splenic cachexia, with a low asthenic type of dysentery. This complication augments the difficulties and embarrassments in treatment, and greatly increases the dangers of the original disease. In temperate climates dysentery is found associated occasionally with continued fevers, and then it

becomes contagious.

The scorbutic complication may be expected to arise in seamen and soldiers, when, in tropical climates especially, they have been supplied for some length of time with unwholesome rations, or with such as consist in whole, or in chief part, of salted meat, Sir Gilbert Blane asserts that the disease "has been known to arise among prisoners of war living entirely on fresh diet, and solely imputable, therefore, to confinement in bad air, a dull uniformity of life, depression of spirits, and indolent habits naturally belonging to a state of captivity." The most terrible instance of suffering from this cause was that of the European portion of the force employed in Ava, during the first Burmese war, where they were for six and a half months fed on salt rations, and where 48 per cent. perished within ten months, principally by scorbutic dysentery.

The symptoms noted by all the medical officers were as follow:
—swollen, loose, and livid gums, with ulcerated and sloughing edges, and fetid breath, pain and hardness in the calves of the legs, constriction of the hams, and purple discoloration of the skin of the lower extremities—edematous swelling of the feet and legs—anarsarca—ascites and hydrothorax. When to such a condition of the system, scorbutic dysentery was added, the speedy and fatal character of the disease may be inferred.

The bowel complaint was characterized by green, or greenish-yellow discharges, becoming by degrees sanious; then dark, bloody, grumous, and putrid. A sudden and universal dropsy marked the latter days of the sufferers: I saw one of Her Majesty's regiments at Prome, in the second campaign, lose forty men in one night from this cause; and at Rangoon, in the first year, the Madras European regiment, containing many old, hard-drinking soldiers, was the first corps to be destroyed by this dropsy which, from its suddenness and fatality, resembled the worst form of the Beribery of Ceylon.

From such a hopeless state of disease as has here been described, there are many grades, down to the very minor degrees; and the medical officer who finds his men fed on bad, poor, and insufficient diet, under tropical influences, should be on the watch for scorbutic taints. They are all dangerous, and they mount rapidly from the least to the most aggravated grades.

PATHOLOGY.—As the symptoms of tropical dysentery during life will depend on the condition of the general habit—on the amount of plethora, of congestion, or of actual inflammation in a portion, or in the whole of the abdominal viscera—so, the results of these morbid actions will, after death, vary with the form and degree of the disease. Generally speaking, tropical dysentery consists in inflammation, more or less extensive, along the mucous membrane, and occasionally in the solitary glands of a portion or of the whole of the large intestine, running rapidly to ulceration more or less advanced, unless checked by treatment, and always associated with disturbance of function of the skin and liver. "Dr. Parkes considers it to be a process of ulceration universally com-

mencing in the solitary glands of the large intestine; others, with Mr. Raleigh, consider it to be a simple inflammation of the mucous coat of the large intestine; but if it were simple it would be more amenable to treatment. Whatever of truth or error may be in these opinions, the appearance presented to us in simple Bengal dysentery, is that of an inflammation of the large intestine, which may be diffusive, ulcerative, purulent, hæmorrhagic, or gangrenous, according to circumstances!" . . . "It should be borne in mind that the state of the solitary glands, as observed by Murray and Parkes, exactly corresponds with their usual appearance in cholera, and that all Murray's and most of Parke's cases, occurred in dysenteric patients suddenly carried off by that disease.—Dr. Macpherson.

In the milder cases which yield to treatment within two or three days, we may fairly presume that diseased action has not proceeded beyond plethora, or the mildest form of vascular congestion, or of inflammation of the mucous surface and glands.

In severe cases, the inflammatory action far from confining itself to the mucous membrane, extends to the serous covering, and even to the solid viscera, while ulceration, and even sloughing of the mucous surface and its glands, may be in extensive progress; and all this, occasionally, without that amount of corresponding fever which might, a priori, be anticipated—a circumstance which gives a character of peculiar insidiousness to a disease already very dangerous in its nature. Abrasions, ulcerations, and sloughs are but varying degrees of the same morbid process, the sloughing sometimes involving all the structures alike. Again, ulceration is found to result from deposits of pus, lymph, and serum below the mucous membrane, as well as from inflammation of this tissue.

"Omental inflammation or congestion," says Robert Jackson, "accompanies dysenteric fever not unfrequently. The indications of its existence are obscure; and its foundations are not moved without difficulty—not without the most extreme measures in bleed-

ing, blistering, &c."

When much tumidness with tenderness of the abdomen is associated with irritability of stomach and vomiting; inflammation of the peritoneal covering may be more than suspected, and postmortem appearances will frequently confirm this view of the case. When again, from want of treatment, or through unskilful treatment, the disease has proceeded unchecked for a considerable time, the least that can generally happen will be the merging of acute into chronic dysentery—or imperfect healing of the ulcerated howel

Certain portions of the large intestine have always appeared to me to be more affected when the complication in dysentery has been of the hepatic nature—namely, the cocum and the rectum. In the General Hospital of Calcutta, a government institution for the treatment of Europeans, I observed so frequently as to attract my special notice, that, when the cocum was the principal seat of pain and swelling, the liver was generally enlarged; but whether as cause or effect I could not say. As with the natives of India, Hindu, and Mahomedan, so with British born subjects in their native climate, hepatic abscess is found to be but very rarely associated with dysentery; whereas with Europeans placed out of their native climate, and under great heats, as in the East Indies, hepatic diseases and abscess have been found to be very frequent complications in dysentery.

In the Millbank Prison, out of "many hundreds of cases," which occurred during seven years, "not one has been complicated with hepatic abscess," says Dr. Baly. These facts would lead to the conclusion that a foreign climate mainly, and its unnatural influences, and more particularly heat, produces the difference of

result, as respects British subjects.

The other seat of irritation and of aggravated disease, associated with, or derived from, the hepatic complication, I observed to be the rectum; and in this portion of the bowel two causes appeared to me to operate unfavourably: the first consisted in the arrest of circulation in the hæmorrhoidal veins consequent on congestion or enlargement of the liver, producing internal or external hæmorrhoids, or both. Through inflammation, thickening, and adhesion, the cocum speedily loses its tone and contractile power, thus becoming the passive receptacle for the lodgment of vitiated and decaying matters; while the angle of the bowel at its sigmoid flexure becomes, through the incessant drainings of the same matters over its surface, the special seat of soreness. Much of the suffering and distress of the patient may always be traced to the irritated states of the sigmoid flexure and rectum, kept in constant tormina and tenesmus by the passage of such matters, often with the addition of acrid scalding bile. The same distresses I have seen produced by the misapplication, or by the protracted use, of calomel and drastic purgatives—by over-treatment, in short.

Death being presumed to occur only in the severer forms of dysentery, and in the worst cases only, we perceive on removing the abdominal parietes a turgid and congested state of the vessels, with adhesions of the serous coverings of the omentum, mesocolon, and mesentery, with varied and more extended adhesions between the different viscera; enlargements and inflammations of the glands of the mesocolon and mesentery, with occasional suppurations in them; inflammation and its results, such as thickening of the coats of the large intestines, softening, ulceration, or sloughing of the mucous membrane of the ileum, or of a portion, or of the entire of the larger bowel—the ulcers having the appearance of

variolous pustules, and being, in fact, glandular ulcerations, sometimes perforating the coats of the intestines into the general cavity; thickening of the coats, and contractions of the bore of the intestines, from effusion of fibrine and lymph, especially of the larger portion, and most frequently at its sigmoid flexure. Adhesions of the omentum to the cœcum, and abscess in the vicinity of the cœcum,

are occasionally observed.

In the hepatic complication the liver will be found congested, inflamed, suppurated, or indurated, according to the degree and duration of the disease. In the dysentery of the Peninsular army, under the Duke of Wellington, the spleen, liver, and mesentery were generally found diseased; and in the epidemic form of the disease, as it prevailed in Ireland, the liver was diseased in half, the spleen in a fourth, the small intestines in two-thirds, and the colon and rectum in all the dissections; but I do not recollect mention of hepatic abscess from phlebitis, purulent deposit, or

other cause, an event so frequent in the East.

In the scorbutic complication the post obit appearances are altogether of an asthenic character, indicative of a diseased condition of all the fluids. The inner surface of the large intestine, and of the ileum likewise, generally exhibiting extensive disorganization and decay, the bowel being in many cases filled with shreds and sloughs of mucous membrane and grumous blood, while the liver has a blue appearance externally, and when cut into the blood which flows is atrabilious and dissolved; but where hæmorrhage from the bowel has occurred, the liver will be found softened and anæmic. The spleen crumbles under the slightest pressure, like a mass of grumous blood. Ecchymosed patches are to be seen on the outer skin, on the external and internal surface of the bowels, while congestions of diseased blood and softening of textures are found in the heart and lungs. Appearances such as these indicate-either that the forces of diseased action have been invincible in their nature, or that the treatment intended for their removal has been inert, or insufficient.

TREATMENT OF ACUTE TROPICAL DYSENTERY.—
The first and greatest care of the medical officers of fleets and armies should be directed to the *prevention* of this formidable disease, by securing such attention from the authorities to matters of diet, clothing, protection from weather, and the conduct of duty, as may preserve the men in the vigour of health. There is no disease in which an early preventive and remedial attention is of more importance than in dysentery—the very great mortality of the disease in many of our tropical possessions being in considerable degree referable to the neglect of early measures of prevention and cure. Soldiers and seamen, when left to their own ways, will go

on with their duties under seriously advancing disease; and it only when "bloody stools, with cutting pains," afflict them, th they will "report sick." It is here that the all-powerful influence of commanding officers should be solicited by the watchful su geon. An inspection-muster, morning and evening, should I ordered, with a view to be assured of the earliest detection of according to the earliest detection of the ea vancing disease. This simple measure will save vast suffering

and many lives.

But however watchful the sanitary precautions, a certain amoun of sickness from dysentery will always occur in naval and militar service within the tropics. It therefore behoves all who have treat this disease to bear in mind that the destruction of thos placed in their charge is at hand-indeed imminent-unless promptly relieved by appropriate medical means. Every consider ration of humanity and of duty to the State should concur an unite in directing our efforts to the speedy and effectual arrest the sufferings and dangers attendant on tropical dysentery. Fo tunately we have, both in the army and navy to treat men in th vigour of life, and the further advantage that the patient come under our care immediately that he falls sick. A speedy an secure convalescence therefore results from a timely, well-regulated and decisive course of treatment, while chronic dysentery, wit lingering sufferings in hospital, or permanently broken health, wi follow on feeble and indecisive attempts at cure. Let it be remen bered also that cases, bearing the appearance of the mildest cha racter of disease are occasionally found to exhibit in course of few hours the real characters of destructive inflammation. Nothing therefore, but the most watchful care, guided by intelligence the selection and use of means, can lead to the patient's safety.

It will be found with dysentery, as with fever, that a sufficien abstraction of blood by venesection, practised at the very onset the disease, will simplify and render easy all the subsequent stage of the cure; so much so that, in mild and uncomplicated dyser tery, as in the violently acute and inflammatory type of the dis ease, blood-letting is the primary as well as the cardinal measure In acute uncomplicated dysentery, a moderate bleeding from th arm, followed by a full dose of calomel and James's powder, that of ipecacuanha, say twenty grains of the latter to ten of calc mel, given at bed-time, a warm bath, or hot fomentations to th abdomen-a morning aperient, followed during the day by sudo rifics conjoined with diuretics—the moderate use of demulcer drinks, allowing no other sustenance-will, in a very few days bring about convalescence.

Where pain remains in any particular region of the abdomer as that of the cœcum, sigmoid flexure of the colon or the rectum leeches should follow the general abstraction of blood, guardin carefully against subsequent oozing from the leech-bites, and using warm fomentations or poultices only where the bites have been thoroughly dried up. While any irritation of the bowels remain, as indicated by frequent stools, containing mucus and blood, the nightly dose of ipecacuanha, diminished in amount as the symptoms mitigate, must be continued, followed during the day by cooling diaphoretics with diuretics, as recommended in the treatment of fever. In the use or the withdrawal of calomel or blue pill at night, along with the sudorifics, we should be guided by the presence or absence of fever with a furred tongue, and by the healthy or depraved condition of the intestinal discharges. Mercury ought not, in short, to be regarded, in every case, as a necessity in the treatment of dysentery. It is not a remedy of necessity in all cases or in all climates: in the West Indies, for instance, the hepatic complication is by no means so frequent as in the East.

While the use of sudorifies, with some mild mercurial, is indicated by the continuance of dysenteric symptoms, with more or less vitiation of the secretions, a mild aperient in the form of castor oil, or of compound powder of jalap, should be exhibited every other morning, while in this mild, as in every form of acute dysentery, the diet must be of the very lowest, presuming that the time

required for the cure will not comprehend many days.

In the highly inflammatory type of dysentery, on the other hand, we shall find that, after free blood-lettings, the occasion for the use of calomel with the sudorific will depend on the constitution of the patient; and when he is youthful and robust, with fever, tumid abdomen, and furred tongue, the repetition of bloodletting, and that of calomel and sudorifies in full doses, will be called for; and the mercurial sudorific must be repeated every night, reduced in dose as the disease yields, until the discharges from the bowels assume a natural appearance, when calomel may be discontinued, and blue pill or mercury and chalk substituted, if any such should still be necessary. The circumstance too that, in certain cases, much blood is voided by stool should not deter the surgeon from the use of the lancet, for, on the contrary, the early and free employment of depletion will greatly conduce to safety. Of blood-letting and mercury it must be observed also that, although they prove most powerful to good when used early in the disease, they will still be found necessary, in their just proportions, where the disease has even existed for some time, or where treatment has been neglected. Blood-letting was repeated by Robert Jackson-"until all that was expected from it had been attained." But all the circumstances of each individual must be so weighed, in reference to constitution, age, habits of life, and length of residence in India, as to guide the medical officer in the amount and measure of every means employed. It is impossible to fix an arbitrary rule or guage for every case beforehand, and it is better that a young practitioner should reflect seriously on all that relates to his patient, and act on principle according to his judgment, than that he should betake himself to a pattern such as is occasionally afforded in narratives of published cases. He must learn that, in our management of an ever-varying disease, we are unable to lay down any catholic rules of treatment which shall be suitable to every case. When we feel assured that inflammation has been subdued, a full opiate may be added to the night dose of sudorific, or the two may be combined in the form of compound powder of ipecacuanha; and thus, provided we have been so fortunate as to see the patient, immediately on being taken with the disease, a few days of treatment, such as is here described, will prove effective of cure. Used in this manner, opiates conduce powerfully to the cure; and their free use, in the manner here recommended, should never be neglected. Robert Jackson frequently ordered the Dover's powder, in doses of eight grains thrice a day. In persons of a less robust constitution, again, and even in such as are delicate of habit, in whom dysenteric purging, griping pains, and pain on exploration of the abdomen are present, repeated abstractions of blood generally, and by means of leeches, will be requisite; the measure of blood-letting, as well as the occasion for its repetition, being estimated and determined by the amount and degree of the disease. In every case dysentery must be speedily overcome by treatment, if we would save our patient.

"When the state is inflammation," says Robert Jackson, "and the termination congestion, suppuration, or gangrene, the physician must not hesitate in striking a balance between the chances of debility and the almost certainty of death. There is here no safety in half measures." It was a maxim of Hippocrates that "in desperate diseases, desperate remedies, energetically pursued,

are the most efficacious."

But while without blood-letting there is no safety for the sufferer so long as evidences of inflammation are in existence, no sensible person would, on the other hand, practise such an operation, or continue the use of powerful remedies, after the subsiding of symptoms has indicated the approach of convalescence; and here we limit ourselves to the use of ipecacuanha as a sudorific, cooling diaphoretics with diuretics, mild aperients, cold water enemata, with or without lead and opium, anodynes by the stomach and rectum, warm baths, demulcent drinks, and farinaceous diet, all which means must follow upon depletion, and their exhibition must be persisted in until we shall have obtained to convalescence, but no further.

When, after the removal of inflammation, soreness along any

portion of the large intestine exists, with a suspicion of ulceration of the mucous membrane, the use of washed sulphur alone, or in combination with the bitartrate of potash, will prove the best aperient given every other morning. This combination seems to have a detergent and healing influence, and even the sulphur uncombined acts as a most gentle and soothing relaxant. Enemata are here also most powerful to good, either in the emollient and anodyne form, given at bed-time, or as an astringent and refrigerant in the form of cold water, having eight or ten grains of the acetate of lead, with two of opium, in solution. This last form of injection with diminished proportions of lead and opium, may also be exhibited three times a day where great tenderness and irritation exist in the rectum. It is always desirable that the injections should be retained, and to this end they should not exceed three or four ounces, and the operation of injecting should be performed in the most gradual and gentle manner; and as to the temperature, whether cold or warm, we should be guided by the feelings and wishes of the patient, these being the true tests.

It is important to pay attention to such adjuvants, for tormina and tenesmus are very wearing and depressing to the nervous system, and they are aggravated during the night so as not only to prevent sleep but, by the frequency of the calls to stool, the patient is exposed to the additional risks of checked perspiration and chills. To guard against such evils it will always be proper, in the case here under consideration, to add the enemata to the night opiate, or Dover's powder. Suppositories of ointment of lead, with two grains of opium, will likewise be found calming of irritation in the rectum, and therefore very grateful to the patient.

On the subsidence of inflammatory action, when fulness, tenderness, or thickening of the bowel exists at any portion of its course, blisters are always beneficially employed. Robert Jackson states, where tenesmus is urgent, with pain and heat in the rectum, accompanied by offensive discharges, he found the following powder very efficacious:—

Finely	powdered	charcoal .		Эj.
"	>>	rhubarb.		gr.x.
	**	ipecacuanha		gr.v.

A drachm of charcoal powder in rice water he also states to be equally effective as an enema. "These," he says, "of all means known to me have had the most instantaneous good effect in this form of disease. When the subject has been prepared by previous evacuation, &c., two or three doses of the powder by the stomach generally effect a cure." In severe cases he gave this preparation every five or six hours, but he adds, with characteristic

patches of sloughing ulceration pervade the mucous membrane, which sometimes cause large perforations of the bowel.

In treating this hæmorrhagic form of the disease we must have instant recourse to the most powerful astringents in the largest doses, aiding their effects by the addition of opiates; and both must be frequently administered by the mouth and anus. Lead may be used more liberally and more safely than is generally supposed. Dr. Christison of Edinburgh has given six grains of the acetate of lead daily for several months, without any ill effects, while Mr. Webber of Norwich informs Mr. Squire that, "in severe cases of bronchitis he administers from three to ten grains of the same medicine in pill, with a mixture containing three drachm doses of distilled vinegar every four hours till symptoms subside, which frequently takes place after the third or fourth dose."

In hæmorrhagic dysentery the lead should be given every two hours, so as speedily to establish its influence, while to each dose from half to a grain of opium is added, using at the same time strong enemata of solution of acetate of lead with opium, thrown with sufficient force to enter the course of the colon; and this course of treatment must be persisted in till the discharge of blood shall have totally ceased. In a form of the disease characterised by a rapid destruction of the organs immediately involved, and by general collapse of the nervous and vascular systems, such a diet must be allowed as may nourish while it supports the sinking powers of the patient; and with these views soup should be added to the farinaceous articles, while even wine must be allowed, in order to guard against sinking. Convalescence will be greatly promoted by the exhibition of quinine with nitrie or sulphuric acid, with the addition of a few minims of laudanum.

Dr. Wilmot of Tunbridge Wells, in a severe epidemic dysentery of a scorbutic character, and attended by sloughing ulceration of the mucous membrane, while he exhibited by the stomach the decoction and compound tincture of bark with sulphuric acid, administered in the form of enema a drachm of creosote in twelve ounces of thin starch; and to this latter means, exhibited every night at bed-time, Dr. Wilmot ascribes very decided results, as a

stimulant, antiseptic, and styptic.

But whether we have to treat dysentery in its milder or most severe form, complications will meet us, either from the commencement of the disease or during its progress, and we must then be prepared to meet each contingency as it may arise;—and first, of the hepatic complication. The liver will be found involved, in some instances, from the very commencement of the diseases, while in others it palpably arises in the course of the dysentery. The daily exploration of the hepatic region, as well

as of the course of the large intestine, will therefore in every case be necessary, in order to the just appreciation of all the circumstances of each case. Here depletion, both general and local. must be carried to such extent as to remove congestion of the portal system, tumidness, and pain of the liver, while by the nightly use of calomel and ipecacuanha, followed by compound powder of jalap in the early morning, and diaphoretics with diuretics during the day, we relax the skin, emulge the liver, and restore the lost balance of circulation. The measures here indicated must be repeated so long as the evidence of hepatic complication is present, discontinuing the use of calomel and blood-letting on its disappearance. In this, as in every other instance, where leeches are used, I would urge on the young surgeon the necessity of applying so many leeches as may be necessary, and no more, arresting the oozing from their bites without delay, while no poultices or fomentations should be applied until all oozing shall have completely ceased.

After the removal of acute symptoms, should fulness or tenderness of the liver still continue, the occasional application of blisters should be made over the region; and to advance convalescence the nitro-muriatic acid, internally and externally exhibited, will prove eminently serviceable. It is necessary in the dysenteries of the east to have constant regard to the hepatic complication; for it may be that, by appropriate means, we can remove inflammation and consequent ulceration of the mucous membrane of the intestine, but if disordered function, congestion, or inflammation of the liver tending to abscess be going on, the cure will be postponed or prevented by any one of the circumstances named, or by the occurrence of dysenteric relapse, a con-

dition to be vigilantly guarded against.

In the cure as in the prevention of scorbutus we have mainly to look to the diet, cleanliness, ventilation, and cheerful occupation; while medically, and referring to the scorbutic complication in dysentery, our first indication of treatment is, by appropriate medicine and regulation of the diet, to restore the health of the

In addition to the liberal use of pure lime-juice, vegetables and fruits should be abundantly supplied, and potatoes especially, as being eminently anti-scorbutic. During long voyages some reduction in the meat portion of the ration might be advantage-ously made, granting a proportionate increase of vegetables and fruits; for it is to the want of these last, even more than to the sameness of the diet and the salted meat, that the production of the scorbutic taint is referred. Scorbutus is not only a serious disease in itself but its complication with fever and dysentery, and the greater liability of scorbutic persons to be affected with the

various tropical diseases, malarious and other, renders its preven-

tion and cure a question of great importance.\*

In scorbutic dysentery fresh lime juice with opiates simply, or in the form of Dover's powder, will prove the best medicine, while a liberal allowance of fruits and vegetables should be granted, along with so much fresh animal food as the stomach will bear. When irritability of stomach exists, the chlorate of potash, the carbonates of soda and potash in effervescence, lime juice being in excess, may be given with advantage; and when tormina and tenesmus occur, with bloody discharges from the bowels, anodynes and astringents combined, or lime juice and opium in the form of enemata, should be employed, while the infusion of cinchona with the nitro-muriatic acid and opium are given by the stomach. When sloughs are discharged by stool along with grumous blood, the scorbutic taint being manifest, the danger to life is imminent. For the removal of the oozing which takes place in scorbutic dysentery from the gums and mucous membrane of the bowels, the astringent and tonic properties of the sulphate of iron, largely administered in solution by the stomach and rectum, and used as a gargle, will be found eminently serviceable; with the same view the muriated tincture of iron is recommended by Annesley and others, but it is not intended that either means should supersede the use of the direct antiscorbutics.

The combination of iron and alum will likewise be found of great service, as will the Sand Rock chalybeate water of the Isle

of Wight.

In default of lime juice the medical officer must seek for substitutes in citric acid, and in all kinds of summer fruits. For more than a century fruits have been justly lauded by our army surgeons in the treatment of dysentery, and the physicians of Spain and Portugal, according to William Fergusson, have for ages employed the juice of lemons with good effect in the endemic and epidemic dysenteries of their respective countries. Dr. Cabbel, a talented young surgeon of the Peninsular army, used no other medicine in treating British soldiers; when severely attacked in person by the disease in Portugal, and though attended by Dr. Fergusson, he refused all remedies other than lemons and oranges, and he speedily recovered. "In Trinidad, too, when a dysentery of uncommon malignity appeared there, Dr. Lynch O'Connor found that lime juice, administered by the mouth and in lavement, was the only remedy that could be depended upon."

<sup>\*</sup> Sir Gilbert Blane states that "the first ample supply of lemon juice to ships of war was as long ago as 1757, to the squadron under Admiral Watson in the East Indies, at the suggestion of his surgeon, Mr. Ives, a gentleman highly educated, of great merit and modesty; and we do not learn that he courted public notice, far less that he solicited a reward in doing what he deemed to be merely his duty."

Pringle, more than a hundred years ago, used grapes freely in the treatment of dysentery, and there have been few since his time who may be regarded as better authorities. On the subject of the Bael fruit of Bengal, and on the use of fruits generally in the East Indies, I beg to refer the reader to what is stated respect-

ing them in the article on chronic diarrhœa.

Dr. Kinlock Kirk, of the Bengal army, speaking of the sickly condition of the 78th Highlanders in Sindh, in 1843-44, states that the soldiers suffered from "large livers and spleens, and were weakened by continued colliquative purging." In this state, and suffering from a low fever, "the men found from their own experience more comfort and more advantage from the use of limes than from any other substance, and many gave back a portion of their rations to receive in lieu its value in limes." This intelligent officer, when serving in the same country with Sepoys suffering in the same manner, made use of the Phyllanthus emblica, or Aoula of the natives, and which is held by them in high repute. The fruit is used in various ways, preserved in its dried state, in syrup, or pickled, and is to be found in all considerable towns throughout India. "An infusion of two ounces of aoùla in twenty of water was given to ten men three times a day, in conjunction with whatever else was necessary, as bark, quinine, opium, &c.; and I found this remedy as useful certainly as lime juice, in cases where the spleen and liver were enlarged." Dr. Kirk found on examination that the fruit consisted of "citric acid. tannic acid, and pectin, or vegetable jelly." For years Dr. Kirk has been using lime juice in the treatment of enlarged spleen, "with the most beneficial effect." This quotation is made from the "Pathologia Indica" of Professor Allan Webb, of the Calcutta Medical College, a work of the greatest value to all who are interested in the progress of tropical medicine.

In treating the complications of dysentery with fevers remittent and intermittent, we must have regard to the type and prevailing character of the febrile movement, as well as to the paroxysms, in addition to the attention that may be demanded by the condition of the bowels. When the fever is of an ardent inflammatory character our antiphlogistic means must be applied with an energy proportioned to the violence of the disease; while in the low adynamic type we must have recourse to bark, quinine, opiates, chalybeates, the vegetable and mineral acids, improved diet, and change of air. In convalescence from dysenteries, acute and chronic, much active benefit will be promoted by the external and internal use of the nitro-muriatic acid. When the spleen is found to be enlarged, whatever the stage of dysentery, and whatever the type of the fever, mercury must be carefully avoided, trusting rather to the means just named, and to all such as vivify and im-

prove the blood. It is almost unnecessary to add likewise that the abstraction of blood, in the complication of these fevers with dysentery, should be practised only when the fever is of the acute or ardent type, the adynamic forms, as well as the splenic cachexia, not admitting of the use either of mercury or of blood-

letting.

As regards the treatment of convalescence from dysentery, I would impress on the inexperienced a careful attention to all that has been urged upon that subject in the conclusion of the article on fever. There is not a word of caution there inculcated that does not apply with equal force to the management of the stage of recovery from tropical dysentery, as to that of convalescence from fever; and too much attention cannot be paid to either case if we would protect the seaman or soldier from the dangers of relapse. It should be a rule with all medical officers to grant the seaman and soldier sufficient time to confirm convalescence and restore health, before returning him to duty. Time is necessary to the absorption of interstitial deposits, and for the cure of functional disorders, the results of acute disease; and it would indeed be a mistaken notion of efficiency which should cause a half-cured man to be sent to labour whether in peace or war, not to speak of the inhumanity of such conduct.

So much having been offered on the treatment of dysentery, simple and complicated, I must once and again urge upon the naval and military surgeon the value of time in dealing with such a disease. In treating fever as well as dysentery it should again and again be remembered that time is either our most important ally, or else our most powerful enemy, just as we may use or neglect it. Time will often be turned into a dire enemy both by the soldier and seaman, if permitted; and in civil life the same neglect will occur through the carelessness or the vices of individuals, so that the patient comes under treatment at all stages of the morbid progress. Excepting under the pressure of actual war, when dislocations and impediments will occur, it is happily in the power of officers in command to order parades of inspection conducted by the surgeons; and thus every man may be looked to on the instant of his falling sick, an incalculable

advantage to the individual and to the state.

I propose now, as in the instance of remittent fever, to present a catalogue-raisonné of the treatment of Dysentery in different ages and countries. This, while it exhibits the limited views of some men, establishes with authority the comprehensive measures of cure of the majority of authors:—

1622. Bontius.—Bleeding, repeated according to occasion—vomit of ipecacuanha, and a purge—the extract of saffron, "the

anchor of hope"-emollient enemata and fomentations-fruit diet.

1696. Sydenham.—Early blood-letting—opiate at night—active purge in the morning, repeated every other day, and followed by a powerful opiate—an anodyne with a diaphoretic every night, or every eight hours in severe cases—emollient injections, and injections containing Venice turpentine.

1760. Dr. Huck.—Blood-letting, repeated according to occasion—purgatives—ipecacuanha and tartar-emetic, in repeated and

full doses.

1768. Pringle.—Blood-letting—vomiting—mercurial purges—

ipecacuanha and opium-grapes.

1773. Dr. John Clark.—Full emetic of ipecacuanha and tartaremetic, followed by mild purgatives—ipecacuanha and opium in small and repeated doses.

1782. Mr. Curtis.-Purgatives chiefly; in the advanced stage.

small doses of ipecacuanha powder-astringents and tonics.

1787-9. Dr. Moseley.—Blood-letting—antimonials freely administered—revulsion.

1791. Mr. Wade, of Chunar.-Solution of tartar-emetic and

purging salts-anodynes and sudorifies.

1791—1820. Robert Jackson.—Copious blood-letting during immersion in the warm-bath, "the sovereign remedy, without which nothing can be done with a fair prospect of benefit," followed by an emetic and mercurial purgative;—the alternate use of evacuants and sudorifics, "Dover's and James's powders holding the first rank of the latter"—hot fomentations to the abdomen, with the local use of cold water, and "tonic applications in glyster"—warm clothing—change of air, especially a sea voyage.

1796. Dr. Hunter.—Mercurial and saline purgatives, followed by opiates. In mild cases this constituted the sole treatment:—in severe cases—draughts of infusion of bark with rhubarb—fo-

mentations-blisters.

1799. Dr. Lempriere.—An emetic—warm baths and fomentations—emollient injections "as a necessary preparation to a purge"—active and persistent purging by common purging salts, by calomel and rhubarb, or by calomel and jalap—diluents—farinaceous diet.

1799. Blane.—Blood-letting—vomit and purge at the beginning—ipecacuanha and opium—purgative salts—small doses of

ipecacuanha powder.

1799. Dr. Whyte.—Profuse blood-letting—flannel roller over the abdomen—careful confinement to bed, the body being anointed with oil—no medicine.

1804. Sir James M'Grigor.—Blood-letting—mercury with sudorifics—flannel roller round the abdomen—aperients—opiates—nitric acid internally, and in the form of bath—enemata of lead and zinc—demulcent drinks—antiphlogistic regimen—change of air.

1814. James Johnson.—Blood-letting, repeated according to occasion—calomel in full doses—sudorifics—occasional mild purgatives—anodynes—cooling antiphlogistic diet—warm clothing—

change of air.

1817. William Fergusson.—Blood-letting—small doses of calomel with ipecacuanha—inunction, till the gums become affected—fomentations—warm clothing—the use of fruits—change of air.

1818. Sir George Ballingall.—Topical bleeding—purgatives—infusion of ipecacuanha—opiates—warm baths and fomentations—enemata—blisters.

1818-22. The Dublin Physicians—Epidemic dysentery.— Blood-letting general and local—calomel, antimony and opium combined—emetics—enemata—counter-irritation—warm baths.

1819. Mr. Bampfield.—Blood-letting—cathartics—diaphoretics

with mercury.

1822. Annesley.—Emetic of ipecacuanha, followed by full doses of calomel, smart purge, and warm bath—then blood-letting general and local, according to constitution and length of residence in India—calomel and opium alternating with purgatives and enemata—ipecacuanha, or antimony, with opium as a sudorific.

1823.—Dr. Latham.—After the failure of all the remedies common to European practice, including ipecacuanha—" calomel and

opium became the settled practice."

1832. Mr. Twining.—General and local blood-letting, repeated according to occasion—simple powder of ipecacuanha with extract of gentian—purgatives.

1833. Dr. J. Smith, Edinburgh—Epidemic dysentery.—Scruple doses of calomel, given so as to fall short of salivation, "the

more common measures having failed."

1833. Dr. Joseph Brown—Cyclopædia of Practical Medicine.—Blood-letting general and local, repeated according to urgency, aided by hot baths and fomentations—gentle laxatives—mercury as a subsidiary to general and local bleeding, and combined with simple or compound ipecacuanha powder—sudorifics—opiates—enemata.

1835. Copland—Dictionary of Practical Medicine.—Blood-letting general and local, repeated according to occasion—opiates "after depletion"—mild aperients—cooling diaphoretics—ipecacuanha and opium—blisters.

It is observed by Mason Good, that "the principles of Syden-

ham's practice it is not easy to improve upon;" nor, indeed, have either his principles or his details of practice been much improved upon, from his day to the present—blood-letting, sudorifies, opiates, and aperients remaining still at the head of our standard remedies.

There can be no doubt that the "putrid dysentery" of Clark, described by him as prevalent in Calcutta in 1773, was complicated, like the fevers of those days, with scurvy. The then protracted voyages to India, the bad diet, the general sanitary neglect during the voyage, and, if possible, the greater sanitary neglects after landing in Bengal, left little hope to the British seaman or soldier when tropical disease was added to scorbutus.

It is stated, on the authority of the records of the Calcutta European General Hospital, that even in more recent times the mortality by dysentery was as follows:—

	Treated.		Died.
1797. March, April, and May	. 37		. 21
" June and July			. 35
" August and September			
" November and December	. 21		. 15
1798. March, April, and May			
" June, July, and August			
" October, November, and December .			
1799. July, August, September, and October	. 31		. 11
	-	-	-
Total	. 238	3 .	. 127

In those days the treatment of fever and dysentery was by calomel and opium, mercurial inunctions, bark, animal food, and wine.

Of the authors quoted in the above catalogue, those who speak most in favour of their success in treating tropical dysentery are—Bontius, Dr. John Clark, Dr. Moseley, and Mr. Wade:—a brief inquiry into the merits of their respective modes of treatment may therefore prove of some interest.

Bontius commenced with the abstraction of blood, and with vomits;—then came the extract of saffron, "than which, I dare to say, a more excellent remedy was never discovered by mankind; and I am fully persuaded that it is the most perfect antidote against this disease, even when of a malignant kind."

Dr. Clark merely wishes "to present facts; but he must add that he has given bark to one hundred and fifty fever patients in Bengal and other places in the East Indies; and of that number lost only one who took the medicine with perseverance." This gentleman was equally successful in the treatment of dysentery;

for out of "a number of patients" he only "lost four."

Dr. Moselev abstracted blood likewise: - "Bleeding," he says, "being an operation of great consequence in the flux, the cure is generally begun with it, repeating it as symptoms authorise, observing only, "Non que ætas sit sed quæ vires sint." After bleeding, a vomit of ipecacuanha is to be given, and then an opiate after its operation is necessary. This is to be followed by a careful and continued use of antimonial wine and laudanum combined, with a view to keep up a sweat proportioned to the violence of the disease, and not in the trifling way of giving them in small doses, whilst the patient is exposed, and their operation neglected." The doctor goes on to relate that the eminent success of this plan was exhibited when the soldier had been suffering from "the worst condition of the disease, with blood running from him as in a hæmorrhage;" and he adds that this demonstration of his superior success had been made before "several of the officers of different regiments in the West Indies who were desirous to be spectators of a fact so interesting to the army." He concludes his triumphant narrative by exclaiming-" Such is the power of revulsion."

The power of revulsion in the cure of dysentery is undoubtedly great, whether produced by agents internally or externally exhibited. I remember, when serving with the army in Ava, to have heard that a British merchant, while prisoner in the capital, was seized with dysentery. After days of unmitigated suffering, under circumstances of barbarous treatment, and when death appeared to him near and certain, an order arrived for the removal of the prisoners to a place of closer and still more severe confinement, where they were literally packed together. Here the sufferer was thrown into a violent perspiration; and from that time all his symptoms left him. Such, again, is the power of revulsion!

Mr. Wade, of Chunar, claims an amount of success—unbroken except by "two cases of unfortunate termination in the treatment of about four hundred cases of fever and dysentery." The treatment for both diseases was by a solution of purgative salts and tartar emetic, followed by anodynes and sudorific night

lranghts.

Mr. Wade observes, that "the medical world may draw their own comments on the cases which are submitted to their examination. The person who has treated and compiled them shall defer his, until the public may have formed some unbiassed opinion of them, he should only venture at present to vouch for their authenticity." This is just what has been done and said at all times by every writer of such cases, and by every pretender to extraordinary cures;—but it is a curious fact, well deserving con-

sideration how it happens—and happen it always does—that those who in their day claim the greatest and most exclusive success in the practice of medicine should, in after times, be the least followed in their modes of treatment.

Some of the causes of error in reasoning and practice, as respects dysentery, are so well stated by Sir James M'Grigor that I cannot do better than present them in his own words:- "My opportunities of seeing this disease have been no common ones. Rarely, I believe, has it fallen to the lot of an individual to see so very many cases of one disease in such a variety of climate and situation. In the 88th Regiment, during more than ten years, I saw the same man the subject of this disease on the continent of Europe, in America, in both extremities of Africa, and in India. Of late, it has afforded me not a little amusement to review my notes as well as my journals of practice in this disease, in all these quarters. . . . . I became convinced in Alexandria that, with change of country and climate, we had a different disease. This is one proof how improper, and how unsafe, it is for the practitioner of one climate to set down and describe the diseases of another. They only who have studied the same disease, in various opposite climates, can fully comprehend the extreme absurdity as well as the fallacy of this. From reasoning of any kind, we are incompetent to decide on the identity of disease. Reasoning from analogy here always deceives. In many of the symptoms diseases may agree; but from thence to infer their identity is taking a very narrow view."

Dr. Macpherson, of Calcutta, speaking of the comparative results of the treatment of dysentery by the free use of opium, and other soothing means, observes with proper candour:—" Our obvious difficulty in arriving at safe conclusions is, that the disease itself varies so much in intensity in different years, as well as in different periods of the same year. Thus her Majesty's 55th Regiment lost at Secunderabad in 1837, one in four and a half; in 1838, one in seven and a half; in 1839, one in ten; yet the treatment was the same, and conducted by the same medical officer throughout."

If dysentery and fever were diseases of an uniform character, having an uniform cause and seat, then perhaps they might be treated after an uniform plan; but a very slender experience of the two diseases, as they prevail within the tropics especially, and even within the British Isles, shows that this is not the case; for although, in the disease now under consideration, some portion of the larger bowel is universally implicated, yet, either from the accession, or during the progress of the disease, for we cannot often say which, the lesser bowel, the liver, the spleen, the pancreas, and mesentery, frequently become the seats of plethora.

and of other morbid actions, so as greatly to complicate the

disease, and modify its treatment.

The pathology of our dysenteries, whether in southern or northern India, as described by our best authors, sufficiently establishes the fact that, in this formidable disease, morbid action is not confined here, any more than in Europe, to the course of the large intestine alone, but that all, or most of the associated organs are found after death to be more or less deeply involved, just in proportion to the extent and severity of the symptoms during life. The truth is, that the dangerous and fatal characters of tropical dysenteries are principally due to these complications. "I have been repeatedly astonished," says Dr. Macpherson of the General Hospital, Calcutta, "to discover after death an immense extent of structural change in boys, whose illness could not be ascertained to have exceeded eight or ten days." It appears to me that to a want of just consideration of these inevitable pathological complications must we refer the system of exclusive treatment, so much reprehended by every author who has had varied and extended opportunities of observation in dysentery; these complications explain also the successive abandonment, by the surgeons of our fleets and armies, of every exclusive plan of treatment hitherto proposed, almost as soon as it has been tried. "In the treatment of no other disease, perhaps," says Dr. Copland, "has the baneful influence of exclusive medical doctrine been more fully exerted than in that of Dysentery."

The adhesions of the peritoneum—the pink colour, thickening and softening of the omentum—the effused lymph—the adhesions between the different intestinal surfaces—the red and thickened state of the mesentery, with the enlargement of its glands—while they indicate the general amount and great extent of inflammatory action during life, leave the fact still demonstrated that, in causing death, the main force of the disease has been expended

on the mucous surfaces.

He who would treat dysentery with success, while he shuns exclusive means, must assign to each remedy its proper value, and neither more nor less. Blood-letting, sudorifics, and purgatives constitute the most universal remedies, and in simple uncomplicated dysenteries they will prove all-sufficient. But when the abdomen is tumid, and there is pain in the liver, or in any other region, while the nature of the discharges indicates advancing inflammation, calomel conjoined with sudorifics, and repeated to meet the occasion, will powerfully aid the curative effect, through its influence on the depurative functions—on the circulation, in unloading, jointly with purgatives, the gorged vessels of the abdominal organs—on the blood and on secretions generally—and on the very sudorific function which we so much desire

to excite. This, in so many words, I believe to be the real value of calomel in the treatment of dysentery; and the inexperienced have been led into much unnecessary doubt, and into no small amount of error, by some writers who regard calomel as a remedy of necessity, and by those who, on the other hand, would exclude its use altogether.

It is hardly necessary to repeat that while calomel is a most powerful agent in its place, as an aid to blood-letting, the pushing it to the extent of ptyalism is here by no means intended; nor should mercury in any shape be used in adynamic forms of the disease, in the splenic cachexia, nor in states of anæmia—for in all these conditions of the system its actions are

most injurious.

Amongst the surgeons in Bengal, blood-letting general and local, takes the lead, and has done so for many years in the treatment of dysentery. It is the standard remedy; and I believe that when the subject comes early and freely under this treatment, and that the case is not complicated with hepatic or other congestion, or with inflammation, little else than a few doses of sudorifics and aperients, as aids to general blood-letting, will be needed for the cure. But as in most cases of this formidable disease, as it appears within the tropics, the morbid action going on in the large intestine is intimately associated with general abdominal plethora, and individual complications, other and important means speedily follow upon the bleeding; and of these the first are those which act powerfully on all the secreting organs, internal and external such as calomel with ipecacuanha, in full doses at first, and given at bed-time—followed by mild laxatives—sudorifics—enemata warm baths and fomentations-blisters, and other minor adjuvantia. I believe this to have been for many years the course of practice in Bengal, and I have seldom seen calomel urged to the extent of producing salivation; neither do I think this degree of effect at all necessary to the cure. Several experienced medical officers have assured me that antimony in combination with the calomel proved, in their hands, quite as efficacious as ipecacuanha; but in my own practice I always preferred the latter.

When, after a few days, inflammatory action has been subdued by blood-letting, with the lowest diet, and intestinal irritation has been overcome by calomel, ipecacuanha, and aperients; then, recourse should be had to milder measures, as a persistence in the stronger ones is no longer either necessary or proper to the cure; indeed, such persistence would prove eminently injurious.

The late Mr. Twining would appear to ignore the associated diseases of the liver, and, in his clinical work, he advocates the use of simple ipecacuanha powder combined with bitter extract, which plan of treating dysentery he states to have been very suc-

cessful. As first assistant, and as officiating surgeon to the General Hospital of Calcutta, during Mr. Twining's service in that institution, I had ample opportunity of witnessing the results of his treatment, and I am not aware that his system was followed at the time, or since, by any of the other medical officers of that hospital; neither would his plan of treatment appear to be successfully imitated in the provinces throughout Bengal. Dr. Macnab, in an excellent practical Report on the dysentery of the native soldiery of Hindustan, when serving in Bengal Proper, says, that "blue pill with ipecacuanha and extract of gentian proved a complete failure, as has generally been the case in my trials of it. Indeed I much suspect that Mr. Twining overrated the value of his favourite remedy, and that he may have also miscalculated the anti-emetic properties of the gentian."

On Mr. Twining's use of drastic purging, Dr. Everard says:—
"Twining has certainly overpraised repeated strong purging.
The Deputy Inspector General, Murray, informed me that in his tour of inspection, he found some cases which had been dysenteric, labouring under one-drachm doses of compound jalap powder given daily at noon, which got well immediately on removing the

cause."

Ipecacuanha has been a favourite remedy in the south of India for nearly a century, although Mr. Twining, like many respectable practitioners, fell into the error of supposing that what he observed was new, while in truth it was very old. Speaking of his cotemporaries, Mr. Geddes, Drs. Mortimer and Baikie, of Madras, he concludes:- "In fact, ipecacuanha has been a favourite remedy with some of the medical men at the Madras Presidency;" and from the context it is evident that a very recent date is here meant. Dr. Whitelaw Ainslie, writing in 1813, after an experience of thirty years in Madras, and an extensive practice amongst all classes of Europeans, says of this drug that "it has no equal in simple dysentery, that is, dysentery not complicated with hepatic derangement; in such cases, given even so as to produce daily a little vomiting, it has the happiest effects." This is an observation of great truth and practical importance consequently, and I believe that it impresses and fixes a just discrimination in the use of ipecacuanha. In speaking of a personal experience of two-andtwenty years in the East Indies, and of an extensive range of observation of this disease as it occurred in hospitals, European and Native, and in extensive private practice in Calcutta; also as it occurred amongst troops serving in the unhealthy provinces of Orissa and Gondwana, and in the fatal campaigns of Rangoon and Upper Ava, in the first Burmese war, I should say with Dr. Ainslie, that it is alone in simple uncomplicated dysentery that ipecacuanha shows its best effects, administered as an aid to blood-letting and moderate purging.

In hepatic dysentery—a frequent complication in Bengal, especially during the cold season-calomel is absolutely necessary to the cure. I remember treating, for this form of the disease, an officer who had suffered severely from the fever of Batavia, contracted at the capture of Java; he was bled generally and by leeches, followed by sudorifies and purgatives; but no amendment took place, and nothing was voided but mucus and blood; two full doses of calomel with antimonial powder were then given at bed-time, which produced copious biliary discharges, followed by immediate relief, a few doses of blue pill and ipecacuanha, followed by mild aperients, concluding the treatment. There existed in this case no enlargement of the liver, nor uneasiness on pressure, but there was a total absence of biliary secretion, and, until that was restored, the other treatment afforded no relief. In another case of very severe hepatic dysentery, requiring remedial measures of great activity, recovery was marred in convalescence by errors in diet; the liver became enlarged and painful as ever, the dysentery recurring, and requiring a repetition of general and local blood-letting, mercury, &c., and this under circumstances of greatly reduced strength. This gentleman was sent to China, where, six months afterwards and owing to continued errors in diet, hepatic abscess took place, from which he died.

Cases of hepatic complication, treated without mercury, frequently terminate in inflammation and abscess of that organ, even where blood-letting has been employed. It is not enough here to reduce the volume of the blood, nor yet, along with it, to diminish the force and frequency of the circulation; we must also emulge

the gland involved in disease.

To such as prefer ipecacuanha in the treatment of dysentery it might prove desirable to ascertain the following circumstances:—

1st. Whether extract of gentian have any and what effect in preventing vomiting under the use of this drug. It is true that, after a few days' use, a comparative tolerance of medicine seems to be established, as in the case of the antimonial preparations; but altogether, as it appeared to me, unconnected with the use or disuse of bitter extract.

2nd. Whether the preparations of the Cannabis Indica may from their sedative, as well as from their powerfully antiemetic qualities, prove serviceable in conjunction with ipecacuanha, or with antimonials. Unlike opium and its preparations, hemp

does not lock up the secretions.

3rd. Whether four grains of ipecacuanha occasion as much sickness as ten grains, and, if so, whether the larger dose should be preferred for the cure of dysentery. If the action of ipecacuanha be purely revulsive, or if, according to Dr. Paris, it be to abate both the velocity and the force of the heart's action, so as to

affect "the whole series of blood vessels from their origin to their most minute ramifications," we may do wrong by being sparing of our dose, in so formidable a disease as dysentery, if the stomach can be made to bear the larger quantity. This subject has not, I think, received the attention due to its importance, in reference to the subject here under consideration. "Ipecacuanha has long been usefully employed," says the Brit. and For. M. C. Review, for July, 1855, "in various morbid puerperal conditions. M. Legroux is said to resort to it with almost uniform success at the Hotel Dieu, in sub-inflammatory and congested states of the uterus following delivery."

For the following interesting table, exhibiting the prevalence and the intensity of dysentery, as indicated by the mortality in

various countries, I am indebted to Colonel Tulloch:-

	Period		Dysentery.			
STATIONS.	of Observation.	Aggregate Strength. Attacked. D		Died.	Proportion of deaths to admissions.	
Windward and Leeward	20 years	86661	17843	1367	1 in 13	
Jamaica	20 "	51567	4909	184	1 in 263	
Gibraltar	19 "	60269	2653	64	1 in 411	
Malta	20 "	40826	1401	94	1 in 15	
fonian Islands	20 ,,	70293	3768	184	1 in 20½	
Bermudas	20 "	11721	1751	36	1 in 48%	
Novia Scotia and New Brunswick }	20 "	46442	244	18	1 in 13 g	
Canada	20 ,,	64280	735	36	1 in 20½	
Western Africa	18 "	1843 22714	370 1425	55	1 in 7	
Cape of Good Hope	0 "	8973	751	69	1 in 321	
Mauritius	"	30515	5420	285	1 in 11 1 in 19	
Ceylon	20 "	42978	9069	993	1 in 19	
Cenasserim Provinces .	10	6818	1460	137	1 in 10#	
Madras	5 ,,	31627	6639	559	1 in 12	
Bengal	5 ,,	38136	5152	411	1 in 124	
Bombay	5 ,,	17612	1879	151	1 in 121	

## CHRONIC DYSENTERY.

ONE has but to behold a soldier in hospital under chronic dysentery to grieve that a disease so destructive had not been cut short in its acute stage; for truly does William Fergusson declare that the sufferer exhibits "a spectacle of distress of as pitiable a kind as can be found in the history of human misery." Let the young naval and military surgeon ponder on the spectacle before him, and determine so to master the subject and manage the treatment of acute dysentery, as that he may have the least possible amount of the chronic to deal with. He will thus by one act cure the acute, and prevent the chronic disease.

It is not intended in this place to enter into the details of chronic dysentery, the reader being referred to an article on that subject in the second part of this work, and which describes the disease as it occurs in European invalids on their return from tropical climates. But chronic dysentery in the East Indies differs from that disease as it appears in persons on their return to Europe in this—that in India a certain amount of congestive and inflammatory action still remains, both in the mucous membrane of the large intestine and in the associated organs, requiring moderate topical blood-lettings, mercurial alteratives, and blisterings; whereas in the same disease, as it appears in England in the person of the tropical invalid, neither of the above conditions hold.

With the important exceptions stated, the treatment of chronic dysentery should be the same in both hemispheres; and in that of the East as of the West I would strongly urge the internal and external use of the nitro-muriatic acid. It is a powerful means as a substitute for mercurials, powerful in maturing convalescence.

## NOTE ON DYSENTERY IN THE NATIVES OF INDIA.

A LARGE proportion of the sickness and mortality of the native troops throughout India occurs from dysenteries and diarrheas, originally contracted as such, or resulting from fevers remittent and intermittent; and this holds especially in respect of the troops recruited in the plains of Hindustan, when they are transferred thence to the low marshy countries along the courses of the great rivers, or into countries abounding in jungle.

The late Mr. John Tytler, of the Bengal Medical Department,

in an Essay on "a most melancholy disorder," which he terms Diarrhea Hectica, states that "it is no exaggeration to say, that of the total deaths among the lower orders of Hindustan, three-fourths are the effect of this disease, either idiopathic or as a terminating symptom. It is the scourge of crowded jails, and of hospitals filled with harassed and fatigued Sepoys; and it is the great avenue through which the swarms of naked and famished paupers and miserable infants that occupy the bazars are continually vanishing."

Mr. Burnard, speaking of the sufferings of the Sepoys in Aracan, under General Morrison, says: "Dysentery and diarrhœa were the diseases which proved most fatal, not in their acute form, but ensuing as a consequence of fever; and their ravages, parti-

cularly amongst the native troops, were very great."

"The average mortality amongst natives in the Calcutta Medical College Hospital," says Dr. Macpherson, "has been 16.9, that among Europeans, 22.5. This accords with general experience, which has shown the disease in natives to be more amenable to treatment than in Europeans."

The symptoms of dysentery in the Asiatic races are precisely the same as in the European, differing only in degree of severity and in complications; and the same may be said as to the patho-

logy of the disease.

The post-mortem appearances exhibit ulcerations in the large intestine, amongst the natives of India, as with Europeans; but the complications in the dysenteries of Asiatics are by no means either so frequent or so severe, if we except that of the spleen in malarious districts, in which dysentery and diarrhoa are commonly associated with the remittent and intermittent fevers of such localities. Here the ulcerations of the large intestine are always of an

anæmic, atonic character.

Excepting in the instances of powerful and athletic Sepoys, and those mostly of the better-fed Mahomedans, I seldom had recourse to general blood-letting in treating natives for this disease—leeching being in the far greater numbers quite sufficient. The same regard to measure holds good in the use of purgative and other remedial appliances, the mildest in moderate doses being effective to the cure; and mercury in any form is seldom required, unless it be to the more robust, especially the Mahomedan, to whom I had often to exhibit one moderate dose of calomel with ipecacuanha or antimonial powder, followed by a brisk purgative, with the view to remove jaundiced appearances of the conjunctivæ, indicative of hepatic congestion. If such indications were not present, I never had recourse to mercurials in any shape.

The great principle of treatment in the dysenteries and diarrhœas of the natives of India, in their acute stages, is what Mosely

would term revulsive—sudorific; ipecacuanha being the best means, aided by gentle aperients, farinaceous diet, fomentations to the abdomen, and warm clothing. When by such means we have overcome the more acute symptoms, the Dover's powder will very fitly complete the cure; and the assistance to be derived from proper diet and warm clothing should never be overlooked. Cold acts most banefully on the native constitution, and too much attention cannot, therefore, be paid to the subject of suitable cover-

ing by night and day, especially during the cold season.

For maturing convalescence from the ordinary dysenteries of India, the nitric, or nitro-muriatic acid in bitter infusions will be found very efficacious; but where convalescence is retarded by continuance of liquid discharges, the acetate of lead and opium will prove greatly effective, the lead being exhibited in full doses. I had recourse to the acetate of lead in treating the native soldiery. after the more acute symptoms had passed, very early in my service in India, and afterwards very extensively in treating the dysenteries of the natives of Calcutta, while surgeon to the Native Hospital of that city. My excellent assistant at this institution, Mr. O'Brien, has for more than ten years subsequently, as surgeon to the 4th Regiment of Native Infantry, Gwalior Contingent, carried out this plan of treatment with great success, from the very onset of the dysentery. The Native Hospital formula was acetate of lead, tincture of opium, and distilled vinegar. This treatment has continued to be very successful in the hands of Mr. O'Brien as well as in mine; and he says that "when the liver is complicated, appropriate treatment is directed to that viscus; but acetate of lead is given at the same time without at all interfering with the other remedies. I do not mean to say that I employ lead solely; for I employ leeches, aperients, and blisters, which are very generally indispensable." The sulphate of copper and opium are also an excellent combination in the chronic stage of the disease. At all stages, indeed, and under every degree of severity and of distress from this disease, we are limited, in the treatment of natives, to the use of medicines by the stomach alone, as enemata they will not consent to use—the very impression of death making no alteration in their fixed aversion to every such remedy.

## ON THE DISEASES OF THE LIVER.

THE most cursory reference to the articles on fever and dysentery must satisfy the reader that, whether as original or secondary affections, inflammation, congestion, and chronic enlargements of the liver constitute in reality very frequent and very important diseases in the East Indies. Nor do the official returns, under the special heads of acute and chronic hepatitis, afford any approach to the actual frequency of hepatic diseases; for when these last occur as complications with, or as sequelæ to fevers, dysenteries, diarrhœa, and cholera, the hepatic complication or sequel remains unnoticed in the numerical returns-the cases being classed and numbered under the primary disease, whether fever, dysentery, diarrhœa, or cholera. It thus happens that the numerical hospital returns do not yield anything like an approximation to a true estimate of the existing amount of hepatic disease. The same observations will, I believe, apply with equal justice to the returns of acute diseases of the West Indies, the West Coast of Africa, the Stations in the Mediterranean, and the French Possessions in Algeria, although in a lesser degree.

INCREASED SECRETION OF BILE.—On the subject of the physiological influences of the climate of India on the functions of the liver, I must refer the reader to what is stated in the introductory chapter on climate. It is there stated as a generally received opinion, that an increased secretion of bile is one of the ordinary results of the exposure of an European to an unnaturally high range of temperature; and this opinion I believe to be wellfounded to the extent, first-that the exaltation of function is not of long duration, and is very much confined to the earlier years of residence, declining thence; -secondly, that, to a certain degree, such increase of secretion is salutary ;-and, thirdly, that it is in part from the sudden suppression of secretion through cold, that congestive and inflammatory states of the liver are engendered, rather than through any serious injury from increase of secretion-a process consisting more of acclimation than of disease, and one which nature adjusts in the progress on residence in India. So much has been said by some writers on tropical diseases as to morbid increase and vitiation of bile, that they have altogether forgotten the more important but less notable deficiencies and suppressions of this secretion-conditions far more serious in their results than mere excess of secretion.

Chemically, we have as yet ascertained nothing as to the healthy or unhealthy qualities of bile; and, practically, all we do know is, that with certain symptoms there is a redundancy, and with others there is a deficiency of that secretion. We know also, from the scalding sensations that accompany certain forms of bilious diarrhea, that the fluid in question is of an acrid irritating quality; and that is all.

TREATMENT.—If, along with an aerid and irritating bilious diarrhœa there be fulness and uneasiness in the region of the liver, so as to excite the suspicion of congestive action, a few leeches followed by fomentations and warm baths, together with a mild mercurial and an aperient will procure relief; but when heat and dryness of skin exist, with irritative febrile symptoms. effervescing draughts, containing antimonials and diureties, will prove both grateful and curative.

In general, the disorder constituting increased biliary secretion, and those resulting from it, are sufficiently cured and prevented by avoiding the direct rays of the sun, by a strict attention to diet, exercise, clothing, bathing, and by the use of demulcent and subacid fruits, as the bael, pomegranate, orange, &c., and by the oc-

casional use of mild aperients.

With few exceptions, and these resulting from exposure to direct solar influences, or from ill habits of life, such simple means of treatment I have always held to be both suitable and effective: and I cannot but think much unnecessary confusion has arisen from wordy descriptions of what certain writers on tropical diseases, as Annesley, appear to regard as an affection demanding a complicated medical management. They talk of "bile" as if it were a disease, and would cure it by "full doses of calomel at bed-time, generally for five or seven days, followed by Pil. Aloes and Myrrh alone, or with Pil. Hydrarg." Of such unmeasured. disproportionate, and unnecessary treatment of a mere functional disorder, it is to be hoped we have seen the last:-it is really worse than routine. The real diseases connected with bile-and very serious they are too—are those in which bile is altogether absent, and wanting to the cure!

CONGESTION OF THE LIVER .- Congestions more or less acute, according to constitution, previous condition of health, and season, are very common to all countries in the East. Sudden changes of temperature, producing chills—the repeated cold stages of fevers, remittent and intermittent,-an over-full and stimulating diet, together with the abuse of vinous and spirituous liquors—violent bodily exercises, especially those carried on in the sun, exposing the individual to the more ready influences of

cold ;-those are the more ordinary causes.

The organ will be found enlarged, generally in its upper convex direction, but sometimes in all directions, as in malignant intermittents—the enlargement resulting from venous and biliary congestion, and the latter being produced by the pressure of the overdistended veins preventing the due diminution of the smaller ducts—a condition of the liver, I repeat, frequently associated with, and following upon, remittent and intermittent fevers.\*

The bulging out of the liver—the evidence of severe vascular turgescence—is rarely accompanied by more than uneasiness on exploration, or of weight on getting into the erect posture:—the respiration on the affected side will be found more or less oppressed and impeded. These symptoms will become well marked when, as occasionally happens, the volume of the liver is much augmented; and when such is the case we may fairly infer that stagnation and consequent viscidity of the bile will take place, both in the gall-bladder and ducts, so as to complicate and aggravate the disease.

The countenance and complexion will exhibit pallor, or else a dusky livid, sallow hue, according to the temperament of the patient, or the duration of the disease—paleness, with sense of cold and shivering, being more characteristic of the immediate result of external cold, or chill. The tongue will generally be found coated—the bowels constipated—the intestinal and viscal excretions depraved—the appetite defective, with occasional nausea or even vomiting, and more or less headache. The pulse is slow, oppressed, irregular, or quick and feeble, according to the age and constitution of the patient, and to the duration of the disease; but the frequency of pulse will not be that of fever. Here, in short, the symptoms are obscure—not declared—and they must be taken in their collective characters, assemblage, and connection, and not separately, or individually.

When, through any or all the causes above enumerated, congestion of the liver has been of long duration, permanent injury to its structure and secreting function may result:—the disease in any form must, therefore, be regarded as one of importance. The signs in this, as in other diseases of the liver, are so obscure that of such as we actually possess a knowledge we cannot afford to overlook the least, if we would do justice to our patients, and to our own characters. On examination after death, the liver is found to be enlarged, principally upwards into the right thoracic cavity—dark from impeded circulation, and full of blood; and,

<sup>\*</sup> Passive congestion of the liver, and torpor of that organ, as they appear in the returned Indian, are noticed in the second part of this work.

when the disease has been of any duration, its texture will be found softened.

TREATMENT.—When from careful exploration, and a due consideration of all the attendant symptoms, we are satisfied that congestion of the liver does actually exist—for its existence must not be assumed—a moderate venesection, followed by leeches if necessary—a few doses of blue pill or calomel, with purgative extract and an antimonial, all three united—a few saline purgatives—hot fomentations to the region of the liver, and warm baths;—these means, together with abstinence in diet, will speedily bring about a cure. Nor must we be deterred—all other circumstances being duly considered—by feebleness or oppression of the pulse, from the use of such antiphlogistic means in these cases; for, under judicious treatment the pulse will rise and expand under the use of these means, and general relief will follow.

Finally—Convalescence will be matured, and an exemption from future attacks secured, by attention to diet, exercise, bathing, attention to clothing, and by avoidance of exposure to direct

solar influences, as well as the chills by night.

HEPATITIS.—Amongst the better classes of Europeans, who are careful in their habits, this disease is not of so frequent occurrence as formerly, when the habits of life were different from what they are now; and this observation has special reference to the acute inflammatory, or adhesive form of the disease. The cases seen in private practice are very much confined to those of fair complexion, and lax fibre. European females of every class are far less subject to this disease, and to every form of hepatic affection, than their countrymen; and women of the better classes are rarely affected comparatively. Amongst British soldiers, and Europeans who lead what is called a free life, and who incur exposure to the direct rays of the sun, and to the damps and chilling influences of the night, the disease, on the other hand, attacks men of all ages, and of every variety of constitution. It is a disease of great importance in a military sense, causing much immediate loss of life, invaliding, and ultimate inefficiency in the European portion of the army. It would appear from various returns to which I have had access that, of British soldiers attacked. upwards of half were lost to the service by death, by ruined health in India, and by invaliding to England. In the Madras European Regiment, according to Mr. Geddes, 280 cases of hepatitis occurred in five years.

SYMPTOMS AND PROGRESS.—The symptoms and progress of hepatitis vary much according as the inflammation is

seated in the upper convex surface, on the exterior of the organ, or in its central parenchyma; but the nature and amount of the inflammation are assuredly the first and most important considerations to be determined, if we would successfully combat so dangerous a disease.

For practical purposes hepatitis may be considered, first—as consisting in a more or less superficial and adhesive inflammation of the organ; and secondly—in deep-seated suppurative inflam-

mation of its parenchyma.

1. This well-marked form of the disease is ushered in by rigor, or sense of cold and constriction of the skin, followed by febrile reaction, the complexion being of a dusky sallow hue, nausea and vomiting, urgent thirst, loaded tongue, occasional headache, pain of the right hypochondrium, increased by pressure, or a full inspiration, and more or less acute according as the peritoneal covering may be more or less involved, pain occasionally of the right shoulder, cough and oppressed breathing, difficulty in lying on the left side, scanty and high-coloured urine, loss of appetite; such are the usual and more urgent symptoms. They will not all be present in any case; but so many of them will be manifested as to render diagnosis generally easy. There occurs frequently a loose state of the bowels, the matters voided being pale, or muddy, and watery; at other times the bowels are even constipated. When vomiting is urgent, with sense of heat and burning in the epigastrium, it indicates an acute state of the disease involving the ducts, duodenum, and pylorus.

As the disease advances the pulse becomes frequent and hard, the skin is hot, dry, and constricted, the pain, cough, and oppression of breathing increase, there is occasional hicup, with an anxious and desponding countenance, all indicative of advancing disease; and if this condition be not speedily arrested by remedial means, abscess of the liver, destroying life, or permanent enlargement of the liver, destroying the general health, will inevitably result.

In this formidable disease we should regard the severity of the hepatic and epigastrial pains, and other distressing symptoms as really invaluable indications; for while they warn us of a pressing danger they explain everything, and thus relieve the mind from what otherwise would be a most painful state of doubt and anxiety. In the form of hepatitis which is attended with so much suffering as is here described, the fate of the patient is comparatively secure; for even the inexperienced will be induced to apply immediate and decisive remedies.

2. It is different in respect to the deep-seated and suppurative inflammation of the parenchyma of the liver; for here the danger is even greater than in the former instance, with no salutary warnings from symptoms. It is one of the most dangerous of

diseases, because of the total absence of urgent symptoms, and consequent insidiousness; the process which leads to destruction is here silent and rapid.

Of various kinds of inflammation the suppurative is by much the more frequent, in Bengal at least. It attacks persons of both sexes, and those even of the most temperate habits; it occasionally

terminates the career of the old Indian.

Whether occurring in the person of the older resident in Bengal, or in the new comer, this form of the disease occurs very generally in the cold season, and is caused by exposure to alternations of heat and cold, by such means, in short, as determine powerfully from the surface to the internal organs. I have seen cases where it resulted from the chilling thorough-draughts of the northern entrances to the Calcutta houses, on issuing from a ball-room, for instance; and other cases where it was occasioned by exposure to damp and cold before daylight for the purpose of hunting.

The disease is sometimes preceded by a perceptible falling off in the general health, such as emaciation, dry cough, and embarrassed respiration, loss of appetite, the complexion gradually assuming a muddy sallow hue, but it more generally comes on in the midst of apparent health. We seldom indeed see the patient till inflammation has actually commenced; when he generally complains of a feeling of abdominal uneasiness, but more particularly of the epigastric region and that of the liver, with some degree of fever preceded by slight rigor or ague; but all these may be so slight as too often to attract but little of the patient's attention. Perhaps he consults his physician on account of diarrhwa, supposed to result from errors in diet; medicine affords some relief. and the patient proceeds in his ordinary occupations for days, or, when the action is less acute, for weeks, though under great depression of the mental and corporeal energies, till at length his altered appearance, hacking cough, permanently dry skin, invincibly rough furred tongue and morbid taste, all expressive of a suppressed and depraved state of the secretions, attract some more serious notice on his own part, and that of his family. The real nature of the disease may still remain a secret to both patient and physician; and it may not be till actual tumour of the liver, a marked succession of rigors, or profuse and clammy sweats announce in unmistakeable terms the formation of abscess, that either party becomes awake to the impending danger; and then it is too late. A sense of uneasiness, hardly amounting to an obtuse dull pain, a sense of weight and oppression, may or may not exist in the region of the liver, according as the disease is centred more or less deeply in its substance or in its upper convex surface; when the former exists the symptoms are more than usually obscure and insidious; in the latter case they are somewhat more of an acute nature. In this form of hepatitis it is

seldom that we discover any pain in the shoulder.

I should say, then, that a rigor, or a diarrhœa,

I should say, then, that a rigor, or a diarrhœa, followed by feverishness, the peculiar harsh state of the skin, the tongue having the roughness of a coarse file with adherent coating, together with the local uneasiness already described, cough, scanty surcharged urine, often of a deep saffron colour, ought immediately to warn the physician of the suppurative inflammation which speedily leads to abscess. In every instance the diagnosis will receive material assistance from the external examination of the chest, especially when the upper convex surface of the liver is the seat of disease. Nor should the exploration of the abdomen be less carefully made; nothing should satisfy the medical attendant as to the condition of his patient short of the most painfully assiduous explorations often repeated; for we have here to deal with a disease deadly in its nature and rapid in its progress.

CAUSES.—The causes of hepatitis will be found in the following circumstances:—residence in hot climates, exposure to high ranges of temperature, followed by cold and damp, causing sudden arrests of the cuticular as well as of the internal secretions, the cold seasons of tropical climates, night exposure while imperfectly clothed, the exposure of the bivouac. These causes become very powerful when fatigue is added to them, and under irregularities of diet, and intemperance in the use of wine and ardent spirits especially. Dysenteries and fevers, remittent and intermittent, a too rich and stimulating diet, producing gastric and duodenal irritation, these are frequent and influential causes of hepatitis; and more remotely may be numbered previous attacks, malaria, various forms of dyspepsia, neglect of the natural action of the bowels, the age of puberty.

PATHOLOGY.—Were the earliest morbid change in the liver ascertainable antecedently to actual inflammation, I have no doubt that that condition would in most cases consist in acute congestion; for, as in the well known instance of hepatic congestion and consequent diarrhea, both resulting from the sudden application of the cold damp of the Himalayahs, for example, so the pale, watery, premonitory diarrhea is one of the most ordinary and immediate precursors of hepatitis; it is generally the earliest and indeed the only symptom which attracts the notice of the patient; and it should always receive the instant and close attention of his medical attendant. In two cases of hepatitis in recently arrived Europeans, and which terminated fatally, Mr. Twining found "the internal membrane of the hepatic veins inflamed, and puriform matter in the right ventricle of the heart." Their com-

plaints, he says, "began as common diarrhoa of a severe description." In the case of an European soldier in Burmah, treated by Dr. Murchison, and in which inflammation terminated in abscess, "an obstinate diarrhoa had been one of the most harrassing symptoms in this case; but no traces of ulceration were found either in the small or large intestine." These observations on the earlier pathological changes, refer to every form of hepatitis, but more especially to the deep seated suppurative inflammation of the parenchyma of the liver; and he who will best ascertain them at their commencement will least be taken by surprise in the progress and terminations of this disease. In the suppurative inflammation the earliest observable lesions I believe to be redness and softening, soon to be aggravated into the formation of abscess.

Mr. Twining declares that he has "never seen a case terminate in abscess without our being able by a careful examination to detect the disease that is in progress long before there was any reason to believe that suppuration existed." This may be generally true, but then the examination must be most rigorous and often repeated, until we are quite satisfied as to the true nature of the case. The patient's recovery or death will depend on the care and discrimination with which this examination is made; a false diagnosis is here a death warrant, they are indeed but convertible terms. I remember that, in the cold season of 1837 alone, in Calcutta, I was consulted in four cases of suppurated liver, in which no such condition had been previously suspected.

The appearances on dissection of Europeans who have died of hepatitis will vary with the intensities and the duration of the disease. Various vascular discolorations of the peritoneal covering are observed, differing according to the degrees of inflammation, and the number of previous hepatic ailments—thickenings of the peritoneum, and effusions under it being very generally discoverable. Adhesions of the liver to the colon, stomach, and diaphragm are also common.

Where inflammatory symptoms have run high, and death has ensued rapidly, there will be great enlargement of the liver, with much redness and vascularity of the organ, with depositions in the interlobular cellular tissue; in other cases its colour will be dark, or even black—bleeding freely in both instances on section being made. In other cases the covering membrane is found of a pale hue, thickened and indurated—the structure underneath bleeding but little on being cut; while induration, more or less extensive, with deposition of coagulable lymph, tubercles, and concretions

on the concave surface of the liver, the gall-bladder is often enlarged and filled with viscid bile; in other instances, diminished in volume, its coats thickened, and the body covered with false membranes.

The observations of Rose and Henry would seem to establish that the quantity of urea excreted by the kidneys is lessened in suppurative hepatitis apparently in a degree proportioned to the extent to which the secretion of the liver was destroyed by the abscess. "If the statement be proved by future observation," says Dr. Parkes, "it is, in my opinion, impossible to overrate its importance in a pathological as well as in a physiological point of view."

Since my return to England I have on several occasions tried to obtain from India information on this and on other important matters connected with the chemistry of the blood and of the secretions in tropical diseases; but some officers said they were over-worked, others that they were situated too far from the Presidencies, and so on. Thus I have failed in my object; but I hope yet to succeed.

TREATMENT.—The maxim of De Retz that "what is absolutely necessary is never dangerous," is true in medicine as in politics: nor is there here any difficulty in perceiving either what is necessary to be done, or yet what is safe in the treatment of acute hepatitis; it is likewise very easy to comprehend what it would be dangerous to leave undone. The greatest danger will indeed be found in feebleness, irresolution, and in do-nothing

However long the disease may have existed, and in whatever part of the liver the inflammation may be situated, provided there be no symptoms indicative of suppuration, general blood-letting, repeated as the symptoms may demand, and copious in relation to age, health, and length of residence in India must be had recourse to; and the sooner the better for the patient, the measure of depletion being, in all cases, a sense of local and general relief—with softening of the skin. These are the only criterions of adequate loss of blood; and it should be always held in recollection that inflammation, suppurative especially, and of the most deadly character, is present, and that, consequently, there is not an hour to be lost.

After the general blood-letting, a full dose of ten grains of calomel, with five of James's powder is to be exhibited at bedtime, followed in the early morning by a strong saline purge, or a full dose of compound jalap powder; and this course must be pursued—using leeches to the side, mercurials with antimonials, purgatives, and diaphoretics with diuretics daily, so long as they are called for by fulness or tenderness of the hepatic or epigastric region. The medical attendant must remember that the subsidence of local symptoms, together with a declared and satisfactory abatement of all the general symptoms, including that of the force and frequency of the circulation, and a permanent relaxation of the

skin, with freedom in the secretions, can alone warrant the discontinuance of antiphlogistic means, including the most spare use of farinaceous food and drink. If the bowels be irritable, and anything like diarrhœa threatens to distress the patient, the Dover's powder, or pure opium, must be added to the calomel, in lieu of the antimonial.

As regards the constitutional influence of mercury, the utmost that I found necessary, even in the worst cases, was the producing a gentle action on the gums, salivation being unnecessary, and in the instances of persons of a feeble or strumous habit, naturally injurious. The precautions already insisted on against oozing from leech-bites must here be carefully regarded; while blisters will be found useful on the entire disappearance of inflammatory action.

Nor should the most strict abstinence in diet be deviated from, until, by the subsiding of all uneasiness, and tumidness of the region of the liver, and of the abdomen generally, by the cessation of all febrile movements, and by the restoration of healthy secretions, we feel assured, on these sufficient evidences, that convalescence is well established:—and even then, the return to accustomed habits should be made very slowly, so as to afford time, under great abstemiousness, for the absorption of instertitial deposits; a reparative process which cannot take place under a full stomach.

Mild saline aperients in bitter infusion, the nitro-muriatic acid in bitter infusions—the bicarbonate of soda in large doses with bitter infusion, where there is much acidity—warm baths, and gentle exercise in the open air—all tend to the maturing of convalescence; and, in the cold season, great attention must be paid to the clothing. Nor should the European soldier here, any more than under fever or dysentery, be permitted to return to the barrack life, or to duty, before his strength has been restored: it is to the neglect of this rule that many of the serious and often fatal relapses are due.

When convalescence proceeds unkindly, and there is reason to apprehend structural injury to the liver, a full course of the nitromuriatic acid, as described in the second part of this work, will be found the surest and best remedy.

Mr. Twining relied too exclusively on blood-letting in hepatic inflammation, and too little, as appears to me, upon other remedial means. There are cases where inflammation will go on notwithstanding the most copious depletions, and others in which the same will occur with a debilitated constitution and a weak pulse. What then is to be done: "What degree of depletion," says Dr. Billing, "will remove a node, or syphilitic iritis, without mercurial and other medicines? What would venesection do for rheumatic pains without antimony, colchicum, opium, bark, mercury, and

other medicines? The question is not so much as to possibility, as expedition and safety to the constitution." Robert Jackson—no advocate of mercurial treatment in tropical diseases—admits that "mercury, after the condition has been prepared for its action, is the principal means of cure where biliary secretion is principal.

pally in fault."

Mr. Twining, who was in the habit of prescribing half scruple doses of calomel at bed-time, in hepatitis, asks the following question respecting the action of mercury in this disease:—"We have been told that mercury is beneficial in hepatitis in the way that the breast-pipe or pump relieves inflammation of the female breast, when milk abscess is impending. But a legitimate parallel cannot be established in the action of the respective remedies; the pipe relieves the inflamed breast by drawing off the superfluous milk, without exciting increased secretion. Has the action of mercury

an analogous effect on the liver?"

No: and if it had but the limited effect of the pump or pipe, it would not be the valuable remedy physicians and surgeons have ever found it to be in so great a variety of inflammatory affections, and in the deadly inflammation of the liver in particular. It is for the very reason that calomel assists powerfully both in "drawing off" accumulations, and in promoting "increased secretion," that it proves of such value in aid of blood-letting. It is, in fact, by this very double action, of purging and increasing secretion at the same time, that calomel relieves the loaded and inactive vessels of the diseased gland; not to speak of the other acknowledged physiological influences of this mineral, such as-its increase of all the secretions and excretions of the body—its influence on the capillary circulation-its febrifuge effect-the peculiar specific power ascribed to it by physicians and surgeons, as an antagonist to inflammations, whether general or local-its stimulant power over the absorbent functions—its power of unloading, at the same time that it gives a new impulse to, the vascular system-its peculiar power in removing viscid and tenacious intestinal secretions -its antiphlogistic, solvent, and alterative effects on the blood;these are the actions and uses ascribed to mercury by the ablest British practitioners and authors, and they are such as to place this remedy second only in order of importance to blood-letting in all the more acute hepatic affections of India.

That mercury enters into intimate union with the elements of the blood is now an ascertained fact: it must therefore "modify its plasticity, and influence all the organic functions to which it is subservient." That mercury has been, and is still, abused by illinformed and inconsiderate persons, is quite true, both at home and abroad; but to argue hence its disuse would not be either

safe or wise.

It is of great practical importance that the real benefits derivable from the use of mercury be impartially set before the inexperienced tropical practitioner, and I have here endeavoured briefly to do so, I hope, with fairness to the subject, and to persons with whom I may happen to differ in opinion. In the pride of pathological science we are induced to believe that, because we see and examine, we therefore understand; and we too often neglect the science of pure observation so inestimable to the physician; and thus, I think, it happens that matters are still in doubt with us which a more chastened and correct observation ought long ago to have fixed and settled.

When inflammation has terminated in hepatic abscess, mercury given with the view to its general influence, is always highly injurious:—it aggravates the irritative fever and general distress, while its specific influence—so hopeful in the previous stage—now but aggravates the danger by increasing the suppurative process, and the general debility, so as to retard, or altogether to obviate the curative efforts of nature.

Shortly before I left Calcutta, a case came under my care, in the month of August, in which slight hepatic tenderness existed in complication with severe dysentery. From both diseases the patient speedily recovered; and to restore his strength, I advised a short voyage to sea, where, having exceeded in diet, the liver became tender and enlarged, there being no recurrence of dysentery. A second time he recovered, but being of a weakly habit, and now much reduced by disease and treatment, I sent him on a longer voyage to Pinang. Here again, despite the most careful written instructions, he took to excesses in eating and in the use of wine: - the result was that, in three months from his leaving Calcutta, an hepatic abscess formed, and discharged itself through the stomach by vomiting, soon after which he died. I mention this case as affording an illustration of the necessity for abstinence in diet, and to show that, even when the hepatic disease is but secondary, it may, and very often does, prove the ultimate cause of death by the formation of abscess.

In 1837, a young staff officer consulted me on account of cough and emaciation, which he considered to indicate tubercular consumption, as some relatives of his had died of this disease in England His respiration and pulse were very rapid, and he had profuse night sweats; in short, I found that he was dying of suppurated liver. The history he gave me was to the effect that, six weeks previously, he drove in an open gig from a ball-room in Calcutta to Barrackpore, and that on getting to his quarters, he was seized with shivering, followed by feverishness and diarrhœa; he continued to perform the duties of his office up to the day on which he consulted me, and the real nature of his disease had not

till then been so much as suspected. He died in a fortnight, and the entire substance of the liver was found to be converted into a huge cyst filled with pus.

Well might Curtis, writing in 1781, speak of hanging up cases such as these "in terrorem, against the attacks of this insidious disease;" or rather, he might have added, as warnings against the crimes of heedlessness and inattention in cases of so pressing a

duty.

I will now present a case the converse of the above:- In the cold season of 1836, a professional friend called on me to consult respecting the case of a patient of his; and just as he was taking leave, he said, as if by accident, that he had "a pain in his back tike lumbago." Being an able experienced surgeon, I might well have said, "go home and take care of yourself;" but there was a something in his expression of countenance which excited a suspicion of diease, and so I made a careful examination. The result was the discovery of deep-seated inflammation of the liver, and that it had existed for three days, during which he had been living as usual, and using the cold bath every morning. All the patient had noticed was a slight shivering three nights previously, followed by feverishness and pain in the back; but he considered his symptoms of so little moment that he felt a doubt as to the accuracy of the diagnosis. I then referred him to our common friend, Mr. Nicolson, of Calcutta, who concurred with me completely as to the nature of the disease. The patient was young and of robust habit, so that with the loss of about eighty ounces of blood in the first twenty-four hours, followed by calomel and antimony gently to affect the gums, strong purgatives, and total deprivation of food, he rapidly recovered; but I think he recovered with difficulty; a few more hours lost to the treatment, and it would have been too late.

The above case is interesting and important, as showing how very obscure and insidious are the symptoms, and how little they possess of urgency to cause a salutary alarm, even to the mind of a professional person. But it is always thus when the inflammation is centered in the parenchyma of the liver; and hence the absence of that acute pain, and those urgent symptoms which characterise inflammatory states of the peritoneal covering of the gland, and which generally give ample warning.

HEPATIC ABSCESS.—The formation of abscess in the liver, as already stated, is generally announced by irritative fever, increased towards night—rigors or shudderings, followed by profuse cold sweats—a tenaciously furred rough tongue—scanty and high-coloured urine, &c. &c.; when the abscess is large, by a perceptible tumor of the right hypochondrium—the mind being anxious and depressed.

by Dr. Budd, in a case of jaundice from a closure of the common duct, and in a case where a cancerous ulcer of the stomach had eaten into the adherent liver; and by Dr. Graves, in an inflamed gall-bladder. Dr. Parkes, again, has seen this state of tension very prominent in a case of phthisical diarrhæa, in which there was extensive alteration in that part of the ileum situated immediately under the muscle; while partial contractions of the abdominal muscles, over a suffering part, are justly stated by Mr. Holmes Coote, and by Dr. Kirkes, to be frequent in jaundice and other affections of the abdominal organs.

We may, therefore, conclude with Dr. Watson, that this sympathetic affection is but one instance among many of that kind of protective instinct whereby a tender part is in some measure

shielded against the infliction of pain by pressure.

TREATMENT.—It is little, indeed, that we can do for the unhappy sufferer from hepatic abscess. When the abscess is small we may favour the efforts of nature at absorption by care in diet and clothing—gestation in the open air, and change of air—the use of the nitro-muriatic acid with bitter tonics, &c. When the abscess has happily discharged its contents into the stomach or bowels, or externally, a more nutritious diet will be proper, with the view to support the strength of the patient, and even a little wine may now be allowed. But nothing should be done which may excite the nervous or vascular functions; and mercury in every form is to be carefully avoided, and even the mildest

aperients should only be exhibited under necessity.

Of the instances of recovery within my personal knowledge, by far the greater numbers were amongst such as had discharged the pus by the bowels; and one married lady I remember to have recovered health rapidly, after vomiting the contents of an enormous abscess. I also saw four instances of recovery where the abscess had opened into the lung. Of those in whom the pus was discharged through the external integument, the majority have within my knowledge died. But by much the greater proportion of the instances of hepatic abscess end fatally-either before the bursting of the abscess, or within a few hours from its discharge into the peritoneal cavity:—and this brings me to the consideration of the propriety of making an opening for the discharge of such abscesses through the external integument. I have often seen this operation performed in India, and I have myself performed it; but in no instance that I remember did the operation appear to me to result in eventual good. I therefore agree with Dr. Budd, that " it is generally best, when an abscess of the liver projects at the side, to allow it to open itself. Nature performs the operation better than the surgeon. When the abscess opens of itself it is

usually by a very small aperture, like those of worm-eaten wo which never closes; and the matter gradually oozes out as the contracts. No air gets mixed with the matter of the abscess, a no violence is done to its walls; and, consequently, no fresh flammation is set up. The discharge is very gradual, and small in quantity as it can be. There is less shock to the syste and less drain from it than when the abscess is opened by knife. The prominent part should be poulticed, and the mat allowed to escape in the poultice, but should not be squeezed pressed out."

These observations accord altogether with my own experier in India where, too often, young medical officers are urged by pressing dangers of the case, and by the importunity of frien

to "do something."

Dr. Budd is deserving of attention on another question co nected with this subject. "In opening an abscess in the si stance of the liver," he says, "there is another, and greater, a more unavoidable source of danger, which has not been notice by the writers to whom I have referred. It is that the so hepatic tissue cannot readily collapse, so as to close the cay when the abscess is opened. When, then, a free opening is ma even into a recent abscess, air almost necessarily enters into cavity, and from the sudden removal of pressure, or, it may from the manipulation employed to empty the cavity, violence done to the walls of the abscess, and there is often some deg of hæmorrhage from them. Air and blood thus become mix with the pus in the abscess, decomposition takes place, and air, or the decomposed pus, sets up fresh inflammation of inner surface of the sac. This causes, of course, a fresh acc sion of fever, and of other constitutional disturbance, and if abscess be large, a profuse, and fœtid, and continuous dischar which may soon exhaust the strength of the patient. The condary inflammation thus excited by the presence of air, or the decomposed pus, may even lead to gangrene, and speed destroy life; and this happened in one of the cases that fell und my care at the Dreadnought. A similar case is noticed Cruveilhier."

Of fifty-seven cases of hepatic abscess, real or supposed, corded in the Medical Journal of Madras, for 1844, the following were the terminations:—

Recovery took place		in	17	cases
Of these the trocar entered the absce	ess .)	in	8	"
No pus obtained		in	3	
Pus obtained on the 6th, 7th, and 11th	h days	in	3	,,
No pus mentioned	)		3	22
Death occurred		in	40	39

The proportion of recoveries to deaths in eighty-one cases of hepatic abscess, recorded by Mr. Waring, of the Madras army, and in which their contents were evacuated by operation, was as follows:—

Recoveries		14	15,	or	18.519	per cent.
Deaths .			66,	or	81.481	"
			81	-	100.000	

Referring to 300 fatal cases of hepatic abscess, of which the details are given by Mr. Waring, in the first part of his work, he considers that operation might have been applicable in the following proportions:—

Of 177 cases in which abscess was solitary, there)	al number of cases re were a plurality of abscess	es in : : :	300 108
			192

Thus it appears that "the proportion of cases of hepatic abscess which hold out a reasonable hope of cure by the performance of an operation for evacuating the contents, is much smaller than most persons are aware of." . . . Out of the whole number above recorded, "only in 116, or little more than one in three, could the operation have been undertaken with any reasonable probability of success; and this number would, of course, be still further diminished by taking into account the cases in which abscess, though solitary, communicated with the lung, colon, or some other viscera, and those in which other organic disease existed."

JAUNDICE.—This disease occurs in India, as elsewhere, as an acute affection, and as a distinct and separate disease—often

unconnected with hepatitis, fever, or dysentery.

Jaundice is usually attended by mental depression and physical lassitude—dyspepsia with loss of appetite, nausea, a yellow furred tongue, and flatulence. In some persons there exists fulness and uneasiness on pressure of the epigastric region and that of the liver; while the amount of sallowness of the skin, and of the urine, together with the degrees of paleness and of absence c

bile in the alvine discharges, will depend on the intensity of cause of the jaundice. The characters of the disease will also with the age, constitution, habits, and length of residence India—age and length of residence being circumstances that certain to operate to the disadvantage of the patient, especia an anæmic state of the system be present at the same time.

The existence of gall-stones obstructing the ducts may be ferred from the following symptoms:—A sudden seizure of visits of pain in the epigastric and right hypochondriac region companied by irregular transient shiverings, and severe fivomiting—the pain at times being excruciating. Feverishne the irritative character now makes its appearance; and, we twenty-four or more hours, sallowness over the forehead, exing rapidly to the trunk and extremities, becomes evident, on the appearance of the jaundice pain usually subsides. well to note here, that the pain attending the passage of stones far exceeds that even of peritoneal inflammation, and the fever, when such is present, is of a subdued character, use the endemic fevers of India—diagnostic circumstances worth attention of the inexperienced.

In the East Indies, and as it generally appears amongst I peans there, the progress of the disease, under appropriate ment, is very generally of a favourable tendency—the cause I in most cases, of a congestive or inflammatory nature, susce of easy removal; and we find that as convalescence proceed urine gradually loses its saffron hue, while the skin also bee fairer by slow degrees. More rarely, though the jaundice is pleasen, so as to require no other diagnosis than is furnished the complexion, the causes of this complaint are nevertheless difficult to trace.

CAUSES.—Whatever causes may produce hepatitis, or especially congestion of the liver, will give rise to jaundice. tain affections of the brain have been known to excite this disast the extremes of despondency, and sudden bursts of pas but by far the more ordinary causes are—cold applied to the faces of the body, or cold applied suddenly to the stomach is form of iced drinks, when the body has been heated by export other exertion. In an analysis of seventy-two cases of it typhoides, by Professor Lebert, one-third of all the cases occi in November and December, or in two only out of the cold me Previous attacks predispose powerfully to jaundice. Form when extreme courses of mercury were used in the treatme venereal diseases, what with the action of mercury on the he function, and with the influence of external cold on the jaundice was a frequent disease in our hospitals at home

every case of jaundice that I have seen in England, amongst the returned Indians, cold has been the immediate cause of the disease.

Jaundice, according to Dr. Budd, may be produced in two ways:—first, by some impediment to the flow of bile into the duodenum, and consequent absorption of the retained bile; and, secondly, by defective secretion on the part of the liver, so that the principles of the bile are not separated from the blood. It is probable, he adds, that jaundice which follows closure of the common duct, does not result merely from absorption of retained bile, but also, in part, from the secretion of the liver being less active, so that the principles of bile are retained in the blood.

"But in many cases of jaundice, perhaps in the greater number, there is no impediment to the flow of bile through the ducts. In fatal cases, it happens not unfrequently that the gall-bladder and gall-ducts are found empty, and their mucous membrane is unusually pale—showing that no bile was secreted. The jaundice results solely from suppressed, or deficient secretion."

I think it right to place before the reader the views of this able writer, as to the causes of jaundice;—adding, however, that in India the more general causes are congestion and inflammatory affections, involving the gall-ducts, more or less directly.

PATHOLOGY.—As few persons die in India of actual jaundice, the opportunities for the examination of the liver and its appendages in such cases can only occur by accident, as when a sufferer from jaundice dies of another acute disease. Such occasions, however, have been sufficiently numerous to establish, on examination after death, the following appearances:—These consist in inflammatory states of the liver, and of the gall-ducts—congestions of the liver—deposition of coagulable lymph within the capsule of Glisson, with enlargement, more or less considerable, of its absorbent glands—effusion of lymph into the areolar tissue about the gall-ducts—and the presence of gall-stones impacting the common duct:—morbid affections of the duodenum are also found to be associated with jaundice.

Dr. Henry Kennedy, of Dublin, after narrating some cases of rapidly fatal greenish yellow jaundice, in which the ducts were found quite pervious, and in which the discharges from the bowels "though probably of a lighter colour than what is natural, were by no means free from bile," suggests that "a much deteriorated bile, the absorption of which would cause serious consequences," may have here caused death. Dr. Cathcart Lees, of the same city, has been led by the examination of the urine of jaundiced patients "to entertain the opinion that the absence of any symp-

toms referable to the nervous system in many cases of jau might be accounted for by the presence of cholic acid in the and its consequent absence from the blood." . . . . " supposition be correct, I think the detection of cholic acid, conjugates in the urine, may prove a very important elemprognosis in every case of jaundice, as assuring us again supervention of cerebral symptoms in cases where it is prophyla, if absent, it may put us on our guard, and possibly out to adopt a prophylactic treatment."

TREATMENT.—Before proceeding to treat this disea must weigh carefully the symptoms, and be well satisfied as probable nature of the morbid action producing the jau Where we find tumidness and tenderness of the epigastric r extending into that of the liver, with a harsh dry skin, ishness, and a furred tongue, the constitution being sound general blood-letting, followed by leeches and fomentation generally relieve the more acute symptoms, after which doses of calomel, of active purgatives, saline diaphoretics diuretics, and warm baths, will in a few days bring about a so of the disease, the diet consisting of farinaceous articles, and diluents, soda water being about the best drink. Con cence will be matured in such cases by the continuous warm baths, a moderately low diet, and by the daily use of aperients in some bitter infusion. In treating the jaundice returned Indian at home, I have found no remedy in the stage of the disease more serviceable than the warm bath, us bed-time.

In treating the calculous form of jaundice we must come with the use of such means as allay pain and relieve spasmeducts, as opium or morphia in full doses, followed by hot for tations to the affected region, and both measures to be repuntil relief shall be obtained. By such means the violent pain and spasm will generally be overcome, while a hot used to faintness, and followed by a dose of calomel and a saline purge will conduce greatly to the relief of the patient prepare for the stage of convalescence.

Should congestive or inflammatory symptoms superveneral appropriate antiphlogistic means already indicated will be necessary, including nauseating doses of emetic tartar.

When, on the other hand, the patient is anæmic, or long dent in India, with a flat abdomen and evidences of enfe health, neither blood-letting nor mercurials will avail to any but indeed the contrary. We must here be content with a purgatives, bitter tonics, diaphoretics, and the persistent use warm bath at bed-time. Should drowsiness or tendency to appear, very active purgatives perseveringly used, sinapisms to the epigastric region, and blisters to the nape of the neck will prove efficient in aid, even in cases wearing an alarming aspect.

When jaundice becomes chronic I know of no means to be compared in effect to the persevering use of the nitro-muriatic acid bath, applied to the whole body, under the influence of which I have seen cases of the most obdurate nature recover, both in India and at home. As stated elsewhere this powerful agent was first used in the treatment of hepatic diseases by the late Dr. Helenus Scott of the Bombay army, who, according to Mason Good, "plunged the Duke of Wellington in a bath up to the chin, for a severe hepatic affection he was then labouring under, and thus restored him to health in a short time." It was through this means that General Wellesley was enabled to return to his command, and to fight what, in proportion to the numbers engaged, was the most bloody of his victories, the battle of Assaye, and thus to lay the foundation of his future greatness.

CHRONIC ENLARGEMENT AND INDURATION OF THE LIVER.—Although the subject of this article is fully detailed in the second part of this work I am desirous very briefly to notice this form of hepatic disease as it appears in Bengal. It is presented as the result of previous inflammation of the liver, more or less acute, of congestions of the organ, or as the sequel

to fevers, remittent and intermittent.

The liver is enlarged, sometimes indurated, in a perceptible manner, its function is greatly impaired, the biliary secretion being scanty and depraved; that of the kidneys being similarly affected. There is frequently a hacking dry cough, dyspepsia in various forms, a despondent state of the mind, and much general ill health, a generally cachectic state, with a sallow pasty complexion and emaciation; more rarely we find dropsical effusion into the great cavities. In this form of hepatic disease we occasionally find also an embarrassed state of the circulation in the form of excitement of the heart's action, or an intermittent pulse, these symptoms being prominent in proportion as the liver is more or less indurated. In both instances the disturbance would seem to arise principally from the impediment to a free circulation offered by an indurated liver. Whenever, therefore, I hear mention of an excited or intermittent state of the heart's action, I make a careful and searching examination of the hepatic region, the only kind of examination that is of any value.

There is a morbid condition of the stomach and bowels described by the late Dr. James Johnson, which is very often mistaken, both in India and in England, for hepatic enlargement or other diseases; it is a mistake of consequence, inasmuch as the

disorder is one of frequent occurrence, and the treatment usually employed, on the supposition of its hepatic origin, is very injurious to health.

In both cases there is tenderness of the right side and epigastrium, increased by pressure, but here the resemblance ceases. There are, in the affection described by Dr. Johnson, none of the other symptoms characteristic of hepatic disease, and the tumidness, besides the difference in anatomical position, wants both the appearance and solid character of the real hepatic enlargement. Abdominal tumors, such as morbid growths of the omentum, or peritoneum, and feecal accumulations in the colon, are occasional sources of mistake; but the situation and fleeting character of such tumours, together with the occasional and spasmodic nature of the attendant pain will, in general, point to the true nature of the case. In this state of the stomach and bowels there will generally be found a red, inflamed, or excoriated state of the tongue, gums, and fauces, indicating a highly irritable condition of the mucous digestiive surface, the nervous system being much excited. There is occasionally also a continuous acceleration of the pulse, while at other times irregular attacks of feverishness occur, preceded by rigors, and followed by irritable dyspepsy.

Purgatives are borne with great distress, especially when of an acrid nature; and calomel, so often resorted to for the cure of the supposed liver disease, is but rasping the already inflamed or irritated mucous membrane of the stomach and bowels, so as frequently to produce discharges of blood and mucus, with great

aggravation of all the symptoms.

Grievous, then, as are the errors of omission, noticed in the article on hepatitis, where acute disease of the most dangerous nature is allowed to proceed undetected, and fruitful as they are in calamity, they are not surpassed in result by the errors of commission in chronic affections, where mercury, bleeding, and drastic purgatives, are quite as injurious to health, as they are necessary to save life in the first mentioned instance; in fact there are no mistakes more detrimental, nor any against which the inexperienced should be more on his guard, than the two here mentioned. In my experience, however, such errors in diagnosis do not arise so much from want of knowledge as from a careless and hasty manner of examining the patient, an unpardonable negligence in exploration.

The treatment of this disease is not well understood. Mercury I believe to be improper: it injures the stomach and bowels, already but too often over-drugged, without exciting any secretion from the organ chiefly affected, and on which this mineral, from repeated use, has lost its influence; indeed, in the great majority of instances, the system is in a cachectic state, so as to render

taken medicine. This plan was altered, and I was put through a course of the nitric acid bath, taking a vapour bath every other night for the first week. The nitric acid bath acted in a few days very powerfully immediately on using it, and in about three weeks both the liver and spleen could no longer be felt, nor did pressure give me much uneasiness;—the stiffness too disappeared, and my skin became less tense and dry. I took an aperient draught twice a week, and nothing else but the bath. I left Calcutta in the end of December for Simla, and had little or no occasion for medicine for two years afterwards, my general health being completely restored."

This was a case of extreme and urgent danger:—there was enormous enlargement of both liver and spleen, so much so that, on using any drink, if the patient walked across the room the movement caused the fluid to pour from his mouth. There were ascites and universal anasarca; and, altogether, the case was regarded as of a very hopeless nature.

In the cold season of 1838, this same gentlemen again visited Calcutta, in a peculiar state of disease. There was an unnatural dryness or suspended function of the surface of the body and of the mucous intestinal membrane, with a doughy enlargement of the abdomen, swelling of the hands and feet, and obstinate constipation, his excretions from the bowels being deficient in bile, pasty, and peculiarly dry. He brought the following report from the surgeon of his station:—

"Arrived at this station in January, 1836, in tolerable health. During the cold season the digestive organs were much deranged, with loss of appetite, obstinate constipation, distension and hardness of the epigastrium, particularly after meals, acidity, and flatulence. The complexion was sallow and pale, and occasionally he complained of a dull heavy pain in the right hypochondrium. These symptoms continued till the cold season of 1837, when some improvement took place; but on the return of the hot season all the former symptoms recurred with increased violence, accompanied by excruciating headache which for the time utterly incapacitated him from attending to the important duties of his office. Towards the termination of the rains, the hands and feet began to assume a dropsical appearance; and for many months he was compelled to take drastic purgatives almost every day, without which no movement of the bowels could be effected."

Regarding the case now, as one of well marked and extreme torpor of the liver, I directed the patient to go through a course of the nitro-muriatic acid bath, and vapour bath, as before, administering mild saline aperients. Presently the skin began to soften, the tumidness of the abdomen to subside; and with these symptoms, a more ready action of the bowels, so that, in about a month,

a fourth part of his original dose acted as an aperient; and before he left Calcutta for the Cape of Good Hope, the bowels were for the first time during two years moved without purgatives, the discharges having become copious, dark, and pitchy;—in other words the first outpourings of a long pent-up liver. This was a case of peculiar interest from its severity, "calomel, alterative courses of blue pill, combined with purgatives and ipecacuanha, and strong irritating medicines," having, in the language of the surgeon's report, "utterly failed of doing any good."

In the chronic enlargement of the liver, change of air and of climate should never be omitted in the treatment. We have, in the protracted stages of tropical diseases, indeed, no other adequate substitute for change of air. "In India," says Sir James Mc'Grigor, "when patients, whose condition of life permits them to take a voyage to Europe, are in this state of disease, they never fail to take it, and most commonly are recovered by it; but there is no

hope for the poor soldier or sailor."

It is so likewise in the latter stages of fevers and dysenteries; and Dr. James Johnson, speaking of the sufferers from the latter disease, in our fleets and armies, says: "they waste away and die for want of the only remedy that possibly could arrest the hand of

death-change of climate."

The distinction so feelingly pointed to by the eminent men just named, as between the seaman and soldier and his officer, is unjustifiable in principle, as it is cruel and barbarous in results; and I trust the day may not be distant which shall see both classes treated alike in their diseases. On any medical, moral, or just financial grounds, the distinction is, I think, unwarrantable.

As in the articles on fever and dysentery, I conclude the present one with a table exhibiting the geography of hepatic diseases, furnished me by Colonel Tulloch. It shows that though diseases of the liver are somewhat more frequent in the South of India, they are more fatal in Bengal. It is remarkable in the geography of this disease that, though there exists so great an apparent similarity between the climates of the Eastern and Western hemispheres, there should yet be so great a dissimilarity of result in the comparative influence on the European constitution:—in the East Indies generally, Annesley states that thirteen per cent. of the effective strength of British regiments are attacked with hepatic disease, while, according to Colonel Tulloch's Reports, but 2½ per cent. of the effective strength in the Windward and Leeward Command, and one per cent. only in Jamaica, are similarly affected.

	Period	Aggregate	Inflammation of the Liver and Jaundice.			
STATIONS.	of Observation.	Strength.	Attacked.	Died.	Proportion of deaths to admissions.	
Windward and Leeward }	20 years	86661	1946	161	1 in 12	
Jamaica	20 "	51567	539	51	1 in 11	
Gibraltar	19 "	60269	759	22	1 in 34	
Malta	20 "	40826 70293	857 1168	47 58	1 in 18 1 in 20	
Bermudas	20 "	11721	168	6	1 in 38	
Nova Scotia and New }	20 "	46442	384	10	1 in 38%	
Canada	20 "	64280	488	12	1 in 40%	
Western Africa	18 ,,	1843	150	11	1 in 14	
Cape of Good Hope	19 "	22714	496	25	1 in 20	
St. Helena	9 ,,	8973	171	24	1 in 7	
Mauritius	19 "	30515	2508	122	1 in 201	
Ceylon	20 ,,	42978	4382	213	1 in 11	
Tenasserim Provinces	10 "	6818	488	29	1 in 17	
Madras	5 "	31627	3372	190	1 in 17#	
Bengal	5 "	38136 17612	2412 1084	174	1 in 14	
Bombay	5 ,,	1/012	1084	62	1 in 17½	

There died in London, according to the reports of the Registrar General during the fifteen years, 1840—1854 of the specified hepatic diseases as follow, and doubtless a goodly proportion of them had their origin in the influences of tropical climates:—

Hepatitis. Jaundice. Disease of Liver. 2270. 2091. 7914 = Total 12,275.

## ON THE ENDEMIC CONGESTION OF THE SPLEEN.

Considering the frequency and fatality of splenic affections amongst soldiers especially, both European and native, and amongst the European inhabitants of our intertropical possessions, it has always been matter of surprise that so little attention has been bestowed by the profession on the consideration of diseases of the spleen. But when we look to our imperfect knowledge of the anatomy and physiology of this organ, we shall cease to wonder that naval and military surgeons have refrained from disquisitions on what could only result, at the best, but in unprofitable speculation.

Mr. Gray, of St. George's Hospital, the most recent and most successful investigator of the structure and uses of the spleen, considers that the organ "serves to balance alike the quantity and the quality of the blood, and is especially adapted for this function by its connexion with that part of the vascular system which is concerned in the introduction of new material into the circulation."\*

Mr. Gray, from numerous experiments, has come to the conclusion "that the weight of the spleen increases considerably during the time when the digestive process is near to its completion, at the time when the new material is about to be, or has become, converted into blood; and that it decreases considerably in weight at varying periods after that process has been finally completed." This conclusion, says Dr. Carpenter, is confirmed by the results of experiments upon highly fed and upon insufficiently fed animals of the same species; for the increase during digestion was carried in the former to an unusual degree, so as to give the organ more than twice its former weight, whilst in the latter no increase after feeding was observable.

The distinguished reviewer concludes:—"we are inclined to believe that the office of the colourless parenchyma of the spleen is not only to serve as a storehouse for the surplus albumen that finds its way into the circulation on the completion of the digestive process, but also to exert an assimilating action upon it, whereby it is rendered more fit for the nutrition of the tissues; and of this assimilating action, we deem the generation of fibrine to be one of the results. And if it be true, as we have elsewhere suggested, that one special function of red corpuscles is to assimilate or prepare that peculiar combination of materials which is required for the nutrition of the nervo-muscular apparatus, the disintegration of

<sup>\*</sup> Review by Dr. Carpenter, British and Foreign Medical Chirurgical Review, for January, 1855.

these corpuscles in the splenic parenchyma may answer the two purpose of regulating their total proportion in the mass o blood, and of diffusing through the liquor-sanguinis the mate which the nervous and muscular tissues are to draw from i their own development." If such be the functions of the sp we can readily perceive how diseases of that organ may, actually do, spoil the blood, and produce a state of general cachesia.

The tumid or turgid spleen, as it appears in Europeans have resided in low marshy districts in Bengal, or as it c under observation in hospital among the children of ind Europeans, and as seen in British and native soldiers who campaigned in unhealthy countries, appears to consist in pa congestion, or hyperæmia of that organ, and is found in its severe forms during the cold season. It may, and does, occu a primary disease, the result of mere residence in malarious le ties, but far more generally congestion of the spleen is foun-India at least, as a complication of, and a sequel to, fevers r tent and intermittent, and especially in their advnamic type on the West Coast of Africa, Walcheren, parts of Italy, Ar and other malarious countries in the East. When complied with endemic fevers, it may be regarded as in its more stage; but, with the subsidence of the primary disease, this c tion gives place to that of passive congestion. Why the term "engorgement," was preferred by Mr. Twining to the truly expressive and old one, congestion, does not appear certain it is that the altered form of expression has in no added to, or improved, our knowledge of the obscure patholo congestion, whether of the spleen or of any other organ. In the dical, as in other sciences, terms ought not to be changed, u they express relations established as certain, which have not determined before; whereas here, we have but a distinction words without any difference in fact.

In the fevers of the West Indies, Robert Jackson found that spleen was "generally distended, sometimes even to rupture."

Hyperæmia and softening of the spleen are found also t very general complications in the congestive typhoid fevers o own country, and of those of the continent of Europe—the being demonstrated in the most extensive post-mortem exan tions.

SYMPTOMS.—In the more acute form of the disease we present some febrile action—a sense of dull pain in the left he chondrium, extending to the shoulder of that side, accompany fulness and increased uneasiness on pressure—all indiconful something more than mere vascular engorgement—the disease we present action of something more than mere vascular engorgement—the disease we present some february action of something more than mere vascular engorgement—the disease we present some february action of the shoulder of t

ance of the abdominal functions being referrible to congestive fever, and to its splenic complication. The symptoms of splenic diseases, whether acute or subacute, are always more or less negative in character. Whether this form of splenic enlargement be the result of fever, or the more gradual influence of malaria, it is a very important affection, owing to the unfavourable state of the constitution which invariably accompanies it. As a complication with fever, too, it is unfavourable in itself, and as putting an instant bar to the use of mercury, however desirable the exhibition of that mineral might prove, as a means of equalizing the circulation-of eliciting increased secretions-or of relieving local congestions

in other organs.

To the practised eye, the very countenance of the sufferer bespeaks his condition. The expression is dull, the mind anathetic; the complexion is of a dirty lemon-colour, the integument being puffed up and bloated, the eye pale, and of a peculiar clearness. the lips, tongue, and fauces blanched and bloodless; in short, we have here a concentration of cachexia of systematic writers—the very type and essence of anæmia. A wound or trifling abrasion, which at another time would escape notice, now becomes a foul and sloughing ulcer, owing to the depraved state of the blood, and the generally diseased state of the system:-hæmorrhages arise from slight causes-sometimes spontaneously; and so altered is the character of the blood-from want of red globules apparentlythat when performing surgical operations of immediate necessity, at the Native Hospital of Calcutta, I always became aware of the presence of splenic disease on making my first incision-the hue of the blood being demonstrative of the fact.

In addition to the depravation of the fluids, we have excessive muscular debility, bodily inertness, and mental despondencythe whole morbid assemblage tending to impair the functions of respiration, assimilation, and secretion. It can be matter of no surprise, therefore, that when fatal terminations occur, they should be preceded by dropsies, hæmorrhages, or gangrenous ulcers of the cheeks and gums-conditions to which persons suf-

fering from splenic disease are peculiarly liable.

CAUSES.—Of all the known causes of splenic disease, the most influential is malaria. The long continued application of the ordinary endemic influences, without ensuing acute disease of any kind, will, as already observed, produce anæmia, enlargement of the liver, of the spleen, and of the mesenteric glands. But the most ordinary cause of splenic congestion, whether active or passive, will always be found in the malarious fevers of the East-remittent and intermittent-which, for longer or shorter periods, and by the recurrences of their cold or congestive

stages, for months or years together-disturb, or eventually destroy, the balance of the abdominal circulation, and with it the in-

tegrity of the abdominal functions.

When to these morbid conditions we add destitution—the absence of comfort in food and clothing—the residence in low, cold, and damp localities—mental depression—those causes, in short, which contaminate the blood, and determine its flow into the abdominal organs; all these causes will powerfully tend to the production of splenic disease. Previous inflammation and acute congestion, are often found in India to terminate in chronic enlargement of the spleen.

PATHOLOGY.—"Dr. Henoch of Berlin recognises two forms of chronic enlargement of the spleen. The most common form is a true hypertrophy, a mere increase of the natural elements of the organ. In a large number of cases, however, he considers the enlargement is due to excessive insterstitial exudation of plasma, probably of a character unfit for healthy nutrition, and he consequently regards this kind of tumour as the result of chronic splenitis."\*

The appearances on dissection vary from the firm and friable condition, to the indurated and banded texture, approaching to scirrhus—the degree of enlargement varying, according to the intensity of the malarious and other external influences, and to the duration of the disease. The substance of the spleen is often found of a dark purple on being cut into, and composed chiefly of granular matter. Dr. Sieveking found crystals interposed among the granular matter, which were soluble in acetic acid; also yellow spots with tuberculous-like deposits.

In the softened spleen we find its substance reduced to the condition of a grumous mass, indicating a diseased state of the constituents of the blood. Thickening of the investment and adhesions of it to that of the surrounding parts, are more generally

found in the Native than in the European subject.

There is reason to conclude that endocarditis is a frequent and serious concomitant of the more active diseases of the spleen—inflammation of the interior lining membrane of the heart, together with vegetations on the aorta and valves, coagula, and other appearances, demonstrative alike of endocarditis, and of a diseased condition of the blood—being often observable after death. I have no doubt that the numerous sudden deaths of both Europeans and natives of the army of Aracan, in the first Burmese war, during apparent convslescence, arose from the causes just noticed—the fever which destroyed the army having been a malignant, adynamic, intermittent. Sudden deaths from the same causes are found to occur in yellow fever.

<sup>\*</sup> Dr. Ballard, Brit. and For. Review, July, 1855.

TREATMENT.—In splenic congestion of recent occurrence. where the system is sound, and exhibits an inflammatory action. or an acute state of congestion, whether of the abdominal vessels generally, or of those of the spleen, a moderate blood-letting, general or local, repeated according to occasion, and followed by brisk eathartics and sudorifies, will conduce powerfully towards convalescence. On the cure of the fever, indeed, we shall generally find that the splenic complication will subside; but if tumidness of that organ still remain, by the continued exhibition of quinine and brisk purgatives, followed by chalybeates and change of air, congestions, both general and local, will be removed. But, on the other hand, if the type of the fever be adynamic, and the state of the general health anæmic, opposite and tonic remedies must be used; every indication of cure resting on the two radical conditions—the local or endemic influence existing at the time, and the sanitary condition of the individual.

Within a few years of my arrival in India, I suffered severely from the dangerous remittent fever of Gondwana, followed by malignant, adynamic intermittent. For the recovery of my health, I was sent to Calcutta, where, notwithstanding the best advice and care, ague recurred twice or thrice every month during a year and a half, when I was ordered to sea. There was then enlargement of the spleen with ædema of the lower extremities. I made a voyage to sea, and was absent but eight months from Bengal; and, although I used not one grain of medicine, all my ailments disappeared. The truth is, that there are fevers, dysenteries, and chronic forms of hepatic and splenic disease, which no amount of medicine can cure, but which change of air will cure with some

degree of certainty.

When the more acute local symptoms have subsided, and the disease has become a simple chronic enlargement, we still find that no corresponding improvement takes place in the health; and in general, according to the duration of the disease, and the size of the spleen, so do we find a complicated parabysma, and the consequent cachectic state to prevail; softening and relaxation being prevalent in the nervous, vascular, and glandular structures. Again, we occasionally see cases where enormous spleens, occupying two-thirds of the abdominal cavity, cause but surprisingly little disturbance of the health, apparently—the patient complaining only of sense of weight in the left hypochondrium; but this state of passive toleration does not last long, the general health being sure to suffer in the end; while in other cases, and with varying degrees, we have all the unfavourable disorders of the system already described, as concomitants of the more acute stages of splenic disease. When the disease seizes upon the children of indigent Europeans, who are unable to command change of air, or removal to Europe, we find every function in life—the development of the mental faculties—the building up of the corporeal frame—every constituent part stunted and retarded in growth; and without any precocity in youth, there is an actual precocity

of old age.

In the chronic enlargement of the spleen, measures of depletion can but very seldom be required; and even those of a local nature should be used but sparingly. The general practice in Bengal is a steady use of the tonic aperient of Mr Shulbred, formerly surgeon to the Native Hospital of Calcutta, aiding its operation by quinine, chalybeates, stimulant and rubefacient liniments to the affected region, and change of air. The formula of Mr. Shulbred, and the watery imitation of it by Mr. Twining, are as given below:—

MR. SHULBRED.	
Pulveris Jalapæ.	
Rhei.	
— Calumbæ,	
- Scammoniæ.	
Potassæ Bitartrat. aa 3j.	
Ferri Sulphatis, 3ss.	

Mr. Twining.
Pulveris Jalapæ.
Rhei.
——— Calumbæ.
- Zingiberis.
Potassæ Bitartrat. aa 3j.
Ferri Sulphatis, 9j.
Tinct. Sennæ, 3iv.
Aq. Menth. Sativæ, 3x.

Of Mr. Shulbred's powder a sufficiency is given to move the bowels twice or thrice daily. It has for half a century been a favourite remedy in Bengal; and Mr. Twining's formula is given to those who prefer the fluid manner of exhibition. He recommends a chalybeate with purgative extract over night, followed in the morning by powder of myrobalan and the black salt of the bazar;—this as a change of treatment. The kala-nimuk, or black salt of the bazars throughout India, is composed of the following ingredients in one ounce:—

Muriate of Soda		444	grains.
Sulphur		14	"
Muriate of Lime .		12	"
Black Oxide of Iron		6	,,

The natives have long been aware of the beneficial effects acids as well as of chalybeates and bitters in the treatment of the common tumid spleen, for which the bazar doctors prescribe the undiluted sulphuric acid in doses of one drop at a time placed on a bit of plaintain, or more generally five drops in cold water night and morning. Another native plan of treatment is by a mixture of aloes, vinegar, and garlic, with a small portion of the bazar sulphate of iron "kuzees:"—this latter formula is said to be very successful.

I found in treating both Europeans and natives, that severe irritation of the mucous digestive surface was often present, attended by a low irritative form of fever. Here the spleen medicines in ordinary use were altogether inadmissible, as causing increase of irritation, or dangerous diarrhea. In such cases, and they were of frequent occurrence, after subduing intestinal irritation, I had recourse to pure chalybeates, as the saccharated carbonate of iron, the muriated tincture of iron with tincture of iodine-frictions with croton oil-warm baths, &c. In this manner cases having a very unpromising appearance recovered; and the largest spleen I ever saw was cured by the iodide of lead internally administered. It was that of an European boy in poor circumstances. His health had been much reduced by an eruptive fever, when he was seized with ague, for neither of which diseases had he any treatment:-the consequence was enormous enlargement of the spleen, and for which the iodide of lead alone was used, internally and externally. The treatment was continued mildly for four months, when the spleen was no longer to be felt; and during the whole process of cure he never took so much as one aperient. The enlargement in this case had come on very suddenly; but when I first saw the patient it had existed during several months.

In the case of a gentleman from Demerara, treated by me in consultation with Mr. Henry James Johnson, of London, and in whose management we had to contend with extreme cachexia, and sloughing ulceration of the penis, the iodide of lead was persisted in for two months, to the extent, indeed, of producing lead colic, when the spleen, previously of enormous bulk, was reduced to the natural size. Iodide of potassium, warm baths, gestation in the open air, stimulating frictions to the affected region, are all beneficial; but in the chronic enlargement of the spleen, with cachectic states of the system, blisters are apt to slough; and as to leeches and mercurials in such instances, they are certain death.

In Europeans the liver is very often involved in disease along with the spleen; and then there is no remedy, which, in my experience, can at all be compared in power and efficacy to the persevering use of the nitro-muriatic acid bath, using the combined acids internally at the same time with bitter infusions, and keeping the bowels freely open. Here, owing to the hepatic complication, quinine and chalybeates prove ineffective to the cure of the splenic, while they injure the hepatic disease. At page 179 will be seen a remarkable instance of rapid cure, by the acid bath, of enormous disease in both liver and spleen associated with universal dropsy. It was one of the earliest cases treated by me after this manner; and I regard it as one of much interest and importance.

A practical fact of some importance, and which I found to have place in India as in this country, is this—that the acid bath acts more speedily and with far greater power in the hotter than in the colder months. Following out the suggestion afforded by this fact, I always directed the use of a warm water or vapour bath twice or thrice a week at the commencement of the treatment in the winter months, with the view to open and purify the pores of the skin, and to increase its absorbent and exhaling functions.

It is almost unnecessary to urge the necessity of attention to the diet in every stage of this disease. So long as heat of skin, or any febrile movement exists, the diet must be of the lightest and easiest of digestion, being limited to articles of a farinaceous nature; and even in the chronic stage animal food should be used

but once in the twenty-four hours.

Some eminent French physicians have used large doses of quinine for the cure of enlarged spleen, exhibiting the salt to the extent of from forty to eighty grains in the twenty-four hours, the quantity of the drug used being in direct ratio to the extent of the disease. When the stomach did not retain the large doses exhibited, the continental doctors administered the quinine in the form of enemata carefully retained in the bowel. Dr. Elliotson, in his lectures, recommends a similar plan of treatment; but I am not aware that this mode of cure has been anywhere established on the foundations of success. From all I have personally observed and studied I am disposed to look mainly to the emulgent plan by nitro-muriatic acid internally and externally applied, for the cure of splenic diseases, aided and followed by purgatives, quinine, and chalybeates. This important subject is recommended to the careful attention of our intertropical practitioners; recollecting that "some of the greatest improvements in medicine have resulted from researches made in hot climates, and that there is not a single fact observed, or a single disease investigated on the banks of the Ganges or the Mississippi, that does not bring its quota of utility to the practice of medicine in our own country."

The following table but ill represents the prevalence of splenic diseases in any of the military commands specified, such affections being too often "lumped under the head of other diseases." I have been unable to procure any returns for the three Presidencies of India; and in the naval force in the East Indies forming an aggregate strength, from 1830—43, of 40,512 men, only thirty-

nine cases are returned under the head of splenitis.

We may nowhere hope to find any record of the splenic complications of fever and dysentery, although these last form together by far the most prevalent forms of the disease.

STATIONS.	Period of Observation (inclusively.)	Aggregate Strength.	Number admitted into Hospital.	Number Died.	Proportion of deaths to admissions.
Windward and Leeward }	1817—36	86661	104	3	1 in 34%
Jamaica	1817-36	51567	: 59	120	0 in 59
Bahamas	1817-36	535	5	**	0 in 5
Honduras	1822-36	320	_	-	
United Kingdom (Dra-) goon Gds & Dragoons)	1830—36	44611.	5	** :	0 in 5
Gibraltar	1818-36	60269	21		0 in 21
Malta	1817-36	40826	21		1 in 101
Ionian Islands	1817-36	70293	133	4	1 in 331
Bermudas	1817-36	11721	2	11.2	0 in 2
Nova Scotia and New ? Brunswick \$	1817—36	46442	19	7.	0 in 19
Canadas	1817-36	64280	25		0 in 25
Western Africa	1819-36	1843	166	7	1 in 232
1	1818-217	1 0 200	1 1 1		3 3
St. Helena	1836—37 (6 yrs.)	5908	1	***	0 in 1
Cape of Good Hope (Cape District)	1818—36	22714	- 49		0 in 49
Cape of Good Hope }	1822—34	6630	5		0 in 5
Mauritius	1818-36	30515	17		0 in 17
Ceylon	1817-36	42978	50	**	0 in 50

In the very able report of a Committee assembled by order of the Governor-General of India, to ascertain whether the Great Ganges Canal, which now unites the Ganges and the Jumna, would tend to increase disease, and thus prove not a blessing but a curse to the inhabitants, we find a test proposed by Mr. T. E. Dempster, of the medical department of Bengal, which seems to yield positive and satisfactory conclusions, and the value of which appears to have been fully established. Mr. Dempster examined the spleen in a certain number of individuals, and by the presence or absence of an enlargement, he determined whether or not the resident in a certain locality had previously suffered from endemic fever. The following table contains a summary of the whole inquiry.\* It will be observed that in each subdivision of the table, the smallest number of spleens and the corresponding small number of previous fever cases, occur in the inverse ratio of the proximity to the river or the canal.

<sup>\*</sup> British and Foreign Med. Chir. Review for October, 1855, pp. 353, 54.

						Adul	A dulla and children	leens,	200	sunering from lever.	Louis	over.	RITTE	anyface of	
	Delhi branch		Within hall Distant mo	Within half a mile of the canal . { Distant more than half a mile		Jo	of all classes. 58	808. 	184. 51 51		1845 45	1846.	ground	ground, in feet.	46
Irrigated from the Western Jumna	Rohtuk branch		Within half	Within half a mile of the canal Distant more than half a mile		1.1	29	11	34	::	34	27	!!	48	
canals.	Bootana branch		. Distant mo	Distant more than half a mile			16	:	41	:	36	22	1	102	
	Hansi branch	:	Within hall	Within half a mile of the canal Distant more than half a mile		111	39	11	50	4.00	31	22 16	11	92	
	Northern division .	:	Within hall Distant mo	Within half a mile of the canal Distant more than half a mile		!!	20	11	37	.:	39 .	27	!!	138	
Irrigated from the Eastern Junna canals.	Central division		Within hal	Within half a mile of the canal . Distant more than half a mile		!!	59	!!	63	::	53	33	11	14	
	Southern division		Within hal	Within half a mile of the canal Distant more than half a mile		!!	25	11	48	4.00	30	17	11	32	
Irrigated from wells	Irrigated from wells in the high lanh land of the Dooab .	of the D			:		80		37		31	. 20	1	24	
	Sikh states		Connected Unconnected	Connected with the canal Unconnected with the canal	::	-	44 29	!!	43	::	52 .	. 30	11	00	
	Delhi territory		. Unconnecte	Unconnected with the canal			=		32	:	82		:	88	
Unirrigated.	Northern Dooab .		High or bu	High or bungor land* Ganges khadir† Near head of Eastern Junna canal		.!!!	21 21 6	!!!	32 41 35	:::	30 42	13 28 27	111	46 0 0	
	Naturally malarious localities (Valleys of Junna and Hindun	ocalities	S Valleys of	jheelst Jumna and Hindun		-	34	11	45	::	683	. 31	11	15	
	Naturally malarious l	ocalities	Nujufgurh	jheelst Jumna and Hin	dur			!!	44	44	46 46	44	34 46 42		34 46 42 31

\* Bungor : The high and firm bank of the river bounding the khadir.

\* Khadir is the balt of moist, low, and often fertile land found alternately on one or other side of large rivers in these provinces. During the rainy season much of this land is advantable to the river frequently and in the river frequently and in the control frequently was a subject to the results of the results and the results are requested to the results of the results and the results of the res

## EPIDEMIC CHOLERA.

SKETCH OF THE EARLIER INDIAN HISTORY OF EPIDEMIC CHOLERA.—From the contemporaneous writings of Hippocrates and the Chinese physicians, as well as from those of the later Hindu and Roman medical authors, it is evident that epidemic cholera was well known and accurately described in

their respective times.

The first notice of this disease we have, by an European writer, is that of Bontius, physician to the Dutch East India Company, and who wrote in 1629, in Batavia. "The cholera morbus," he says, "is extremely frequent: in the cholera, hot bilious matter irritating the stomach and intestines, is incessantly and copiously discharged by the mouth and anus. It is a disorder of the most acute kind, and therefore requires immediate application." In its severer form "the animal spirits" are described as speedily "exhausted, and the heart, the fountain of heat and life, is overwhelmed with putrid effluvia; those who are seized with this disorder generally die, and that so quickly, as in the space of four-and-twenty hours at most. This disease is attended with a weak pulse, difficult respiration, and coldness of the extreme parts: to which are joined great internal heat, insatiable thirst, perpetual watching, and restless and incessant tossing of the body. If, together with these symptoms, a cold and fetid sweat should break forth, it is certain that death is at hand." Bontius treats of the "cholera morbus" and "spasm" as separate diseases; yet it is presumed that both were frequently united, and that, in reality the spasm was a common complication with cholera, as indeed he describes it in the case of Cornelius Van Royen, who "was suddenly seized with the cholera, about six in the evening, and expired in terrible agony and convulsions before twelve o'clock at night."

I shall now lay before the reader, in as condensed a form as possible, the results of inquiry instituted by order of the Governments and medical authorities of Bengal, Madras, and Bombay, with a view to obtain the best possible information as to the his-

tory, nature, and cure, of epidemic cholera.

How ably the medical services of the three Presidencies responded to this call, need not now be pointed out; neither is it

necessary to dwell on the original merits of the author of the "Essay on the Influence of Tropical Climates on European Constitutions," further than by stating, on the authority of Dr. Copland, that "the nature and treatment of the disease were very imperfectly known, until Dr. James Johnson described its symptoms, and pointed out a more successful method of cure, than had previously been employed."

In the reports of the medical boards of Bengal, Madras, and

Bombay, we find the following notices:-

#### BENGAL REPORT.

1781. In March of this year a disease, in all respects resembling the recent epidemic cholera, seized on the Bengal force under Colonel Pearse of the Artillery, while marching through the Northern Circars to join Sir Eyre Coote's army. Seven hundred men died within the first few days, and three hundred convalescents were left behind; this out of an original force of five thousand men.

"Men in perfect health dropt down by dozens: and those even less severely affected were generally dead, or past recovery, within less than an hour. Spasms of the extremities and trunk were dreadful; and distressing vomiting and purging were present in all."

1783. Cholera is said to have broken out at the sacred bathing spot of Hurdwar on the Ganges, in April of this year.

It happened to be one of the twelfth years deemed peculiarly propitious by the Hindus, and the assemblage of pilgrims was consequently far beyond the common annual average, amounting, it is said, this year, to nearly two millions.

"The disease broke out on the springing up of an easterly land-wind during the night, and carried off innumerable persons." In less than eight days, twenty thousand victims are said to have fallen; yet so confined was its influence, that it did not extend beyond the place of bathing, and ceased on the dispersion of the multiple of the multiple of the dispersion of the dispe

## MADRAS REPORT.

1774. The "true cholera morbus" is described by Dr. Paisley, of Madras, as being "often epidemic" amongst the natives, whom it destroyed quickly. In the first campaigns, it proved very fatal to both the European and native troops. He describes it as "operating on the system as a poison, and brings on a sudden prostration of strength, and spasm over the whole surface of the body." The pulse he found to "sink suddenly, and bring on immediate danger."

Sonnerat speaks of "an epidemic disorder which reigns" on the Coromandel coast, and his observations embrace the

period from 1774 to 1781.

After describing the symptoms, and that the poorer and more negligent classes are most obnoxious to its seizure, he states that, "above sixty thousand people from Cherigan to Pondicherry perished." "The Indian physicians," he says, "could not save a single person."

One epidemic is described as peculiarly severe. "Those who were attacked had thirty evacuations in five or six hours, which reduced them to such a state of weakness that they could neither speak

nor move

"They were often without pulse; the hands and ears were cold; the face lengthened; the sinking of the cavity of the socket of the eye was the sign of death; they felt neither pain in the stomach, cholics, nor gripings. The greatest pain was a burning thirst."

1787. A disease is reported as prevalent this year at Arcot, "similar to an epidemic that raged amongst the natives about Palicola, and in the army of observation in January of 1783, in the Bengal detachment at Ganjam in 1781, and several other places at different times.

The characteristic marks were spasms of the precordia, and sudden prostration of strength.

## BENGAL REPORT.

1790. In the middle of April of this year, a fatal cholera of the spasmodic form attacked the detachment under Colonel Cockerell, marching from Bengal to Seringapatam, in the same country in which it had previously proved so fatal to the force under Colonel Pearse—the Northern Circars.

The above three notices, two of which refer to seizures beyond the bounds of the Bengal Presidency, are all of which any record could be obtained by the talented author of the Calcutta Report. Rumour, indeed, had spoken of severe cholera in Bundlekund some forty years previously to the date of Mr. Jameson's report, or about the year 1780, and of another some time near the end of the last century; but as no authentic history exists respecting their nature, no attempt is made to trace them.

The Bombay Report enters into no history of epidemic cholera, beyond a casual reference to a letter from Dr. Taylor, who gives an account of a disease, taken from a Sanscrit work, "which leaves very little doubt that it has not only been long known to the natives, but proves its identity."

#### MADRAS REPORT.

The treatment consisted of stimulants and hot baths, supporting the vis-vitæ."

1782—90. During this period, cholera under various names, but always with the same symptoms and results, would seem to have prevailed in various parts of India.

"The mort de chien, or cramp," prevailed in Sir Ed. Hughes's flect in 1782. The sufferers were "soon brought to great weakness, coldness of the extremities, and a remarkable paleness, sinking and lividity of the whole countenance." There was "great thirst, or rather a strong desire for cold drinks, but there was no headache, or affection of the sensorium commune throughout."

From July to September 1782, Mr. Curtis describes the same disease in the Madras Hospital, and in the fleet.

1782. Mr. Girdlestone describes a disease exactly similar, as destructive to the newly arrived troops from England, more than 50 of whom were carried off by it "within the first three days after they were landed."

1787. Dr. Duffins gives details of the symptoms and treatment of "cholera morbus," which then prevailed at Vellore with great violence—a disease "so rapid in its progress, that many of the men are carried off in twelve hours' illness."

The same epidemic prevailed fatally at Arcot, to which station Mr. Davis, member of the Hospital board, was deputed to investigate its nature. "He understood from the Regimental Surgeons that the last disease had proved fatal to all who had been attacked by it."

Mr. Thompson gives a similar account to that of Mr. Davis's, both having remarked on the singular contraction and emptiness of the bladder on dissection, it being found no larger than "a walnut."

1814. Two notices of cholera are found in this year—one by Mr. Wyllie of the 24th Regt., and the other by Mr. Cruickshanks of the 9th Regt., in which it appeared with great severity.

"The disease in the 9th Regt., in 1814,

BENGAL REPORT.

#### MADRAS REPORT.

resembled in every particular, with exception to the heat at the pracordia, the cholera at present" (1819) "so common, although it could not be called epidemic." It appears to have broken out in a brigade of N. I. while on the march from Jaulnah, about the 10th of June. The patients "exhibited all the symptoms now so well known of persons labouring under the advanced and fatal stage of the epidemic cholera, the skin cold, and covered with a cold perspiration, the extremities shrivelled, cold, and damp, the eyes sunk, fixed, and glassy, and the pulse not to be felt. These persons all died, and I find on referring to such notes as I have preserved, that, influenced by consideration of the vascular collapse, and total absence of arterial pulsation, I had denominated the disease, asphyxia."

The report of Mr. Cruickshanks is peculiarly valuable on many grounds, as well as for an early notice of one of those singular and unaccountable features which it has frequently manifested since, namely, that of the two corps composing the brigade, apparently under similar circumstances, one only was attacked, while the other escaped altogether.

Two distant and separate notices of the epidemic cholera, one 25 years, and the other 34, previously to 1817, are given by Mr. Staff-surgeon Hay. These visitations were in the Travancore country.

Dr. Burke, also, in a report from the Mauritius in 1819, where he was chief of the medical staff, states that a disease in all respects similar to the recent epidemic is reported to have occurred in that island in the year 1775. "The symptoms, fatal and sudden effects, and duration of the disease would seem to be the same."

It thus appears clearly, that epidemic cholera prevailed at various remote periods, and at many of the principal stations throughout British India, sometimes coming as a wide-spread pestilence, and at others, desolating only particular localities.

That the disease, in its epidemic form, was not altogether unknown to our forefathers in Europe, is rendered more than probable also, by the writings of various of the older physicians; and many of the cases described by Sydenham, in 1669, would seem to have been of the true spasmodic nature; but, as none of the visitations described by former writers approached in duration, extent, or severity that which took its rise in Bengal in 1817, no detailed record of them is preserved. From the date of its

first outbreak on the shores of the Ganges, till in its course it embraced those of the Indus, Euphrates, Nile, Danube, Volga, St. Lawrence, and Mississippi, this great epidemic has visited every nation of the earth; and whether it fastened on the delicate frame of the Hindu, or that of the robust European, the disease has been essentially of the same nature, nor have the results of medical treatment differed much. Everywhere throughout Asia, Africa, Europe, and America, it has been thus, no difference of race, climate, or temperature exercising any very marked influence on its nature and progress.

Condition of life, as in all other epidemics, has everywhere exercised its vast influence for good or evil; amongst the poorer classes, in their greater liability to this deadly disease, and amongst the wealthier classes, in their comparative exemption

from disease and death.

In each and all the countries traversed by this scourge, certain low, damp situations, filth, destitution, and misery, have exercised their usually fatal power; while the more comfortable conditions

of life acted as powerful preventatives.

It was amongst the poor, ill-fed, ill-clothed, and crowded inhabitants of Jessore that epidemic cholera made its first appearance; and it is under similar circumstances, and in similar localities, that the disease has ever since, and in every country, proved

most rife, most virulent, and remained the longest.

The cholera epidemics which have ravaged various parts of Hindustan since 1817, have always originated in, and issued forth from India. They have never, in any instance within my knowledge, been even supposed to have been imported into any of the ports of India, by ships from infected countries, or through any other manner of human intercourse. It may therefore be inferred, that the cause of the disease, however latent or submerged for a time, is never actually absent from the soil of India, or from some of its localities.

Without disputing the possibility of contagion in cholera, even in the midst of and notwithstanding the thorough ventilations prevalent throughout India, I am bound to say that although I may be said to have lived in the midst of the disease during many years, I have never seen anything which, in my opinion, warranted the belief; nor have I ever communicated with any Indian medical officer who believed in the contagiousness of epidemic cholera in India.

In the European General Hospital of Calcutta, in which I served as assistant-surgeon and surgeon, it was well known that of the five native keepers and washers of clothes who had, during twenty-five years, kept and washed the hospital clothing, not one had cholera; nor had those who assisted them. The same immunity attended the native dressers, averaging from twenty to thirty

men, who, during the same number of years, were in constant and close attendance on the cholera sick, all day and all night; nor were the sweepers who washed and dressed the patients, and who removed the matters vomited and ejected by stool, ever affected with cholera.

I served in the General Hospital in March, 1827, the time referred to by Mr. Twining, when the house was filled with cholera patients, and when all of us, European as well as natives, including native medical students employed for the occasion, were exhausted with the labours of attendance on the sick; but none of us suffered from the disease. To these facts many others of the like nature might be added; but, in respect of contagiousness, who shall dispute the possibility of occasional exceptions, under peculiar susceptibilities, and under different circumstances of climate and other contingencies.

GEOGRAPHICAL PROGRESS OF EPIDEMIC CHO-LERA.\*—The epidemic cholera, which originated in the delta of the Ganges, in July, 1817, true to its birth-place the town of Jessore, with its low, marshy, jungly, crowded, and ill-ventilated conditions, spread rapidly from villages, towns, and cities, in no better sanitary state than their prototype; and from that day to the present, epidemic cholera has proved greatly more fatal in Asia than in Europe.

> It commenced its march from Bengal in August, 1817, Reached Bombay . . . 10th August, 1818. And Astrakan, where it died away, 18th September, 1823.

In Persia several local epidemics appeared at intervals in the seven years, 1823—30. The great epidemic that traversed Europe is said to have sprung up in June, 1830, on the low western shores of the Caspian Sea. The velocity of the two epidemics only differed in Asia; it was the same in Europe.

The epidemics of 1830, 31, and of 1848, 49, travelled at the same rates westward, as appears from the following table:—

Astrakan			July 20th, 1830	June .		1847
Moscow			September, 1830	September 18	3th	1847
Petersburgh			June 16th, 1831	June .		1848
Berlin .			August 31st, 1831			1848
Hamburgh			. October, 1831			1848
Sunderland			October 24th, 1831	October 4th		1848
Edinburgh		1	January 22nd, 1832	October 1st		1848

The progress of the disease did not stop here; it ravaged the nations again and again, its course, like that of the epizooties, blights, and mildews, being altogether unknown. Infection has

<sup>\*</sup> It will be seen that, in the body of the following article, I have made free use of, and copied largely from the admirable works of Dr. Farr, of the Registrar-general's office, and from those of Drs. Baly and Gull, framed by the desire of the College of Physicians.

not been proved to have had any share in its progress and propagation; on the contrary, it has marched onwards through an inscrutable atmospheric and terrestrial agency, uncontrolled by all opposing physical obstacles, and by all the efforts of man. Nor does there exist any means of modifying its force, other than such as are well known to lessen the power and duration of all epidemics, viz., the removal of all those conditions, whether natural or artificial, that produce and maintain impurity and stagnation in the atmosphere we breathe and in the water we drink, conditions that everywhere, and in all times, have opened the human frame to the admission of disease, endemic and epidemic.

SYMPTOMS AND PROGRESS.—There are two circumstances which render the assault of true cholera both insidious and dangerous in the extreme:—it is preceded by a painless diarrhæa of more or less violence and duration, according as the epidemic may be at its height or decline; and its seizure is generally by night, or in the early morning. It thus happens that, too often, the real nature of the case is mistaken by the patient for some

slight disorder of the bowels.

When once the true cholera has fastened on the patient it proves the most unmistakeable of diseases—one case being the counterpart of all others-so that an error in diagnosis is here scarcely possible. The disease in its most malignant form is the same in all countries. It is ushered in by muscular debility, tremors, and vertigo, occasional nausea and spasmodic griping pains in the bowels, and marked depression of the functions of respiration and circulation, and sense of faintness. These signs are speedily followed by copious purging of serous fluid, and this again by vomiting, with sense of burning heat at stomach-coldness and sweating-dampness of the whole surface of the bodycoldness and lividness of the lips and tongue-cold breath-a craving thirst-a feeble rapid pulse-oppressed and difficult respiration-suppressed urinary excretion-extreme restlessnessblueness of the entire surface of the body-a sunken and appalling countenance—a sunken and peculiarly suppressed voice a peculiar and indescribable odour from the body-partial heats of the præcordia and forehead; -such are the signs of a fatal collapse.

Such is the ordinary course of this fatal disease in its most virulent form, or at the onset of its epidemic invasion. But in less concentrated visitations, or towards their decline, there will be found a premonitory diarrhœa, varying in duration from three to twelve hours, or more, to be followed by true cholera. During the stage of diarrhœa the quantity of serous fluid voided is some-

times immense, and in one case noted by Dr. Parkes, in which the patient had but seven stools, eighty-eight ounces of fluid were

discharged from the bowel.

But the relation between purging, vomiting, and cramps, and the final algide collapse, is exceedingly variable. In some cases the patient will have been purging for twenty-four, forty-eight, or more hours, before vomiting, muscular cramps, and collapse occur; while in other instances the whole duration of the disease will have been gone through unto death, in seven or eight hours, with marked collapse almost from the commencement of the seizure.

A Hindu, attended by Dr. Parkes, lived 108 hours from the time of the first stool, and ninety-one hours after the positive development of algide symptoms—the patient having voided altogether twenty-one stools. Another Hindu, in contrast with the above, and who had only four stools, died in ten hours from the first discharge. All observation in India goes to show that the more concentrated and malignant the disease the less purging and vomiting; and so also in the worst and most rapidly fatal cases, muscular cramps are very rarely observed as prominent symptoms.

After an indefinite duration, and varying infinitely in the degrees of severity of the several symptoms, we find, in favourable cases, that the oppressed functions of respiration and circulation begin to show signs of more freedom of action;—feverish reaction commences, to be developed into actual and sometimes severe fever, accompanied by headache and gastric symptoms—the results of previous congestions and arrested functions of secretion.

PREDISPOSING CAUSES.—Although it be quite true that the cholera poison requires no definite predisposing conditions in the system to enable it to produce its effects, and that it proves fatal at all ages, and nearly equally so to both sexes, and that neither the weakness of infancy, the vigour of manhood, nor the decrepitude of old age, are a safeguard against its inroads;still, in common with fever and dysentery, it may be truly said that whatever depresses the constitutional powers will predispose to the invasion of cholera; as the use of crude, indigestible, or ill-prepared food-fatigue, want, filth, and misery,-but, above all, intemperance in the use of spirituous liquors, so common to the lower orders of English in all parts of the world. It was found by Dr. Farr that on Saturday, Monday, Tuesday, and Wednesday, the deaths from cholera in London and other cities were above, and on Thursday, Friday, and Sunday below the average. The week's wages are generally paid on Saturdays; and the Mondays are days on which a certain proportion of the population indulge in intoxicating drinks—the Fridays being days of comparative abstinence.

It thus appears that the depression of the nervous system and the derangement of the digestive organs following immediately upon occasional intoxication, had a marked predisposing influence. But it is difficult to estimate separately the various noxious influences which in large cities are naturally associated with intemperance. This vice soon entails all those conditions which not only impair the vigour of the body, but favour the dissemination, and probably the generation of the cholera poison. There is, indeed, sufficient evidence that not only habitual drunkenness in itself, but also the immediate effects of an occasional carouse, will induce a predisposition to an attack.

The influence of long marches in predisposing to cholera in India, would appear to indicate something more powerful than mere physical depression resulting from the labour of marching; for, in the laborious trades and occupations of civil life, the mortality from cholera in England has not been proved to have been necessarily high, or to have been in any manner associated

with excessive labour.

It has been proved in India, and recorded by Mr. Orton, Drs. Balfour and Lorimer, that troops, both European and native, suffer more from cholera on the march, or soon after it, than they do at the military stations. Dr. Balfour has proved, that of the native soldiers of the Madras army 32 died of cholera in cantonment. and 86 when marching, to an average of 10,000 of strength. The attacks were respectively 85 and 200 in 10,000. Dr. Lorimer's reports show that the troops were more frequently attacked on long than on short marches:-thus the troops in 219 marches of 20-40 days were attacked 39 times; while in 14 marches of 100-120 days they were attacked seven times. If we take a hundred marches as the basis, they were attacked 18 times in about 30 days, in the one case; while in the other case they were attacked 50 times in about 110 days-that is, at the rate of 14 times in 30 days. This is no proof, says Dr. Farr, that fatigue increases the liability to attack; -it only proves that, on the long marches, the men are exposed a longer time to the causes of disease. We would assuredly expect more men to be wounded in a battle of three days', than in a battle of one day's duration. When a march is conducted with undue haste, in heavy order, or under direct solar exposure, in hot climates, it will produce fatigue and other injurious results :- but in general a well-ordered march is conducive to health; and there are well-known instances in which the health of an army has even improved materially on the line of march.

Soldiers are exposed to many injurious influences during long marches, which frequently lie by rivers, on low marshes and jungly grounds; and when to these circumstances we add sleeping on the ground, we perceive many causes other than fatigue in operation. As elsewhere observed, camp beds, which should raise the men off the ground, would prove a great advantage; and so

would a light order of march, and short marches.

But referring to the special subject of this section, I believe that we are seldom so much in error as when we would ascribe sickness and mortality to one cause only; while in reality disease and death are always, and under all circumstances, justly to be referred to the assemblage of many causes. In truth, the medical topographer, when investigating the difficult and obscure causes of disease, must ever hold in recollection the requirement of Cabanis—to inquire into all the circumstances natural and physical in the midst of which we live in a given place.

COMPARATIVE DURATION OF FATAL DIARRHŒA, SUMMER CHOLERA, AND EPIDEMIC CHOLERA.—The average duration of 4,045 fatal cases of diarrhœa in males, in the epidemic of 1848, was 16.04 days—of 3,851 females 16.69 days. The summer cholera, according to Dr. Farr, has an intermediate duration of about five days. The difference he exhibits in the subjoined table of cases taken indiscriminately from the returns of 1848: more than half the cases of Asiatic cholera terminated in less than a day; half the cases of summer cholera did not terminate in three days; and half the cases of diarrhœa lasted more than six days. This is a new and important element in diagnosis.

DURATION OF FATAL CASES OF DIARRHŒA, SUMMER CHOLERA, AND ASIATIC CHOLERA, IN LONDON, 1848.

-	Total	Under		Numbe	r of Cas	es terr	ninating	in the s	everal D	ays.
	Deaths.	Day.	1	2	3	4	5	6	7	8
Diarrhœa Summer Cholera Asiatic Cholera .	116 116 129	1 9 66	1 12 21	8 14 12	17		9 5 3 11 2 6	6	13 7 5	8 4
		Ç.,	Nu	mber of	Cases t	ermina	ting in t	he seve	ral Days.	-
		9	10	11	12	13	14-21	21—28	28—35	35 & up.
Diarrhœa Summer Cholera Asiatic Cholera .	116 116 129	1 4	5 1		2	ï	12 2 	13 1	7 2 1	8

Fatal diarrhoa is thus seen to be a disease of much longer duration; and it agrees with cholera, being of somewhat longer duration in females than in males in both cases, between the age of 15—35, than at ages further advanced.

DURATION OF FATAL CASES OF DIARRHŒA AT VARIOUS AGES.

						Days.
Males, all ages		- 1			7	16.044
Females, all ages		*				16.692
Males and Femal	es,	age 15	-35			18.668
Males and Femal	es, a	ge 35	and	upwa	rds	17.544

The duration of life, after well marked symptoms of fatal cholera have set in, appears to diminish as age advances. Dividing these cases into three groups, the following results are obtained:—

DURATION OF FATAL CASES OF CHOLERA AT VARIOUS AGES.

	Mean	Duration.
	Days.	Hours.
Persons of the age of 15-35	2.121 =	= 50.904
Persons of the age of 35-55.	1.954 =	= 46.896
Persons of the age of 55 and upwards	1.973 =	= 47.352

The fatal cases of cholera were preceded frequently, if not invariably, by a stage of diarrhoa, which attracted little or no attention. That stage is, it is believed, not included in the cholera duration, which is reckoned by the medical attendants who made the returns, from the manifestation of the first characteristic symptoms of the epidemic.

Of 19,232 fatal cases of cholera in males, 315 terminated in six hours, leaving 18,917 alive; 3,030 died in the next six hours, leaving 15,887 alive at the end of the twelfth hour; 4,965 died in twelve and less than eighteen hours, leaving 10,922, or rather more than half the number of those who die, alive at the end of eighteen hours. If a man is to die of epidemic cholera, it is probable that he will die in the first day (twenty-four hours), as out of 19,232 males who die, 10,187 die within, and 9,045 survive that period.

RELATION OF ELEVATION OF THE SOIL WITH MORTALITY FROM CHOLERA IN LONDON.—Dr. Farr states that the elevation of the soil in London had a more constant relation with the mortality from cholera than any other


known element; the mortality having been in the inverse r the elevation. The mortality of the nineteen highest d was at the rate of thirty-three in 10,000, and at the nineteen districts 100 in 10,000. The elevation in the two groups seventy-one to ten feet above the high-water mark of the TI or as seven to one; while the mortality was as one to three the inverse ratio.

In the two groups of the six districts, supplied with the of the Thames at Kew and Hammersmith, the mean elevation 35 and 175 feet, the mortality from choleran intereen and in 10,000. In the two groups of twelve districts, supplied Thames water between Battersea and Waterloo Bridges, the elevations were one-eighth foot and ten feet; the mortality and 77 in 10,000. In the two groups of twenty districts, supplied with the waters of the New River and the Lea, the mean elewas 24 and 59½ feet; the mortality from cholera was 59 at While the effects of the water and of the wealth of the diare apparent, we do not, in this analysis, conceal the effected elevation.

LONDON. GROUP OF DISTRICTS.		Population (Persons to	Elevation in Feet above luco High Water (Yea Mark, (Trinity).
6 Districts supplied with Water taken from the Thames above Battersea	15	72	105
20 Districts supplied with Water from the New River, the Lea, and the Ravensbourne	48	137	42
12 Districts supplied with Water taken from the Thames, between Battersea and Waterloo Bridges .	123	73	5

Cholera was excessively fatal in all the four districts whi on a level with, or below the Trinity high-water mark; it dest 144, 161, 164, and 205, in 10,000 inhabitants. The foll diagram and description are from Dr. Farr's report:—

# A DIAGRAM.



"The figures in the centre express the number of deaths from cholera to 10,000 inhabitants living at the elevations expressed in feet at the side of the diagram.

The length of the black horizontal lines shows the calculated relative fatality of cholera in districts, at relative elevations indicated by the height from the base of the diagram. The dotted lines indicate the mean mortality observed in the elevations given. Thus:—in districts at 90 feet above the Thames, the average mortality from cholera was 22 in 10,000 inhabitants.

The relation discovered between the elevation of the soil, and the mortality from cholera, is so important that it was thought right after the above calculations were made to submit the principle to another test, by comparing the elevation and the mortality from cholera of each *sub-district*. The population of the sub-districts, in 1851, having been enumerated, it became possible to construct the tables which, although they make the mortality no the lowest level less, and are deranged by the deaths in hospitals and workhouses, entirely confirm the announced law.

We must look with extreme interest to the confirmation, within our intertropical possessions, of the law here deduced by Dr. Farr; the question being one of extreme importance as regards the welfare both of the native inhabitants, and of the European troops. The low-lying towns on the coast, says Dr. Farr, were all attacked by cholera; and the disease proved three times more fatal on the coast than in the interior of the country. Cholera also appeared first in the ports of England, and gradually extended into the interior.

The epidemic is stated by this authority "not to have been so fatal on the primary geological formations—on the granite, the Silurian, or the Devonian systems; while Herefordshire, however, on the old red sandstone, escaped, Cornwall and South Devon, on the same formation, suffered severely. Plymouth, and the districts around the Sound into which the Tamar flows, were one of the principal cholera fields; but it is a port, and the mouth of a river which rises partially in the millstone grit, forming a part of the carboniferous system."

THE SIX THEORIES AS TO THE CAUSE AND PRO-PAGATION OF EPIDEMIC CHOLERA.—The varieties of opinion as to the remote cause of Asiatic Cholera, according to Dr. Baly, and the causes of its spreading are very numerous, but all that need be referred to may be reduced to six principal theories which must be briefly stated, since it is only by reference to them that the value and import of the facts to be examined can be estimated.

"1. The first theory is, that the disease spreads by an 'atmospheric influence or epidemic constitution,' its progress consisting of a succession of local outbreaks, and that the particular localities affected are determined by certain 'localising conditions,' which are, first, all those well-known circumstances which render places insalubrious; and, second, a susceptibility of the disease in the inhabitants of such places, produced by the habitual respiration of an impure atmosphere.

"2. The second theory, following the analogy known to be due to morbid poisons, regards the cause of cholera as a morbific matter which undergoes increase only within the human body, and is propagated by means of emanations from the bodies of the

sick, in other words, by contagion.

"3. The third theory—that propounded by Dr. Snow—gives a more specific form to the doctrine of contagion. It supposes that the poison of cholera is swallowed, and acts directly on the mucous membrane of the intestines, is at the same time reproduced in the intestinal canal, and passes out, much increased, with the discharges; and that these discharges afterwards, in various ways, but chiefly by becoming mixed with the drinking water in rivers and wells, reach the alimentary canals of other persons, and produce the like disease in them.

"4. The fourth theory assumes that the cause of cholera is a morbific matter or poison, but supposes that it is reproduced only in the air, not within the bodies of those whom it affects, and that

its diffusion is due to the agency of the atmosphere.

"5. The fifth theory is a modification of the fourth. It admits that the cholera matter is increased by a species of fermentation, or other mode of reproduction in impure, damp, and stagnant air, but maintains that it, nevertheless, is distributed and diffused by means of human intercourse; it being carried in ships and other vehicles, and even in the clothes of men, especially the foul clothes of vagrants, and the accumulated baggage of armies.

"6. The sixth theory combines the second and fourth, assuming that the material causes of the disease may be increased and propagated in and by impure air, as well as in and by the human

body."

Such are the theories; and the object of Dr. Baly's Report to the College of Physicians is to inquire into the facts, and to learn which of these theories is most in accordance with them. The conclusions arrived at are as follow:—

1. The theory that the cause of the disease is a general state of the atmosphere—a general "atmospheric influence," or "epidemic constitution" has been found untenable. The wide extent which the epidemic soon attained in England in 1848, the obvious dependence of its increase in the summer of 1849, and of its decrease and subsequent disappearance in the autumn and winter of that year, on some influence exerted throughout the country, and the association of the disease, when it was very rife, with those characters of site and defects of sanitary provisions, which have been termed "localising conditions,"—these three general facts constitute the sole support of the theory in question. Other facts, as general and as well established, are irreconcilable with it. The duration of the epidemic for many months in a country, and even in a single city, is obviously inconsistent with the belief that the cause is an influence belonging to the moving body of the atmo-

sphere; and if the atmospheric influence is regarded as independent of the moving air, though acting through it-some "electrical state" or "telluric influence"—the following features of the epidemic are still difficulties little less than insuperable in the way of adopting this theory :- first, the very partial distribution of the epidemic. even among places presenting equally the so-called "localising conditions;"-second, the outbreak of the disease at different times, in several parts of a comparatively limited tract of country. in several parts of the same town, or even of the same building ;third, the cessation of the disease, also at different times in different localities; -fourth, its lingering in a few spots where the general conditions favouring it have obviously ceased :- fifth, its extension from centres in different parts of the country ;- and. sixth, the evident connection in many cases subsisting between the successive outbreaks in neighbouring places, and especially in different parts of a town or of a public establishment, where the occurrence of the first outbreak of the series has evidently been the condition determining those which followed. Still more difficult it must be to explain satisfactorily by this theory, in any form which might be given to it, the comparatively slow and stepby-step progress of the epidemic in a determinate course over a continent; its slower progress in winter than in summer; its frequent arrest in the winter; its often affecting a mere narrow tract of country, leaving unscathed the regions on either side; its extension through a country in different directions at the same time : its march for months together against the direction of the winds: the rapidity with which it has crossed the Atlantic Ocean, contrasted in a striking manner with its slow progress on land: the constancy with which, in invading an island or fresh continent, it has appeared at the largest seaports before extending inland; the great leaps it sometimes seems to make over entire countries in passing from one seaport to another; and, lastly, many facts to be noticed presently, which clearly connect its extension with the intercourse and locomotion of mankind.

2. The persistence of the epidemic for a certain time, even in localities of small extent, and its very partial distribution in a country, a town, and even parts of towns, are two facts which at once suggest the inference that the cause of the disease is a material substance, and that it is only partially distributed. This inference is confirmed by the characters above referred to, as those presented by the epidemic in its progress over a country, and by the fact that, within a limited area, many spots have remained free from it, which exactly resembled the localities attacked in respect of the supposed localising conditions. It is certain, however, that cholera is so far connected with the conditions of low site and defective sanitary provisions, that it is

never very rife, except where they are present in a marked degree. Now, all these conditions alike tend to produce a damp and impure state of the air in the places where they exist; while the variations of the intensity of the epidemic, attending variations of season and temperature, and the slighter effects of other known meteorological changes, are most easily explicable by referring them to the increase or decrease of moisture and impurities in the air, which necessary result from a rise or fall of temperature. The natural inference is, therefore, that the matter, which is the cause of cholera, increases and finds the conditions for its action, under the influence of foul or damp air with the aid of some degree of warmth; and this being premised, other facts become intelligible, namely, the persistence of the disease in the winter, for the most part in the interior of large establishments, where there is warmth, together with impurity of the air produced by the accumulation of many human beings within a limited space; the preference manifested by the disease throughout an epidemic for low and denselypopulated districts, especially the tracts of countries about the mouths of rivers, and for crowded towns and dirty ill-ventilated parts of towns, or single dwellings of the same character even in elevated situations; and likewise its appearing at the commencement of an epidemic, as a general rule, first in places of the character described, since in these places the impure and damp air would be found earliest and in greatest abundance.

The increase of the epidemic at its climax results chiefly from the increased number of individual localities then affected; and since this increase in the number of separate outbreaks takes place, as a general rule, most rapidly at that season when a high temperature, and other meteorological conditions, tend to increase the impurity of the air, and in those districts where local conditions affect the state of the air in the same way, it may be inferred that, while the increase and action of the cause of the disease depend on the state of the air in small localities, its transmission from spot to spot is facilitated by the state of the atmosphere over larger areas. This view is, of course, corroborated by the more rapid extension of the epidemic over the continent of Europe in the summer season, by its extension, especially along low tracts of country bordering on rivers, while it avoids mountainous regions; and by the fact that in this country it was very late in attacking some districts lying far inland, and on a high level, though they were densely populated, and eventually, owing to their defective local sanitary conditions, were visited severely.

The state of the atmosphere which thus favours the transmission of the cause of cholera from one country to another, and that opposite state which, at other times, and in other places, prevents its transmission, thus arresting the progress of an epidemic, or in a country or town where it already exists, causing it to become extinct, are in part known only by their effects. The former state is generally associated with a high temperature, the latter with a low or falling temperature; but there occur very remarkable variations in the intensity of the epidemic not referable to temperature, which show that some unknown conditions of atmosphere, though not the cause of cholera, exert a powerful influence over it, either by acting directly on the material cause of the disease, or by affecting other matters in the air which enable it to exist or increase.

It may further be inferred from the very local and persistent character of the outbreaks of cholera, that its cause is not a gaseous, diffusible substance, but rather matter in the form of solid or liquid particles capable of attaching themselves to surfaces of other bodies.

3. The results arrived at with reference to the means by which this cholera poison is conveyed from place to place, have next to be noticed. A large body of evidence renders it certain that human intercourse has, at least, a share in the propagation of the disease, and that it is under some circumstances the most important if not the sole means of effecting its diffusion.

The progressive advance of the epidemic along great lines of human traffic, the rate of its progress varying according to the activity and means of human intercourse in different countries, but in no case surpassing the rate at which men travel; and its appearing first at the seaports of any island or continent which it is newly invading, are evidence only of the presumptive kind. A nearer approach to actual proof is made in the facts that in India the disease has travelled for some hundreds of miles, and during several months in the teeth of the monsoon, and is therefore, at least in some cases, independent of the influence of the wind; and that it has continued to prevail in ships during voyages of many weeks after they had left the infected parts, and also among bodies of troops for several weeks, while they marched through districts till then healthy.

Evidence of a more direct kind is afforded by the circumstance that the origin of the disease in the seaports of the British Isles and America has, in a large proportion of cases, been immediately by the arrival of ships from infected ports, and of ships actually bringing persons already affected with cholera. In several instances, too, the first patients attacked in these ports had had communication, more or less immediate, with sick persons brought in the ships; and in some cases the facts have been such as to admit of no other explanation than that the disease was imparted by the infected ships.

The information collected relative to the origin of the disease

in different towns, parts of towns, and public establishments in England, also leaves no reasonable room for doubt that the disease is propagated by human intercourse on land as well as by sea; and the published works, giving accounts of recent epidemics in France, Germany, and America, support the same view.

The facts, however, by no means sanction the belief that cholera is always propagated in this way. On the contrary, it is certain that the extension of the disease over large towns, if not over large areas, may take place independently of communication between the sick and the healthy. This is proved by the frequent outbreak of the disease within public establishments, such as prisons and lunatic asylums, in almost every case without a source of infection being traced, and likewise by the rapidity with which the arrival of an infected ship, or the occurrence of the first indigenous case in a large city, is followed by the appearance of the disease in various and distant parts of the city; the extension of the epidemic having, in some of these cases, been so rapid that several hundreds or even thousands of persons have perished in the course of between two and three weeks.

In the cases where human intercourse cannot have been the means of diffusing cholera, the agent most likely to have conveyed the poison from one spot to another is the wind. The poison of cholera being so dependent on the states of the air for its existence, increase, and power of action, and having the capability of passing from place to place, must, it would seem, not only be exposed to the air, but, even though it be in part attached to the surface of bodies, must, in part, also float in the air. The statistical facts, showing the extension of the disease from centres, and the successive attacks of different localities, though they might be owing to the transmission of the cause of the disease by human intercourse, are quite in accordance with the view that the poison is scattered by varying atmospheric currents from the foci in which it had been developed and increased; while the extension of the disease through a large city with the rapidity above mentioned is, indeed, explicable by no other agency, if the cause of the disease be a substance of the nature supposed. There are, however, few direct observations which tend to confirm this view, by showing a correspondence between the position of a place newly attacked with regard to an existing focus of the disease, and the direction of the wind at the time or just previously; and of the few observations which exist, the majority are unsatisfactory.

If, however, the atmospheric currents, as is most probable, share with human intercourse the office of disseminating cholera, their part would seem, from the facts communicated to the Cholera Committee, as well as from theoretical considerations, to be rather the diffusion of the disease over limited areas, its transmission

from some spots to others near at hand, than its conveyance to distant places, which is probably effected, in a majority of cases, by the locomotion of men. But the proportion of instances in which the introduction of the epidemic into towns, parts of towns, and individual houses or public establishments has been due to the one or to the other mode of diffusion, cannot at present be determined.

4. The propagation of the disease by human intercourse does not prove its contagious nature. If the poison of cholera increases in, or under the influence of, damp and impure air, and is likewise capable of attaching itself to the surfaces of bodies, to the walls of rooms, and to furniture, it will also be collected by the clothes of persons living in infected dwellings, will be carried by them from place to place, and, wherever it meets with the condition favourable to its increase and action, will produce fresh outbreaks of the epidemic. That its propagation in such a mode as this is at least more frequent than its communication by virtue of true contagion is to be inferred from the impossibility of tracing communication between the first and subsequent cases at the commencement of the epidemic in a large city; from the apparent impossibility that any direct communication can have taken place in many of these cases; still more decidedly from the great rapidity with which the disease sometimes spreads at once through the whole population of a city; from the influence of season and temperature, and of the characters of localities on the rate of the diffusion of the epidemic; and from the occasional alternations of its intensity during its prevalence in a town. The ultimate cessation of an epidemic throughout a country, and even a continent, the restriction of its course in crossing a continent to a tract of comparatively limited extent, furnish, perhaps, still stronger objections to the theory of contagion; for not only are they, like most of those before mentioned, characters which diseases known to be in the strict sense contagious do not present, but they suggest the belief that the propagation of the disease cannot be maintained by any matter emanating from the bodies of the sick.

Some facts, which constitute presumptive arguments, of more or less force, in favour of the dependence of the epidemic on contagion, namely, the relation, as a general rule, borne by the numbers of the population of a town, and even of a public establishment, to the duration of the epidemic there, the successive attacks of the different inmates of a house, or of the ward of a lunatic asylum, the ultimate cessation of the disease after a limited number of days in each house or ward, and the fact that, in the cases where the introduction of the disease into a locality has been traced to human intercourse, the supposed vehicle of the infection has usually been a person already suffering from the disease, or clothes or bedding

which had been used by the sick in other places, all these facts have been found susceptible of explanations in other ways; though the explanations offered have in some instances been necessarily of a conjectural nature.

With reference to two other arguments which, if established, would only prove that cholera is in some cases contagious, the

evidence examined has been found contradictory.

The frequent communication of the disease by the clothes or bedding of the sick, to the persons who handle or wash them, under circumstances rendering other sources of infection, than emanations received from the bodies of the sick, improbable, appears to be by no means proved. The preponderance of evidence is, in fact, opposed to its occurrence. On the other hand, the evidence respecting the especial liability of nurses or others attending on the sick to suffer from cholera, though conflicting, is in some instances of such a character as to preclude the absolute rejection of the view that the disease has a contagious property, even though it

does not usually spread by virtue of contagion.

5. The question, whether the poison of cholera enters the body through the lungs, or through the alimentary canal, has not been conclusively solved; but no sufficient reasons have been found for adopting the theory that the poison is swallowed with the food or drink, is reproduced in the alimentary canal, and being discharged with the secretions of the stomach and intestinal canal, propagates the disease by finding access in the same vehicles to the stomachs of other persons. This theory has been found especially inconsistent with the great mass of evidence which establishes the influence of the different conditions of the atmosphere, and with nearly all those facts which would be equally opposed to any view which regards the human body as the exclusive nidus for the reproduction of the morbific matter. So much, indeed, as attributes the extension of the disease among the inhabitants of groups of houses or of towns, to the contamination of the water used for drinking or culinary purposes, undoubtedly explains very readily some facts in the history of the epidemic in London; but even this part of the theory is found to be supported by very scanty evidence, and to be contradicted by other facts which, if they do not prove that the character of the water drunk is altogether destitute of influence with reference to the diffusion of cholera, at least show that its power and effects are very inconsiderable in comparison with those of other conditions.

Of the six theories then, above-mentioned, that alone is supported by a large amount of evidence which regards the cause of cholera as a matter increasing by some process, whether chemical or organic, in impure and damp air, and assumes that, although, of course, diffused with the air, it is also distributed and diffused by means of human intercourse.

This theory explains much that would otherwise seem capricious in the course of cholera; and it elucidates the relation subsisting between cholera and other epidemic diseases. Several epidemics of cholera have sometimes been immediately preceded in some countries or cities by the prevalence of fevers or of diarrhœa and dysentery; and this has been made an argument in support of the vague notion of an "epidemic constitution." For it has been supposed that this epidemic influence, in the course of its development, gives rise at one time to fever or diarrhœa, and at another time to cholera. It has been imagined, therefore, that the prevalent diarrhea preceding cholera resulted from the slighter of the peculiar atmospheric influence which subsequently produced the more formidable epidemic. But it is not by any means a general rule that an increasing prevalence of diarrhœa, or of any other epidemic disease, precedes the appearance of cholera; and the occasional association of cholera with such diseases is capable of being otherwise explained. The fact in question, and the similarity of the local conditions favouring cholera and epidemic diseases generally, together with other facts examined, seem to agree best with the view that these several diseases are caused by different poisons, all of which find their means of increase in similar states of the atmosphere, though there probably are modifications of the atmospheric conditions more essential or more favourable to some of these diseases than to others.

In the statement that the theory above indicated is the only one supported by a large amount of evidence, it is not implied that this theory is adopted to the exclusion of all others. For the possibility that cholera is occasionally communicated by a virus produced in, and emanating from the sick, has already been admitted; and other questions relative to the means by which the cause of the disease is disseminated, and its introduction into the human body effected, have been left open for further inquiry."

PREVENTIVE MEASURES.—The ultimate object of an inquiry into the subjects treated of, are stated by Dr. Baly to be—the discovery of the means by which the onward progress of cholera may be stayed—its increased diffusion moderated, and individuals protected from its attacks. The partial attainment of this object is all that at present can be hoped for; but the principles by which endeavours to attain it thus partially should be guided, are, for the most part, free from doubt.

"From among the great features of a cholera epidemic three stand forth as of paramount importance:—one, the undoubted influence of locality and of the sanitary condition of towns and dwellings on the degree of severity with which the epidemic visits them; a second, the equally certain influence of season and tem-

perature, together with some unknown condition of the atmosphere on the general prevalence and rate of extension of the epidemic; and a third, the share taken by human intercourse in determining not only the progress of the epidemic, and the direction of its advance, across a continent, but also its extension from continent to continent, and most probably its communication from one town to another in the same country, and from one locality to another in the same town. This third feature of the disease is less generally admitted than the other two; but the question here is not whether the evidence is so complete as to compel belief, but simply whether it amounts to so great a degree of probability that it cannot merely be neglected in the consideration of the measures to be taken for the prevention, arrest, or mitigation of a destructive pestilence.

1. With regard to the *first* of the features of the epidemic, there cannot be a doubt with regard to the course of action which it calls for, as the duty and interest of every portion of the community. The more fortunate classes are, it is true, exposed to proportionably little danger, since they are often able to leave the neighbourhood of spots in which the disease is raging, and usually dwell in more elevated, open, and airy parts of towns, and in more spacious and

cleanly and less crowded houses.

But it is also true that the power of the disease through a town is increased in proportion to the degree in which the conditions of insalubrity referred to are present in various parts of it; and not merely the poor, who live in the spots where moisture and foul air feed the cause of the disease, but all the inhabitants, are exposed by the existence of these evils to a greater risk of becoming its victims. And further, it is certain that the more intensely the epidemic prevails in a large town, the more does the whole district for miles around suffer, and the more danger is there of its

being propagated to other districts.

By improving the drainage in low parts of the town, opening close courts, thinning the buildings in the more crowded parts, putting a stop to the burial of the dead in large cities, keeping even the smallest streets constantly free from filth, covering drains and sewers, and abolishing cesspools, and other sources of foulness in the air and soil; by improving the dwellings of the poor in respect of ventilation, giving them the means of maintaining a due warmth in their rooms, without excluding the external air, promoting the general substitution of good water-closets for open privies, inculcating cleanly habits among the poor, and affording them that most important requisite, an abundant and constant supply of good water, by means of which they may attain cleanliness; by adopting these measures, it cannot be doubted that the ravages of the disease would not only be lessened among the people dwelling in the localities thus improved, but also greatly weaken the force

A close inspection of all such vessels should be made on arrival in port; and it would not be unreasonable to require that, in consideration of the restrictions of quarantine being abrogated, there should be brought with each ship, coming from an infected port, an official certificate of its having been inspected, and found cleanly and not overcrowded, and the crew healthy at the time of

its sailing.

On the arrival of ships having persons ill of cholera on board, or having had deaths from that disease during the voyage, more active measures must be adopted; and the best that have been recommended seem to be the removal of the sick to a hospital ship, moored at a distance from the other shipping in the harbour, or to a special hospital in an isolated and airy situation on shore; 2, permission to the rest of the crew to land after exchanging their dress for fresh clothes provided from the shore; 3, the thorough exposure of all articles of dress and baggage to the air and disinfecting agents before they are removed from the ship; and 4, the thorough cleansing of the ship itself, with the free use of disinfecting agents in every part of it, but especially in the parts occupied by the crew and passengers, or their baggage.

If, notwithstanding such precautions as these, cholera finds its way into the country, then the low lodging houses frequented by vagrants and the vagrant-wards of workhouses should be narrowly watched. For these especially are the places in which the disease is fostered, and whence it seems to be distributed widely to other localities. In these establishments, then, the most scrupulous cleanliness and free ventilation should be maintained, and even the personal cleanliness of the inmates as far as possible

enforced

When cholera appears in the places referred to, or within dwellings of the poor, intercourse with the surrounding population cannot of course be interdicted. But still it is possible to adopt measures which would not only check the extension of the disease among the inhabitants of the infected houses, but greatly diminish the risk of its propagation to other localities. Of these the most important is the provision of spacious and well-ventilated buildings in airy dry sites for the reception of the inhabitants of the infected spot, while their dwellings are cleansed and disinfected. These "Houses of Refuge," it cannot be doubted have saved many lives from destruction by cholera, both in this country and on the continent. No considerable town should be without one; and several should be prepared in the environs of the larger cities. The "Houses of Refuge" would receive the healthy, but for those already labouring under cholera other asylums must be found.

There has been much difference of opinion respecting the desirableness of establishing Cholera Hospitals. But it surely can-

not be disputed that those struck with cholers amongst the poor, ought to be carried to some hospital, if they are at all in a fit state to be removed. They cannot be properly treated at their homes, and mere change to a purer air offers them a better chance of recovery. Moreover, in the rooms in which the poor are struck with cholera, those who nurse them, and in a less degree those who visit them, are exposed to danger, probably not from contagion, but in most cases from the pestiferous atmosphere of the locality; while, if the sick are placed in the spacious and wellventilated ward of a hospital, nearly all danger from approaching them is at an end. Wherever, therefore, general hospitals do not exist, or cannot afford sufficient space, Cholera Hospitals should be established.

The buildings selected for the purpose should, if possible, be situated in the least crowded parts of the towns or districts in which they are needed. They should have spacious rooms, with provision for permitting the free circulation of air through them; and during their occupation, the most strict attention should be paid to their ventilation and cleanliness: otherwise they may

prove an injury rather than a benefit.

In the general hospitals, too, the same conditions of ample space, free ventilation, and scrupulous cleanliness, are the essential requisites. If they are provided, it is probably a matter of little moment whether the cholera patients are placed together in special wards, or dispersed among the ordinary patients in the

general wards of these establishments.

Whether the sick be removed to Cholera Hospitals, or to wards of general hospitals, it should equally be an object of care to prevent the accidental introduction of infection together with the patients, and with this object it is desirable that the clothes brought with the patients should be removed from them in a special receiving-room, or at once be either destroyed, or subjected to a disinfecting process. And even in the case of the healthy removed from infected dwellings to houses of refuge, care should be taken that the foul clothes be not carried with them, and that the clothes they wear should as soon as possible undergo an efficient cleansing.

The propriety of adopting measures founded on the belief that human intercourse aids in the dissemination of cholera, has been urged upon those who may still doubt the propagation of the disease in that way. And, on the same principle, other precautions, suggested by the view that the discharges from the stomach and intestines contain the cholera poison, must here be recommended, although the theory based on that view has been found generally untenable, and, at most, susceptible of very partial application. The precautions referred to are, the immersion of the

soiled linen of the patients, and subsequently of their bedd water to which some disinfecting liquid has been added; can either the food nor the drinking water can in any way be minated by the discharges of the patients, and especially the on the part of nurses and others about the sick be taken to their hands before taking food; to which may be added measure more feasible in hospitals than in the dwellings poor, the placing a small quantity of chemical decomposing in all the vessels into which the discharges of the patients ceived.

One other method of combating the pestilence has bee posed and partially carried into effect; namely, the house-to visitation of infected districts, with the view of discovering treating all cases of diarrhoad, some of which may be presured be cases of cholera in an early stage. This measure, how based on principles which do not properly belong to the soft this report. It would, therefore, be out of place here quire into the amount of success with which it had be tended."

PREVENTIVE MEASURES IN THE ARMY.—The lowing "Instructions," framed by desire of the Director-Gof the Medical Department of her Majesty's Army, were under authority of the Secretary-at-War, and dated War August 31st, 1848. "These Regulations are to be considered General Orders, and are to be observed as such by all personant whom they may apply:—none of them are to be modified regarded, unless special reasons shall exist for so doing, an not then, without the consent of the Director-General, unled delay which would be necessary to obtain that sanction, was to be the occasion of injury, either to individuals or the In the event of there being reason to anticipate the latt alterations considered necessary may be effected, but their and extent must be immediately reported by the medical who adopts or advises them."

"When Cholera shall have been officially reported to be previous the country, but distant from Military Stations, the jing Instructions are to have effect:—

1. Medical officers will exercise more than common vig in the discharge of their professional duties; they will more than ordinary attention to the interior economy of corps, the constitution of the men, and to every circums however remotely affecting their health. They will be exp to be cognizant of every military arrangement involving the health of the troops, and to put themselves in communication with their commanding officers on any alteration of arrangements in the ordinary duties, which they may consider beneficial or likely to afford greater security.

2. Every possible precaution must be taken to guard against intemperance, crowding in small ill-ventilated rooms, use of unwholesome food, deficiency of clothing, bedding, fuel, &c.: and, if found to exist, measures must be taken to correct or remove

them.

3. Medical officers are to attend the ordinary parades, and ob-

serve the health of the men, without exciting suspicion.

4. Great attention to personal cleanliness is to be enjoined, and the men cautioned carefully to avoid unnecessary night exposure and damp, and to change their clothes when wet.

5. Especial care is to be observed in the cooking of vegetables;

and lamb, pork, and stale fish interdicted.

Roasted or baked meat is to be provided twice a week, instead of boiled meat.

7. Drains, dust-holes, privies, and the removal of accumulated

filth, are to be specially attended to.

8. The barrack-bedding is to be aired daily, but not in the open air in winter; and great caution observed as to the perfect dryness of fresh supplies.

9. As attacks of the disease seem to be frequently determined by exposure to wet, damp, unusual intemperance, and other irre-

gularities, every means should be taken to prevent those.

10. At this period all ailments, particularly diarrhea, require the strictest attention; and the commanding officer should explain to the men the importance of immediately reporting themselves when they feel in any way unwell.

11. Where there is any suspicion of the approach of cholera, immediate steps are to be taken to establish a separate ward in hospital, to which every case not strongly marked is to be sent on

admission.

12. An abundant supply of hospital bedding, dresses, and medicine, is to be kept in readiness.

# When the Disease has appeared among the Military of a Station, or the Inhabitants in its immediate vicinity.

13. The avoidance of all unnecessary alarm cannot be too strongly enjoined; and it is hoped that medical officers, by their own example, will endeavour to allay apprehension in those suffering from, or in any way connected with, the disease.

14. On the appearance of cholera in a corps, health-inspections are to be made at morning and evening parades, and a daily inspection of every individual attached to the regiment.

15. Each soldier is to be provided with two cholera belts, as part of his necessaries. Flannel-waistcoats, if thought necessary

for individuals, are to be provided at their own expense.

16. Married men (if out of mess) should each be provided with a ration the same as the single men; and it is also desirable, in barracks in which the disease exists, that the women and children should have sufficient and regular meals.

17. Soldiers should be cautioned against intemperance; and drunkards and all men of weakly or susceptible habits, should be limited to a certain quantity of liquor at the discretion of the me-

dical officer.

18. As a precautionary measure, but particularly in infected localities, drills, parades, and duties generally, should, as much as consistent with discipline, be reduced, and favourable hours and weather chosen for them; but, above all, the number of sentries, especially at night, should be diminished so that in no case shall the men have less than three nights in bed.

19. Good fires are to be provided in the barrack-rooms, to increase ventilation, and to diffuse a cheerfulness,—which last

should be promoted in every way.

20. Coffee or warm drinks are to be provided to the men before morning and night duties. A hot evening meal enforced, and breakfast, if possible, supplied at the usual hour to every man be-

fore leaving his barrack-room.

21. In the event of the appearance of cholera among the civil population in the neighbourhood, the troops should be confined to barracks, and all intercourse prevented. When the disease prevails in a corps, it may be found beneficial to encamp it—a proper site being selected for the purpose.

22. During confinement to barracks, the minds of the men should, as much as possible, be amused and occupied; and, under proper regulation, occasional marches into the country, and trap-

ball or other games in an adjoining field, permitted.

23. Officers' servants are to be under the same restrictions as others, and no person from the town to be admitted into barracks.

24. If practicable, a considerable reduction of the numbers in barrack should take place, as well to ensure a purer atmosphere and more thorough ventilation, as to make room for the accommodation of women and children, and, if necessary, for a temporary hospital, or observation ward. Over-crowding, under any circumstances, is to be avoided—great attention paid to cleanliness and ventilation, and the floors and passages dry rubbed—not washed.

25. Personal washing, and that of clothes, is to be done in sheds or store-houses appropriated for the purpose, and not in the barrack-rooms.

26. The quality of the beer to be used by soldiers is to be ascertained by a competent person, and the acid in porter or ale,

corrected by chalk or carbonate of soda.

27. Where cholera is present in the neighbourhood, the women and children are to be accommodated in barracks, or aired houses in the vicinity, and to be put under similar restrictions as to intercourse. Where this cannot be done, soldiers with their families who are permitted to live out of barracks are to be excluded therefrom, or placed in a temporary barrack until it is considered safe to admit them-their quarters during this separation being frequently inspected.

28. Where a house is hired for the women, they may be required to pay their usual rent, and coals supplied by Government

at prime cost.

29. Diarrhœa, either as precursory, co-existent, or prevailing by itself, should as a measure of safety, be regarded as closely allied to cholera; and where bowel complaints prevail in a corps stationed in an infected district, a spare room should, if possible, be set apart for the accommodation of the healthy of any room that may become unusually subject to that complaint; and the room thus vacated must be whitewashed, cleansed, and fumigated, so as to be ready for any similar occurrence. This applies still more strongly where cholera actually exists in a corps. Cases of common cholera are to be removed to the Observation Ward.

30. The regimental hospital is to be appropriated to the treatment of cholera, and the ordinary cases of sickness accommodated in barracks; or where this cannot be done, in a hired house, the authority of the Director-General for the latter arrangement being

previously obtained.

31. The appearance of cholera in places where troops are stationed is immediately to be reported to the Director-General, without waiting for it being officially announced by the local Board of Health.

32. From the moment that a case of common or spasmodic cholera occurs, a daily report is to be forwarded to the Director-General until further orders.

33. A full report of each case among the troops, with detail of previous habits, intercourse, diet, exposure to cold, wet, &c., and other particulars, is to be forwarded to the Director-General.

34. Daily reports of the progress of cholera among the civil population where troops may be stationed, will be required from

medical officers.

35. Cases among women and children in barracks are to be

admitted into hospital; those in quarters to civil hospital, und such regulations as may be adopted in parishes; for the former separate ward, or barrack-room, is to be held in readiness.

36. Cases among the troops of the East India Company are be admitted into hospital, as also those specified as entitled

medical attendance in the hospital regulations.

37. Medical officers are to visit their hospitals frequently, at to state the hours of visit in the monthly sick report. They be required to be always available for any sudden call of the svice; and when cholera prevails in the corps, are not to leave barracks, except on imperative duty.

38. Smoking, to those habituated to the indulgence, may

permitted.

39. Corpses are to be removed to the dead-house without del buried as soon as possible, and conveyed in a covered cart—on men's shoulders.

40. Post mortem examinations are not to be discontinued, performed under such modifications as the occasion will read

suggest.

- 41. Bodies of patients dying of cholera are to be sprinkled we chloride of lime, to be wrapped in a coarse sheet steeped in strong solution of the same, and some of the powder should be just the coffin.
- 42. Medical officers are not to visit civilian patients in toy and are to conform to the same regulations and instructions other officers.
- 43. In infected places, Divine Service should preferably be p formed in barrack, instead of marching the troops to church.

44. The old clothes of recruits are to be washed and fumigate

or if necessary, destroyed, and new clothes issued.

45. The clothes and bedding of cholera patients are to be mersed in cold water for forty-eight hours, then washed steamed in boiling water, and dried in the open air.

46. The place whence a patient is taken is to be thorough washed, the bedding and bedstead removed, and fumigation may be chlorine gas, if possible, if not, by nitric acid gas, or fumes vinegar.

47. The barrack bedding is to be removed with cholera or s

picious cases to the hospital, or Observation Ward.

48. Provisions brought from town should be delivered at

barrack gate.

49. Deserters, recruits, and men from escort or furlough, rejoining, should be separated from the other soldiers for a per varying from seven to twenty-one days, according to circumstance

50. Attention is called to Her Majesty's orders, in the book instructions for regimental hospitals, on contagious diseases, ext

ordinary sickness, inspection of barracks and quarters, ventilation, and fumigation.

For the purification of drains, privies, clothes of recruits,
 c., chloride of lime will be issued by the Ordnance Department,

on the requisition of the commanding officer.

52. Bearers are to be applied for to the Ordnance Department, and a horse and covered cart hired for the conveyance of the sick, should the distance from the hospital require it.

I purpose next to present such extracts as are deemed necessary, from the report of Dr. Gull, on the Morbid Anatomy, Pathology, and Treatment of Epidemic Cholera. "The design" of his portion of the work, "has been to set down the principal ascertained facts, and to draw such conclusions as were possible from them."

MORBID APPEARANCES AFTER DEATH IN COL-LAPSE.—In the former, as well as in the latter epidemie, it was noted by several observers, that the icy coldness of the body, in the stage of collapse, passed away after death.

No precise thermometrical details were given; but the surface of the body was observed to have become actually warm. Latterly, it was proved by Dr. Barlow, of Westminster Hospital, and others, by careful measurements, that a rise of temperature on the surface of the body did occur after death. In one case, in which the coldness of the surface during life was very marked, the thermometer indicated 102 12 two hours after death; in another case, ten minutes before death, the temperature in the axilla was 103 1°, and five minutes after death, it rose to 104°; in a third case, the temperature of the uncovered body, twenty-five minutes after death, was 89°; forty-five minutes after death it was 90°; and fifty minutes after death, it rose to 91°.

Among the phenomena presented by the body Massair conafter death from cholera, are the well-known contractions of the muscles, which often occur to so great an extent, and last for so long a period, as to excite horror in the ignorant, and add in such minds a further mystery to this disease. The muscles affected were principally those of the extraonities, but contractions were occasionally observed in the other voluntary muscles; the duration of the contractions varying from a few minutes to two hours.

This phenomenon was observed more commonly in those who died rapidly of the disease, in the middle period of the, when the

muscular system was vigorous and well developed, which will, perhaps, account for their greater frequency in males than in females.

Cadaveric rigidity often supervenes very quickly; Rigor Mortis. in one case it began at the end of an hour, and in another after forty minutes. Its occurrence was not retarded by the high temperature which the body retained.

Cruveilhier says that putrefactive changes were Putrefactive slow, as in all cases where much blood has been changes. lost, and adds,-" In the alimentary canal, on the contrary, decomposition was rapid, as commonly occurs where the digestive organs are the seat of considerable sanguineous congestion." Most observers confirm the former part of this statement; but with respect to the intestinal canal, the conditions were variable, and often the contrary of that noted by this great authority.

The stomach was pale, and generally more or less Stomach. distended. It contained turbid, mucoid fluid, grey or colorless, or tinged of a chocolate or reddish-brown hue, by admixture with blood. The surface of the mucous membrane was covered with tenacious mucus, having in some cases a puriform character, from the large amount of exfoliated epithelium. In some instances it was pale; in others, hyperæmia, in different degrees, existed.

The membrane was generally rather thickened and opake, the

texture firm, and the surface mammillated.

The state of the œsophagus and pharynx was not Pharvnx and generally reported. In one case, fatal after twelve Œsophagus. hours of acute symptoms, it was observed that the lower part of the esophagus was deprived of its epithelium.

The small intestines had generally a pink or rose Peritoneum. external tint from hyperæmia of the portal venous system. In some instances they had a remarkably dark color, the venous trunks being large, and full of pitchy blood. The different

tints soon changed upon exposure to the air.

The coats of the small intestines were thickened Small intesand pulpy, from ædema of the mucous membrane tines. and sub-mucous tissue. The duodenum and ileum were more commonly affected, and to a greater extent than the jejunum. In some instances the mucous membrane was pale throughout; in others, the lower part of the ileum only, or the duodenum only was hyperæmic.

The vascularity of the mucous membrane presented itself under two conditions, as uniform arborescent venous injection, affecting large tracts of the intestine, particularly the lower part of the ileum; or as patches of variable extent, in which the redness was punctate, and of a bright color, frequently with spots of ecchy-

mosis, and an exudation of tenacious mucus.

The villi were also small and prominent from ædema, especially throughout the jejunum. Their appearance, and the condition of the epithelium covering them, was minutely described by Böhm, during the last epidemic of 1832-33. Quoting Dr. Gairdner, the report says :- "The most frequent of all the abnormal conditions of the mucous membranes, was the prominence of the intestinal glands, both of the aggregated and solitary, but especially of the latter. This condition, the 'psorenteric' of some French writers, was found in about two-thirds of the cases."

The changes in the patches of Peyer were similar The patches of to those occurring in the villous surface of the mucous membrane generally, and in the solitary glands.

They were often thickened and prominent, from infiltration of serous fluid, and the included glands, when visible, presented the conditions found in the solitary glands of the mucous membrane elsewhere. There is no evidence that they were the seat of morbid changes, peculiar to them in kind or in degree.

The large intestine was more rarely affected with Large intestine. hyperæmia and ecchymosis than the small intestine.

In many instances it presented nothing abnormal beyond a greater

distinctness of the solitary glands.

In only four, out of thirty-four cases, were the solitary glands prominent throughout the whole extent of the large intestine; in eleven others they were slightly enlarged about the eccum and ascending colon. They were often translucent, from distension of their cavities with serous fluid; sometimes their distension had been followed by rupture, producing an appearance of a small rounded ulceration.

The state of the spleen is noted in twenty-three cases fatal in the algide state :- of those, in three it was of a natural size, and in eighteen it was small, or very small. It weighed but two ounces, two and a-half, and three ounces, in seven cases recorded. In most cases there was no obvious change in the tissue, beyond that arising from a want of blood, giving its capsule rather a wrinkled appearance, and enabling it to resist laceration, the texture being often described as pale, firm, and dry.

After death in the algide stage, the liver was generally diminished in bulk, the tissue flaccid, and the capsule finely wrinkled. The larger veins, both the hepatic and the portal, but especially the latter, were often full of dark viscid blood. The lobular appearance of the secreting structure was indistinct, and the whole tissue of rather a lighter red than usual; but to this rule there were exceptions.

The secreting structure, on a microscopical examination, pre-

sented nothing abnormal.

curred after reaction; but as, in general, these last do not appear to have had much of important distinctness and of difference from the morbid appearances already described, I must here refrain from quoting them. In the instances of persons dying during the febrile reaction consequent on the cholera collapse, we should naturally expect to find congestions of the gastro-intestinal mucous membrane, with hyperæmia and serous effusions in the brain.

PATHOLOGY OF CHOLERA. - "The first unequivocal symptoms of cholera indicate that the gastro-intestinal mucous membrane, with its ganglionic nervous centres, is the focus of the morbid action.

The morbid appearances characteristic of cholera are most marked in the small intestine, duodenum, and Pathology of stomach, and the general symptoms indicate an early the cholera and severe depression of the ganglionic nervous process in the intestines.

centres of these parts.

The principal phenomena which arrest attention are the ædematous state of the mucous membrane, and the more or less extensive patches of capillary and venous hyperæmia, and ecchymosis. These, together with the character and amount of the fluid effused, demonstrate an important lesion of the circulation through the affected parts. How this is produced, can at present be no further elucidated than by the hypothesis of a specific poison, acting upon the ganglionic nervous centres, or upon the mucous membrane itself. The former appears to us the more probable supposition, from being more in accordance with some other phenomena of the disease, as the profuse sweating and the sudden and severe collapse, which, as will be seen hereafter, is not in a necessary and constant relation to the discharges from the mucous membrane. Either would, however, suffice to explain the altered condition of the circulation, since it is well known that if one of the agencies in operation in the capillary circulation be abnormal, there will be a corresponding change in the ease and rapidity with which it is sent through the tissue. In this respect nearly the same result is produced, whether the blood be unfitted to stimulate and nourish, or the nutrition in the part itself be perverted or defective, or the vessels be deficient in nervous supply. We no longer look for mere mechanical causes of vascular retardation; and whatever light may hereafter be thrown upon this stage of cholera, it will no doubt come from a more intimate knowledge of the pathology of the capillary circulation.

These morbid changes in the first stage of cholera are of the simplest kind;—the tissue is infiltrated, and the glandular follicles are distended with the same watery fluid as escapes into the cavity of the intestine;—the epithelium, by this maceration, readily separates after death, and to a certain extent during life, but not so largely as Böhm supposed. Exfoliation of the epithelium appeared to this excellent observer to constitute an essential part of the primary morbid action, but further investigations show that during life it occurred to only a very limited extent, and was in itself an unimportant change. The coats of the capillaries and smaller veins are often ruptured, giving rise to ecchymosis;—this, with the punctate and stellate character of the hyperæmia, denote a want of tonicity in the vessels.

As death occurs only when the phenomena reach their greatest intensity, and when the circulation in the nervous system generally is much retarded, the ramifications of the intestinal veins in par-

ticular are often distended with dark blood.

An examination of the fluids effused from the mucous membrane gives no evidence of active plasmatic changes taking place in them. On the contrary, the large amount of fluid thrown out, its low specific gravity, and its other physical characters, indicate an almost passive exosmosis, as through a dead membrane. Some observers have referred the morbid changes to a catarrhal condition, others have regarded the disease as a form of serous hemorrhage, and the Berlin pathologists, whose attention was particularly arrested by the occurrence of amorphous granular fibrin in and upon the affected surface of the mucous membrane, designate it a destructive diphtheritic inflammation. We believe that, for the present, such generalizations, however plausible, are of little value, and that we arrest inquiry by their adoption. The depression of the capillary power—the extreme exhaustion of the great ganglionic nervous centres in the abdomen-the passive character of the lesions of the mucous membrane-its normal action being reversed to a fatal exosmosis—are peculiar to cholera, and give it an individuality which forbids our merging it for the present in any general category.

Although the intestinal tract is the principal seat of the morbid actions, they are not limited to it. The kidneys, at an early stage, are sometimes affected; the urine having been found albuminous prior to its suppression, and the secreting tissue and the lining membrane of the pelvis present occasionally, after death, lesions of

the same character as those observed in the intestine.

In the female the uterine organs are similarly affected, the lining membrane of the uterus and vagina being frequently much congested and ecchymosed, with commencing diphtheritic exudation upon it.

The liver is free from disease, except in some rare cases, where the lining membrane of the ducts and gall-bladder is the seat of the cholera process, their contents being then of the rice-water character. The distension of the gall-bladder with bile is nearly constant, but cannot be referred to as a pathological indication of any moment, as such a condition is common when the digestive function is long interrupted, and indicates a passive rather than an active state.

There is a hypothesis regarding the nature of cholera, based upon the supposition of a suppression of the hepatic secretion and consequent congestion of the liver; this is altogether unsupported by anatomical facts. The absence of bile from the evacuations is not a necessary phenomenon of the early stage of cholera. In a large proportion of cases the diarrhœa ceases to be bilious only when the more intense symptoms set in, and even then, the rice-water fluid often gives, with re-agents, distinct evidence of the presence of bile; but when it is no longer passed in the evacuations, the secretion is not altogether suppressed, as shown in the contents of the gall bladder and ducts after death. The hepatic function does not appear to be subject to any further derangement than that which naturally follows upon the retardation of the circulation during the stage of collapse.

It would have been unnecessary to advert to this hypothesis had it not widely influenced the treatment of the disease. Dr. Avre, the great advocate of the administration of calomel, gives the following rationale of the cholera process. He says:-"Now, there is one condition which is uniformly and conspicuously present in malignant cholera, and is, indeed, characteristic of itnamely, a suppressed or suspended secretion of bile, as shown by the diminution, and at length the total disappearance of it from those watery discharges which are poured so profusely from the stomach and bowels. As a consequence of this cessation of the hepatic function, an accumulation," he adds, "will take place in the liver, of venous blood, and an impeded circulation result from it, producing a congested state of that organ, and subsequently, by a retention of the blood in its course through them, of those abdominal organs whose circulation is associated with it. Now the congestion thus produced in the portal venous system of the liver and its associated organs, constitutes the stage of collapse, and under various modifications and grades of intensity, whose real nature and amount are unknown, forms the essence of it in all."

Such a train of reasoning is unsupported by any evidence. The serous rice-water character of the cholera stools is obviously due to special pathological changes in the mucous membrane, and not to any merely mechanical congestion of a secondary kind, as here stated. What these pathological changes are, we have

endeavoured to show, and have referred them to a specific action of the cholera poison operating through the blood; a more minute

account we are unable to give.

The appearances, after death, in the chest and cranium show that the viscera in these cavities were not primarily affected. The occasional emphysema of the lungs, their emptiness of blood, except at the posterior and inferior parts, and the inelastic compressed character of their tissue observed in some cases, are explicable by the state of the respiration and circulation at the last stage of life. The congestion of the veins, and the effusions under the membranes and into the ventricles of the brain, are referrible to the same cause."

Cholera has been classed among zymotic diseases, but its clinical history and morbid anatomy are opposed to the theory of its being due to a zymosis, in a strict zymotic or consense of the term. In a zymotic disease, the Lun intagious disduces certain plastic changes, from which results its augmentation. In cholera we have no evidence of such changes; the alterations in the blood, so far as they are known, being referrible to the loss of its fluid parts, in accordance with the physical laws of exosmosis. The local morbid action appears to be of a negative rather than of a positive kind. The marked depression of the organic functions, and the morphological characters of the effused fluids, as well as their general physical properties, indicate a passiveness almost peculiar to cholera. In cholera, the onset of an attack is frequently sudden, and the effects apparently direct, such as would follow the immediate action of an extraneous poison upon the body. The zymotic process is more gradual, and the symptoms follow a more constant rule with respect to them.

Cholera appears to consist of but one single series of actions, which may vary in intensity through every gradation, but, throughout, maintain the same character of passiveness. There is no febrile stage to which zymotic changes, as ordinarily observed,

and the production of a materies morbi may be referred.

Certain prodromata of a general kind have been enumerated as occurring prior to the symptoms which arise from the local morbid action in the intestinal mucous membrane, such as vertigo—headache—general malaise—muscular fatigue—faintings—wandering pains in the limbs—cramps—gripings in the abdomen—sense of weight at the precordia—and nausea. A cursory inspection of such a category suffices to show that, even if generally present, these indications would be unsatisfactory. Many of them arise from mental emotion, so prevalent during an epidemic, and others are not so much premonitory of an attack as the first symptoms of its onset, and the result of the morbid action already begun in the abdomen.

In most cases, a painless diarrhea, lasting for a variable time, precedes the more characteristic symptoms. Although all observers admit the frequency of this precursory stage, there has been some difference of opinion, and much unfruitful discussion about its pathology, whether it be a part of the disease, and due to the action of the poison, or merely a common diarrhea, upon which the specific effects may at any moment be engrafted. There are no characteristics which enable us to draw these distinctions with any certainty.

Experience has sufficiently shown, that during the epidemic the stages from a mild and apparently simple diarrhea to the ricewater purging and collapse are not definable, and that the former, if unchecked, does, in numerous instances, gradually pass into the latter, with its attendant collapse and fatal results. The diarrhea which prevails when cholera is epidemic is due to the same cause as cholera itself.

Numerous replies were received, all serving to establish the frequency of a stage of diarrhœa, lasting from a few hours to several days. A table fixing the period definitely of the duration of premonitory diarrhœa, exhibits a range of from one hour to ten days.

The evacuations were generally more profuse and liquid than usual, but otherwise of a natural appearance, often unaccompanied by pain, and passed without effort, the painlessness and passiveness giving a false security to the patient. It was not until the nervous system began to be depressed, and the feculent character of the stools was lessened or lost, and they became alkaline, watery, and flocculent, that they were distinctive. On this point, the experience of the profession appears to be uniform, and hence we may draw the following conclusion—that during the prevalence of the

epidemic, every case of diarrhœa arising without obvious cause,

may be regarded as a probable result of the specific poison. The cholera poison is not known to produce its fatal effects without the characteristic affection of the intestines. Cholera sicca, in a strict sense, does not occur, for although How far colthe disease may be fatal without any evacuation, the lapse depends upon loss of intestines, after death in such cases, have been found fluids. to contain the rice-water fluid. In some cases which came under our observation, on a post-mortem examination, the larger intestines contained healthy fæces, whilst in the upper twothirds of the small intestine, the mucous membrane presented the ordinary changes induced by the cholera process, and the ricewater effusion was abundant.

As many of the symptoms of the stage of collapse depend upon the loss of fluid, it has been too absolutely inferred that the general phenomena of the disease are always in a necessary relation to the amount of these effusions.

THE TREATMENT OF CHOLERA.—"It is well ascertained, that in the largest proportion of cases, at least in European countries, the poison of cholera produces its first effects on the system gradually, as indicated by diarrhæa, varying in duration from a few hours to several days before the intense symptoms supervene. This period may be called the period of invasion.

The numerous communications received by the College establish the importance of recognising this stage at its commencement, and render it highly probable that the morbid effects may then be often successfully commonly the period of investion.

bated.

The popular theory that the discharges are an effort of nature to throw off a materies morbi, is not only unsupported by any known facts of the disease, but, when applied to practice, is found

to increase the violence of the symptoms.

At present no antidote or specific medicine is known to neutralise the cause of cholera, or with certainty to arrest its early effects; but the communications received, show that the diarrhœa was, in a large number of cases, arrested by various combinations of such remedies as generally suppress discharges from the intestines, and prevent the exhaustion of nervous energy.

Opium was an almost constant ingredient, and was given in conjunction with astringents, aromatics, and diffusible stimuli.

A recumbent position is proved by experience, and also from the nature of the case, to be a most important measure. It prevents exhaustion, favours the circulation, and lessens the frequency of the evacuations. It is highly probable that cases, which otherwise resisted the action of medicines, would have readily yielded, had the horizontal position in a warm bed been strictly enforced. Enemata were not, as might have been anticipated, of much service, since the small intestine and duodenum are the principal source of the effusion.

The indications of treatment in this stage are formed entirely from the mucous membrane itself, independently of any hypothetical derangement of the liver. This organ does not appear to be in any way the cause of the symptoms; the diminution of the bile in the evacuations, as they become more and more fluid, being an effect, rather than a cause of the disease. The character of the fluid thrown out—the attendant symptoms—the absence of pain—the passiveness of the evacuations, and their occasional retention to a large amount in the intestine, indicate a depression of all the energies of the affected surface.

The amount of success attained by early treatment is not yet

determined; there is a general opinion that it was very githis must be received with some limitation, as the facts upon it is founded are not unequivocal. By far the greater nu cases of diarrhœa would probably never have passed beyon stage if no medicines had been administered; and, on the coin many instances the symptoms were uninfluenced by an ment, and fatal collapse came on in spite of every effort vent it.

Notwithstanding this uncertainty, the general results ventive measures were apparently very favourable, as sh the small proportion of cases which passed into the severe of the disease subsequently to early treatment."

The following are the results of the house-to-house visit the metropolis from September 1 to October 27, 1849:—

Cases of diarrhœa discovered.	Cases approaching cholera discovered.	Cases which passed into cholera discovered.
43,737	978	52

In various cities, as Dumfries and Glasgow, the same faveresults attended the system of house-to-house visitation; a no longer matter of doubt that the earlier the disease is a tered, the greater in an infinitely high ratio, are the advunder which medicines are employed to counteract it.

"On this division of our subject we can do little more that out, according to the evidence before us, what of treatment seem to have been of some service useless, or injurious; and by thus clearing the prepare the way for better directed efforts.

An enumeration of all the means proposed for the treat this stage would be useless; it is, therefore, only upon the pal of them that we shall make any report.

Calomel stands foremost, from having in this country used than any other remedy. The of the disease which has chiefly led to its e ment, is not supported by anatomical facts. The absence from the evacuations appears to be merely a subordinate Calomel can be administered only on empirical grounds, a value must be determined by the results so obtained; for appears to be no argument in favour of its exhibition either analogy or pathology.

The results in 365 cases treated by this remedy in sma frequent doses were 187 deaths, and 178 recoveries.

The numbers given by Dr. Ayre to the College of Physicians, do not sustain his favourable opinion of the treatment of cholera by small doses of calomel. The deaths were 365 out of 725 unequivocal cases.

Under various and opposite plans, the recoveries, even in severe cases, averaged from 45 to 55 per cent., according to the period of the epidemic; they should therefore exceed the highest of these numbers before they can be adduced in proof of the value

of any particular method of treatment.

In general, no appreciable effects followed the administration of calomel, even after a large amount in small and frequently repeated doses had been administered. For the most part it was quickly evacuated by vomiting and purging, or, when retained for a longer period, were afterwards passed from the bowels unchanged. Salivation but very rarely occurred, and then only in the milder cases.

We conclude that calomel was inert when administered in collapse; that the cases of recovery following its employment at this period were due to the natural course of the disease, as they did not surpass the ordinary average obtained when the treatment

consisted in the use of cold water only.

lants.

This is termed the "rational method" of treatment, and was intended to combine those remedies which seemed Treatment by best fitted to fulfil the supposed indications of this calomel, opium, stage. The calomel was given to restore the funcand stimutions of the liver, and as an alterative of the morbid action in the gastro-intestinal mucous membrane;

the opium to allay irritation and arrest the discharges; and the stimuli to counteract the depression of the nervous system. Experience did not confirm the theory: the results were unfavourable, and not altogether so indifferent as when calomel was exhibited by itself.

Although opium and diffusible stimuli, brandy, camphor, and ammonia were useful at an early stage of the disease, as collapse set in, they not only failed to produce any favourable result, but often aggravated the symptoms.

It seems well ascertained that opium in large doses was at this period injurious, by increasing the cerebral oppression, and embarrassing the system during reaction. It was probably less and less applicable as the disease advanced to its characteristic development.

Stimuli, especially the various preparations of alcohol, did not act as restoratives in collapse, but even increased the irritability of the stomach, and added to the sense of oppression at the præcordia.

The expectations excited by the early success apparently obtained by the use of chloroform, were not realised in its subsequent employment. It not unfrequently allayed the vomiting and cramps, but did not in any degree arrest the course of the disease.

The perchloride of carbon in five or ten grain doses, and a solution of camphor in chloroform, acted as powerful stimuli, but the results did not indicate that they possessed any specific therapeutic value. Although they produced symptoms of reaction, this apparent improvement was generally, in severe cases, but transient, and their continued use seemed to exhaust the little remaining power rather than to restore the patient.

The obvious requirements of the system, and the urgent thirst, were sufficient indications for the use of diluents, and the experience of the profession appears to be uniformly in favour of permitting patients to gratify their appetite for them. Cold water was generally preferred, and good results were often observed when it was taken freely in repeated and copious draughts, although it excited vomiting. In smaller quantities, and iced, it was refreshing to the system, and allayed the irritability of the stomach.

Ice was generally grateful to many patients in impending or approaching collapse, and probably acted favourably upon the mucous membrane, and served to arrest the discharges. The use of ice and cold water appears, in most cases, to relieve the burning thirst, while it favours reaction. Dr. Arnott administered a mixture of ice and salt in large quantities, and in the cases of two patients so treated, both recovered; but there is no further experience respecting this mode of treatment.

Salines of low specific gravity were sometimes given instead of common water, the intention being to restore to the blood a fluid similar to that lost in the early stages of the disease. We have no evidence that they possessed any influence over the local morbid action in the mucous membrane. It was not until this surface had in part recovered its function of absorption, that any good resulted from their employment. When given at an early period, and in a more concentrated form, they appeared to favour the discharges.

Emetics. Emetics were sometimes given at the onset of the disease, with the intention of cutting short the morbid action, by distributing the blood to the surface, and relieving the congestion of the intestinal mucous membrane. It is probable that, when administered early, they were occasionally of service; but the results appear to have been too uncertain to admit of their forming any part of a routine system of treatment. In collapse they were inadmissible, being often followed by increased exhaustion; and even when they appeared to produce symptoms of reaction, this effect was but of short duration.

Bleeding was employed in the premonitory stage of cholera, for the purpose of arresting the discharges by relieving the congestion of the intestinal mucous membrane. This was not much resorted to in the last epidemic; and the communications to the College contain but little mention of it. The only reports in its favour, by Annesley and other writers on the disease as it occurs in India, have not been confirmed by further observation; and hence, except in rare and exceptional cases, it is not usually had recourse to in the premonitory stage, or at the onset of the more severe symptoms. Its general inadmissibility is to be inferred from its almost entire disuse in the last epidemic.

Even in the consecutive fever blood-letting in any form requires much caution. Leeching the epigastrium and temples, or cupping the back of the neck, were sometimes of service in obviating the cerebral congestion of this period; but if carried to any great extent, were injurious by exhausting the patient.

The general inference, from all we have observed of the disease and collected from the experience of others, is, that large losses of blood, in the consecutive fever, are injurious; but that occasionally good results follow its local abstraction, according to the indications of the symptoms.

Quinine, strychnia, arsenic, sesquichloride of iron, nitrate of silver, nitrous acid, chlorine water, sulphur, sulphuric acid, bichloride of mercury, charcoal, &c. &c. The failure of these methods of

Empirical treatment. Specific remedies.

treatment, which, from being based upon some supposed indications of the disease, may be called rational, led naturally to the employment of almost every active medicine in the materia medica.

It is notorious that the results have been discouraging, notwithstanding the bold assertions to the contrary. The communications made to the College contain no data for determining the inquiry, nor is anything deserving the name of evidence in favour of the value of these means, to be gathered from the numerous journals and published treatises in this country and on the Continent. This part of our report must, therefore, be defective, although an unlimited amount of time and labour has been bestowed upon a perusal and comparison of the statements brought before us."

The report goes on to say, that the therapeutic history of cholera leads the authors to guard against the submission of remedies to what is termed a "systematic trial in a series of cases," by expressing their conviction that for empirical inquiry to be pursued to a good end, the most favourable opportunities are required for noting the state of the patient at the commencement, and at different stages of the treatment, by which alone discrepancies in the results can be reconciled.

Again, the employment, without definite scope, of one remedy after another, with the vague hope of at last finding a specific, is to be deprecated, not only because it can lead to no good result, but because it deprives the patient of that assistance which esta-

blished experience affords.

The state of the patient in the collapse of cholera is so unfavourable to the absorption of medicines, that even if we knew the remedy in itself most appropriate, we could not anticipate great results from its administration by the mouth at this period. Every consideration of this formidable malady urges upon us the paramount importance of obviating the causes which give rise to it, and of arresting symptoms at the onset.

External The application of heat to the surface has been means; heat. largely tried. The hot bath alone, or with mustard or salt, &c., the vapour bath and the hot-air bath, have been principally employed. It appears to be the uniform experience of the profession, that in collapse these means are but of little value.

In the earlier stage of the disease, or when it is less severe, and the pulse indicates some degree of power, the hot and vapour bath may allay cramps, and prove grateful to the patient; but when the nervous and vascular functions are greatly depressed, and the surface cold and clammy, although heat may be imparted to the body, it rarely excites reaction in the system itself; on the contrary, it is oppressive to the patient, and increases the exhaustion.

It is not found by experience that the degree of reaction, which may be sometimes excited by such means, and by internal stimuli, is of any permanent advantage, nor was this to be anticipated from a consideration of the conditions of the system in the collapse of cholera. The state of the blood, and the extreme depression of the nervous power, necessarily render abortive such attempts at restoring the circulation. The whole tendency of the evidence yet acquired from the treatment of this stage, is towards a more restricted use of powerful excitants of the kind alluded to.

Cold affusion. On the Continent, in the former, and in the last epidemic, cold affusion was highly spoken of as a means of producing reaction. The patient was placed in a warm hip-bath, and cold water poured or thrown over the head, back, and chest. This was done quickly, and the patient then placed between two warm blankets. If the first application was followed by any improvement, the operation was repeated every three or four hours. The results appear to have been, on the whole, more satisfactory than from the hot bath.

The wet sheet. The "wet sheet envelope" was more commonly used in this country. The effect varied according to the state of the patient; and in the milder cases it favoured re-

action, but when the disease was severe, it was useless or injurious. The sweating caused by it added to the exhaustion, and had no influence in arresting the intestinal discharges.

In the milder cases, stimulating epithems of mustard and turpentine were of some use in relieving local symptoms, and for obviating nervous depression. Frictions, chloroform liniments, and warm fomentations allayed the cramps. In severer cases these means were quite ineffective.

Two cases are reported in which oxygen was inhaled in the stage of collapse with asphyxia, at the same time galvanism was employed to stimulate the respiratory function. The effects upon the circulation and respiration were such as might have been anticipated. The heart's action was, for the time, increased, and there were slight symptoms of general reaction, but no permanently-favourable influence was exercised, by these means, upon the course of the disease in the majority of cases.

This method of treatment was not much investigated during the last epidemic. The results were, as in 1832—33, generally unfavourable. Its nature, however, cannot be determined by statistics collected from various sources. The operation in all its details is a delicate one, and requires not only a careful discrimination of the cases to which it is applicable, but also an exact attention to the physical characters and composition of the fluid to be injected, and other collateral circumstances. Until these points receive greater elucidation, the results obtained can form no sound basis for an opinion respecting its merits.

The arrest of the function of absorption in the stage of collapse, and the consequent inertness of all medicines administered by the stomach at this period, naturally led to the suggestion of injecting them into the veins. The trials which have been made in this way are too few to admit of any deduction from them. In the former epidemic, laudanum, camphor, quinine, &c., were used unsuccessfully

From the discovery of the circulation to the present time various experiments of this kind have been instituted by different observers, which prove that medicinal substances can thus with safety be employed.

Of those who resist the fatal influence of the cholera poison in the stage of collapse, a large number yet fall victims to its protracted effects. Probably half the cases fatal after the first forty-eight hours of severe symptoms belong to this category.

Treatment of imperfect re-action.

We may, therefore, infer that much care is required during the early period of reaction, since circumstances, in themselves apparently trifling, may at this time determine the issue. A strict observance of the horizontal posture, moderate external warmth, stimulating applications to the extremities, region of the heart, or epigastrium, and the administration internally of diffusible stimuli in small quantities, with the free use of ice, cold water, and other diluents, appear to constitute the principal part of the treatment as far as it is yet determined.

The cerebral oppression and delirium of this period are probably due to exhaustion and the asphyctic condition of the capillary circulation, the blood being imperfectly aërated and deficient in water. The state of the brain which gives rise to these symptoms is to be distinguished from that which occurs at a later stage from retention of the urinary excretions. It is not benefited by local or general depletion, but is slowly removed as the circulation is re-established.

Treatment of "consecutive fever" or uraemia.

The fruitless search after specifics which occupies the mind in the preceding stages of the disease is now tacitly given up, and we are content to be guided by the pathological indications, as in other diseases. But whilst our attention is directed to the local and more tangible complications, it must not be forgotten that they are the effects of an agent which has remarkably depressed

they are the effects of an agent which has remarkably depressed the whole of the vital powers, and, consequently, rendered necessary modifications in the activity of the measures otherwise adapted to their removal.

The next important indication for treatment is the depuration of the blood from the urinary excretions. The conditions following collapse are favourable to their retention, viz. the blood is of high specific gravity, the circulation is embarrassed, and lesions of the kidneys are frequent.

The treatment must include such means as allay irritability of the stomach and promote absorption, restore circulation, and remove the lesions of the kidneys. Of these may be enumerated the use of ice and cold water, effervescing draughts, or weak solutions of the alkaline salts, the warm bath, emollient enemata, dry-cupping to the loins, and counter-irritation over the stomach.

Stimuli are sometimes necessary for the purpose of counteracting the nervous depression, but their employment, whilst the system is poisoned with urea, obviously requires careful regulation. Stimulating diuretics, as squills, cantharides, &c., are not adapted to the great majority of the cases. The spontaneous occurrence of profuse diuresis, when the intestinal mucous membrane recovers its functions, shows that the matters to be excreted are, in themselves, sufficient to stimulate the kidneys, if the conditions favourable to their action be present.

Occasional doses of mercurials, followed by laxatives, were found to be useful, probably by quickening the circulation through the liver and promoting absorption, as well as depurating the blood, but their use was necessarily limited, whilst the excretive

function of the kidneys was defective.

During the consecutive fever, the intestinal tract is frequently, in different parts, the seat of persistent congestion with diptheritic inflammation and ulceration. The termination of the ileum is most commonly thus affected, and is the frequent source of the bloody stools observed during reaction. In some rare instances, limited sloughs of the mucous membrane result from effusion of blood into the submucous tissue. Such have been observed in the large intestine and in the stomach. The character of the local action indicates a want of reparative power.

The treatment of these lesions is probably that generally employed with success in other forms of gastro-enterite, such as gentle support, ammonia, serpentaria, and a moderate allowance

of wine.

Such is an abstract, as copious as I could render it, of Dr. Gull's ably-conducted portion of the Report to the College of Physicians.

EPIDEMIC CHOLERA IN CALCUTTA. — There is no branch of medical inquiry more interesting, nor any more important, than the history of epidemics, nor one, the accurate observation of which would prove more useful to physicians, in every climate. How true it is in our day, as in that of Sydenham, that no one has treated this great question in proportion to the dignity

of the subject.

An accurate history of epidemies in Bengal would prove of great value, as enabling us to trace their connection with changes of climate and condition of the surrounding localities, or with the social condition of the people, European and native. Had such a history existed, it would have helped to an earlier establishment of general principles, and to a more rational plan of treating the local fevers especially; for, though all the epidemic fevers within my personal recollection differed from the ordinary endemics of the country, still, there was in most of them so much of the savour of the soil (if I may be allowed the expression), as to render a knowledge of their history and treatment a matter of interest and importance.

It is these sweeping epidemics, aggravated by endemial and social influences, that swell the bills of mortality in our Eastern Metropolis, and that did so in former times more especially, to so frightful an extent: in truth, endemics are very often the parent-

stock upon which epidemics are engrafted.

Under a system of local sanitary improvement, such as has for years past, at the original instance of the author of this work, been established in Calcutta, the endemic sources of disease will be gradually diminished, and along with them the force of epidemics may also be reduced, so as greatly to improve the health and enhance the value of European and native life, within that rising emporium. Let want, filth, crowding, and misery be removed, and epidemics will have lost their chief power. It is these and other well-known sanitary defects that everywhere, but especially under tropical influences, give such destructive power to endemic and epidemic agencies, and that gave rise to the observation of Cabanis—that the effects of climate are not the same with the rich as with the poor.

CHOLERA FEVER OF 1834.—As bearing on the subject of this article, I would briefly mention a form of epidemic fever which visited Calcutta in 1834, and which we termed the cholera fever.

It arose in May—the worst of the cholera months—and continued through that and the following month, disappearing only on the full establishment of the rainy season.

Its symptoms were, heat and dryness of skin, frequent soft pulse, clammy white tongue, and generally a slight diarrhœa.

So mild a fever I seldom recollect to have seen; yet it was most dangerous to treat—the irritable condition of the bowels, which formed so prominent a symptom of the epidemic, being readily aggravated into fatal cholera whenever purgatives of a saline or drastic nature were exhibited.

At first the peculiar character of the fever was not known; and I remember hearing at the time that several deaths occurred where strong purgatives had been exhibited over night; indeed, one such case came accidentally under my observation. It was that of a gentleman, the patient of a professional friend, into whose house I was called in haste on passing at 6 A.M. His countenance and voice at once testified to the collapse of cholera, and to his approaching end. He had been feverish during the previous day, and had taken a drachm of compound jalap powder at 9 o'clock of the previous night.

Others fell victims, in like manner, to the hasty exhibition of purgatives; but within a few days, and when the nature of the epidemic became generally known, it proved a manageable form of disease, and after that few deaths occurred.

A stout mercantile gentleman, whom I treated with a moderate dose of castor oil, had fifteen stools between 8 and 11 A.M.—the former having been the hour of exhibition. Had the above dose, small as it was, been administered at bed-time, the result to the patient would have been fatal.

Not only was the peculiar fever here noticed prone to merge into true cholera, but I had to remark on three several occasions in Calcutta, that, when the cholera poison was present in a concentrated form, we were for the time precluded from treating the cases of ordinary remittent fever of Europeans in our usual manner. Moderate blood-lettings were now liable to be followed by extraordinary collapse, and purgatives could not be administered over night, lest the fever should be changed by morning into a deadly cholera. The state of the system here indicated, showed that the fevers, and also the diarrheas of the country did, under epidemic or other peculiar influences, merge into the true cholera, the cholera poison being at the moment the more powerful influence. In times like these, we found it difficult to persuade ourselves that the endemic and the epidemic diseases were absolutely "caused by different poisons."

No fever of the nature of the epidemic just noticed had been known in Calcutta previously; nor has it recurred since the year specified. The treatment I adopted was by mild doses of calomel and James's powder, with a grain of opium, exhibited at bed-time,

and an occasional mild aperient in the forenoon.

All the epidemics of which I could trace any record, occurred in the hot months of March, April, or May; and thus they correspond, as to time of occurrence, with almost all the known European epidemics. Andral states, that of fifty epidemics known in Europe, thirty-six occurred in summer, twelve in autumn, one in winter, and one in spring.

INFLUENCE OF HIGH TEMPERATURE.—Independently of the proved influence of heat in promoting the existence and continuance of epidemic cholera, it is remarkable that this disease should have as frequently commenced in Bengal at the season during which ventilation is most free and effective. In Europe we are told that a free ventilation is the most efficient means of destroying the cholera poison; yet, in Bengal, the epidemic has been seen very generally to commence about the middle of March, when the South-west Monsoon comes sweeping with power and purity over the plains, fresh off the Bay of Bengal. The freedom of ventilation at all seasons in India, will account, I think, for the circumstance, that so little suspicion of contagion has at any time arisen in that country-from the first origin of the epidemic to the present day. It is a fact that, "of between 250 and 300 medical officers in Bengal, most of whom saw the disease largely, only three were attacked, and only one attack was fatal." In Calcutta, as in London, the highest range of temperature was that of greatest prevalence of cholera. "The reader of the meteorological tables," says Dr. Farr, "will not fail to observe that in the thirty-sixth

week of 1854, when the cholera raged, and the deaths from a causes rose to their maximum (3413), the average daily range temperature was 30°.9, consequently the greatest in the fifty-tweeks; the highest temperature of the week was 81°.2, the lower was 43°.1, therefore the entire range was 38°.1." It is worthy remark, says the same authority, that in the hot season of 184 when the cholera epidemic acquired great force about the Indusummer cholera and diarrhæa prevailed with great violence England.

THE INFLUENCE OF SEASON.—The following tables a interesting, as exhibiting the influence of season on sickness at mortality from epidemic cholera amongst the inhabitants Calcutta, native and European.

Mortality by epidemic cholera of Hindus and Mahomeda, for each month during seven years, from 1832 to 1838 inclusive:

Months.	Hindus.	Maho- medans.	Total.		REMAR	KS.	
In 7 Januarys	 572	125	696	Mean	temperature,	66°.2.	D
7 Februarys	 620	196	816	,,	"	69°.8.	
7 Marchs	 1873	439	2312	19	"	80°.0.	
" 7 Aprils	 2707	482	3189	99	"	85°.4.	D
" 7 Mays	 2170	464	2634	1,	**	85°.7	R
7 Junes	 615	217	832	>>	"	83°-7.	R
7 Julys	 914	133	1047	"	"	81°.8.	R
" 7 Augusts	 806	146	952	7,	**	82°.0.	R
" 7 Septembers	 785	121	906	19	"	82°.8.	R
" 7 Octobers	 1030	198	1228	111	"	79°.%	D
" 7 Novembers	 1687	230	1917	"	"	74°.2.	Di
,, 7 Decembers	 1425	161	1586	"	"	66°.6.	Dı
Total	15204	2911	18115				

It thus appears, that while the proportion between the Hind and Mahomedan inhabitants of Calcutta is but  $2\frac{1}{2}$  to 1, the ratio mortality amongst the former is, to that of the latter, as  $5\frac{1}{2}$  to this great difference in favour of the Mahomedans can only ascribed to their better habits of life—to their better diet, clothin habitations—comparatively.

The influence of the highest temperature of the year on endemic cholera is exhibited in a still more striking manner on the sickness and mortality of Europeans; for, by the following table it appears that for the twelve quarters, ending 30th June, the

were admitted into the General Hospital of Calcutta, 492 cases of this disease, of whom 220 died; while, for the remaining 36 quarters, there were but 311 admissions, and 152 deaths. This is a remarkable fact.

Table of admissions and deaths of Europeans from cholera into the General Hospital of Calcutta, for each month during twelve years:—

Range of Observation.	Total Admissions.	Total Deaths.
In 12 Januarys	28	11
" 12 Februarys		5
" 12 Marchs	56	34
" 12 Aprils	86	57
" 12 Mays	268	125
" 12 Junes	138	38
" 12 Julys	32	6
" 12 Augusts	37	18
,, 12 Septembers	10	5
,, 12 Octobers		27
" 12 Novembers	59	32
" 12 Decembers	32	14
Grand Total	803	372

It should be observed, that the patients admitted into the European General Hospital of Calcutta are—soldiers belonging to detachments—invalid soldiers from the Upper Provinces—soldiers forming the garrison of Fort William—European townsmen, and men belonging to the shipping in the river. With exception of soldiers on duty in garrison, all the Europeans sent into the General Hospital are exempted from that salutary regulation and discipline which prescribes that every man shall come under medical control the instant he falls sick; and hence, doubtless, a great increase of deaths from cholera, as well as from all other acute diseases within this institution.

The result of an extensive series of observations, ranging over twenty years, shows that, in the Presidency of Bengal, the annual mortality of European soldiers, from epidemic cholera, is about 1.15 per cent., while of the Bengal Sepoy army it is but 0.22. This remarkable difference—so opposite to the popular belief—is worthy our careful investigation. Here we perceive that the boasted strength of the European does not serve him. On the contrary—in his power to resist this terrible scourge, he stands

vastly below his comparatively feeble native comrade. Why is this? We can only refer it to the habits of life of the British soldier—to his habitual indulgence in an over-full animal diet, and to his abuse of ardent spirits—thus keeping the digestive, nervous, and vascular systems in a constant state of over-excitement. He is thus rendered more prone to collapse of the vital powers; for, whatever lowers these last, reduces the general nervous energy, deteriorates the blood, or interferes with the processes of digestion and nutrition, will dispose to attacks of cholera.

PREDISPOSING CAUSES.—Some of the predisposing causes, as already stated, have been ascertained with some degree of accuracy—but not yet the exciting cause. Here, despite our science, all is mystery—alike in the fleet on the ocean, or in the army on the march—all is inscrutable. Why does the epidemic come in one year and at one season, and not for years or seasons thereafter? These and such questions no one can answer.

The number of instances which I observed in Calcutta of Epsom or Seidlitz salts leading to profuse serous diarrhea, and subsequent fatal cholera, is a circumstance worthy of notice. The dose was always taken in the early morning, and the patient has gone through his ordinary avocations throughout the day, mistaking for the operation of a saline aperient what was in reality the induction of choleraic diarrhea, true cholera, and deadly collapse.

In the end of March, 1833, I was requested to visit the head partner of a large house of business—a healthy and powerful man. The note, which was not expressed with any urgency, was delivered to me at 5 p.m., near to the house of the patient, so that I was with him within a few minutes of its being written. I found the gentleman at his desk, at work. His countenance and voice at once expressed fatal collapse, and he died at 8 o'clock that evening. He stated that at 5 of that morning he had taken a dose of seidlitz—his health previously having been quite good.

PREMONITORY DIARRHŒA.—I have never attended a case of cholera in which premonitory diarrhæa, of longer or shorter duration, had not been plainly manifest; but I have seen people—villagers and camp followers—struck down with sudden death on the line of march during April and May, in the deadly jungles of Gondwana, and there the relatives and attendants declared emphatically that there had been no antecedent purging or vomiting whatever. All I observed personally was, that the persons lay dead by the road-side, and that there were no evidences about them of there having been any diarrhæa or vomiting. The cholera was raging at the time in our moving camp, of which I had charge.

A gentleman of the civil service, describing to me the fatal out-

break of cholera, which visited Calcutta in 1842, says :- "It some time in March took quite a new character,-those attacked had not the usual purging and vomiting, but it came on with a sort of faint sinking-the strength was, without outward effort or apparent cause, totally prostrated, and the patient died without suffering, as if labouring under the effect of poison." In the terrible epidemic which seized on Her Majesty's 86th Regiment, at Kurrachee, in June, 1846, and which destroyed, within sixteen days, 238 men out of 399 who were attacked, it was noted that "the peculiarity in many cases was, that there was neither vomiting nor purging, and in the latter cases, by no means profuse. Also, among some of the first cases, before the natural hue had been replaced by the livid look so characteristic of the disease, the pulse at the wrist was almost imperceptible, the eyes turned up, and the voice hollow and feeble. In an hour after, such cases were dead." Others are described as attacked of a sudden, "and the collapse preceding the vomiting and purging of serous fluid." Dr. Lindsay, of Perth, declares that "cases of cholera undoubtedly occur without any rice-water evacuations: nay, it does not appear essential to cholera that there should be intestinal evacuations of any kind."

In the onset of many epidemics, we learn that sudden deaths occur, apparently from the concentration of the poison. Men are thus struck down at the outbreak of yellow fever and plague, as well as in cholera. During the onset of epidemic plague, almost all who are attacked with it perish—death occurring in from twenty-four to forty-eight hours; while, after a little time, death is deferred to the fourth, sixth, fourteenth, and even to the twentieth day—many cases recovering health: at length the pestilence

loses its malignity altogether.

e of

20000

5005

SITE

sub-

The

t has

Stale-

the

IDSK.

bend

erāal

THE R.

the

ound

that

en a

ed a

Orter

ple-

th on

les ei

apping-

Detime

about

holem

harge

al out-

PATHOLOGY.—The first circumstance that will strike the observer in cholera is, the astonishing and fatuous prostration of the mental sensibilities, on all questions relating to the disease, and to the condition of individual danger belonging to it. The person who, before the seizure, was all anxiety, alacrity, and alertness, on the subject of cholera, both as affecting himself and others, is now fatally apathetic; the feelings of hope, sympathy, and fear having vanished, and that of self-preservation even being deadened by the morbid impression of the poison on the nervous system. With all this, the faculties of the mind are in perfect integrity for the arrangement of business, and the disposal of property—the patient making his will with a cold indifferent justice—the mind being as benumbed and prostrated as the body.

The first impression of the morbific cause would appear to be made on the system of the organic nerves and their functions; for, almost immediately, we have the vital actions of circulation, respiration, the generation of heat, secretion and absorption, conspicusly depressed; and that probably, amongst other causes, thrown some unknown changes in the proportion of oxygen, or in electric condition of the atmosphere. We have here, to use language of John Hunter, the death of the blood. This condit of the circulating fluid produces torpor of the brain: the voice veiled, as when great hæmorrhages occur, and very soon cyand becomes complete, the complexion assuming a layender hue.

Fevers, both symptomatic and idiopathic, are referred by Billing, to the loss of the functions of the nervous centres, subsequently to that of the organs depending on them.

In all the severer visitations of this epidemic, we have syntoms at the very onset of the disease which, as it appears to can be referred to no other circumstance than this primary in ence of the poison of cholera on the great nervous centres—give the death-blow to the organic functions. All the secretions suppressed; and in their place we have a general and exhaust transudation from the bowels and skin. In short, all the grunctions of life are first altered and perverted, and then copletely and fatally arrested. The very stomach and bowels see dead-drained of their serum sanguinis;—all now tends, in fit of death.

The shock to the nervous system, paralysing its function causes the blood to be detained also in the great trunks, and the functions of secretion and excretion become additionally suppressed—the power of generating heat to be lost—and eventuate the death of the blood, as John Hunter termed it.

TREATMENT.—It is impossible to say anything satisfact on the treatment of a disease which would seem every year to come a subject of greater difficulty to medicine. Like the Pe lence of the fifth century before the Christian era, described Dr. Arnold, this epidemic has in a manner travelled up and do over the habitable world, unaffected by human endeavours, retuing often to the same place after a certain interval; pausing sor times in its fury, and appearing to stop, but again breaking out some point or other within its range, till, at the end of its appoint period, it disappears—we dare not say—altogether.

Dr. Elliotson declares that "if, as respects this country, all patients had been left alone, the mortality would have been me the same as it has been. We are not in the least more advance as to the proper remedies than we were when the first case cholera occurred."

Certainly, every year makes us more and more familiar with phenomena of the disease; but this circumstance has in no v improved our acquaintance with the means of checking its progr

and opium; and where, nevertheless, the disease still proce the repeated use of smaller doses of calomel with highly sible stimulants was persevered in, so as to rouse the ne and vascular actions.

It seems admitted in actual practice, in all portions of the E Empire, and by the great body of the medical profession during the premonitory stage, or that of simple diarrhea, opi the sheet anchor of reliance. Some give this remedy alone—with chalk and aromatics—some with calomel; but all give of in some form or combination, thus following still the original tice in India. Opium thus used acts at once as a cure of diar and of the earlier stage of cholera, and as a prevention of coming and fatal collapse.

After all our experience, in all parts of the world, the treat by calomel and opium at the commencement of the disease calomel with diffusible stimulants in its subsequent stage originally practised in the East, is that treatment which has the most steadily and the longest approved, by the profession

In the report on the results of different methods of treat pursued in Epidemic Cholera, and addressed to the Presidenthe General Board of Health, by the Treatment Committee Medical Council, we find the following summary:—

"The evidence of these Tables condemns the eliminant

ment altogether, as a principle of practice.

It testifies against the stimulant principle, excepting as source in extreme cases.

It displays a decided advantage in the alterative prin especially as carried out by calomel and opium; and it sh still superior advantage in the astringent principle, as at through the means of chalk and opium—the *general* per-ce of deaths following each plan of treatment being—

Of Eliminants			71.7	per cent.
" Stimulants			54.	,,
" Alteratives, ca			36.2	
" Astringents, c	halk and op	ium .	20.3	"

In order to judge correctly of the value of this evidence, necessary to examine, as far as may be possible, the degree severity of the cases brought beneath this test. The only near the command (on the present occasion at least) to make examination, is to consider the relative proportion which the of collapse bear to the number of deaths of their own clarespectively. Examining, therefore, the collapse cases with number of deaths, we find that calomel and opium stands him the scale of success, and the order of preference appear follows:—

Calomel and opium .		59.2	per cent.
Calomel (larger doses)	4	60.9	
Salines		62.9	
Chalk and opium .		63.2	
Calomel (small doses)		73.9	
Castor oil .		77.6	
Sulphuric acid .		78.9	

According to this result, the superior success of calomel and opium in severer cases, appears a distinct fact elicited by the present inquiry. It is accompanied by other facts, viz.,—the relative advantages of those other modes of treatment which follow in their order of success. This order marks the use of calomel in small doses, of castor oil, and of sulphuric acid, as actually to be deprecated in severe cases.

Chalk and opium, as shown above, stands at the head of the list, in the general per-centages, both in hospitals and in private practice; but in the comparison of the collapse cases with the number of deaths, the average declines to the fourth rank."

The names appended to the report, of which the above is an extract, are:—John Ayrton Paris, President of the College of Physicians; James Alderson; Benjamin Guy Babington; Alex. Tweedie; Nathaniel Bagshaw Ward; all Fellows of the said College. In the "Report on the treatment of Epidemic Cholera, in the provinces throughout England and Scotland, in 1854, being Supplementary to the Metropolitan Report, addressed to the President of the General Board of Health, by the Treatment Committee," all the above facts are substantially corroborated.

I have only to state, in conclusion, that in cholera, in common with the last stages of violent fevers and dysenteries—as, in fact, in all diseases of great exhaustion—the patient will always owe much to the horizontal position, and to careful and unremitting nursing. The most careful nursing, and the most attentive watching of the patient, are both of the utmost importance in this disease; and so easily is the balance of circulation fatally overturned, that a strict attention to the recumbent posture is absolutely necessary to success. In no other diseases are these simple matters of so great importance to be attended to; and in the disease under special notice, I have seen many a life apparently lost from inattention to them.

For the following Tables 1 am indebted to my friend Dr. Balfour, of the War Office. They exhibit the average sickness and mortality from Cholera in certain of our foreign possessions; and show by how much more fatal Epidemic Cholera has proved in the country of its birth than in any other.

TABLE I.

	-					
Periods Aggregate	Aggregate	_			Ratio per 1000.	1000.
of Strength. Observation.	Strength.		Admitted.	Died.	Admitted.	Died.
1830—46 98985	98985		171	75	1-7	ė
	93400		459	131	6.7	¥.
1817—46 26494	26494		None	7	*	P
1825—46 7678	22945 7678		None None			
1817—46 43248	43248	_	210	29	4.8	Ξ
1818—36 29344	154736 29344		356 None	127	<b>6</b> 1	άο
1817—36 86661	86661	_	None			
	21267	_	None			
1818—36 5908	8069 1	_	None None			
33	30515	_	898	88	<b>80</b>	1:1
<b>3</b> :	42978		88	22	18.3	•
1820 45 117588	117588 88390		217	1952	2	2 =
	20087		::	88	::	5
		_		687		7

TABLE II.

				Ceylon		Mauritius	Jamaica		Canada	Nova Scotia and N. Brunswick .	Malta	Gibraltar	* · · · *	Un Cin	Troops generally . *	ed on	n. = = = .	Dragoon guards and Dragoons		
1832	1829	1825	1820	1819	1820	1819	1850	1834	1832	1834	1837	1834	1834	1833	1832	1834	1833	1832		Periods of Observation.
2056	2279	1448	2703	2533	1395	1652	756	2588	2909	2071	2159	3034	38587	42614	45583	6261	6379	6408	St	rength.
344	48	111	42	236	155	113	51	97	259	210	209	459	154	223	886	25	32	114		Cases of cholera.
107	14	31	12	89	12	20	29	- 33	94	59	47	131	61	71	233	11	14	29		Deaths.
167-3	21.1	76-7	15.5	93-2	1111-1	68.5	67-5	37.5	89.	101-4	96-8	151.3	4.	5.2	19-4	4.	5.	17.8	Cases.	Ratio per 10 of strength
52.	6.1	21.4	4.4	35.1	8.6	12.1	38.4	12.7	32.3	28.5	21.8	43.2	1.6	1.7	5.1	1.7	2.2	4.5	Deaths.	er 1000 ingth.
							10 ,, 17-6		,	,	,		,	:	=		:	ii ii	attendance of	Proportion of deaths to

## THE YELLOW OR HÆMOGASTRIC FEVER.

GENERAL AND TOPOGRAPHICAL OBSERVATION -In most, if not in all, European countries, we find that fevers the continued, typhoid, remittent and intermittent characters, pr vail, with bilious remittents, in the autumnal seasons. In Egy and Turkey, again, we find the plague; on the West Coast Africa, and in the West Indies, along with remittents, we ha occasional outbreaks of destructive epidemic yellow fever; in the East Indies, formidable bilious remittents.

All these fevers, while they take their origin from peculiariti of soil and general climate, have certain features of resemblance and yet many wide and essential differences and distinctions: the all unite under the common generic FEVER-only to demonstra to us, for the millionth time, the surpassing difficulties of the

great subject.

The yellow fever-"the hurricane of the human frame," as Blan terms it—would be as difficult to import into the climate of En land, as would be the typhus of our crowded cities into those Jamaica or Bengal; such are the peculiar characters imparted local circumstances to those pervading and dangerous disease The yellow fever, for instance, is altogether arrested by a modera degree of cold, while plague is nipped by cold and dryness,

also by great heat and dryness of atmosphere.

William Fergusson, speaking of the expedition to St. Doming in 1796, "as one of the survivors of that war," says :- "Our hea quarters were the town and its adjunct Brizoton, as pestiferous any in the world; and here we had constant vellow fever in all fury. At the distance of a mile or two, on the ascent up the cou try, stood our first post of Torgean, where the vellow fever a peared to break off into a milder type of remittent. Higher t was the post of Grenier, where concentrated remittent was ran and milder intermittent with dysentery, prevailed; and high still, was Fourmier, where remittent was unknown, intermitte uncommon, but phagedenic ulcers so frequent, as to constitu a most formidable type of disease; and higher still were t mountains above L'Arkahave, of greater elevation than any

them, far off, but within sight, low down, in what was called the bight of Leogane, where a British detachment had always enjoyed absolute European health, only it might be called better, because

the climate was more agreeable than in higher latitudes.

Here were the separate regions or zones of intertropical health mapped out to our view as distinctly as if it had been done by the draughtsman. Taking Port au Prince for the point of departure, we could pass from one station to the other, and with a thermometer, might have accurately noted the locale of disease according to the descending scale, without asking a question among the troops who held the posts. It was just as impossible, or more so, to carry a yellow fever up the hill to a post in sight, as it would have been to escape, had they been brought down and located amidst the swamps of Port au Prince." . . "Even here at home, in certain peculiar seasons of heat and drought, vellow fever may not have been entirely unknown. The year 1807 was one of those, when two soldiers of a militia regiment, in the garrison of Sheerness, in Kent, died of a fever under my own inspection, whose cases might have passed muster for the endemic of St. Domingo." The opinion here suggested accords with that of Frank, who regarded yellow fever as an intense variety of the worst form of the autumnal bilious remittents of temperate regions. Dr. Dickson, speaking of the destructive epidemic vellow fever, of Marieglante, in 1808, says, that "the different types of fever were CONVERTED into each other, of the worst and most aggravated species he ever witnessed. Yellow fever, in the continued form, others with comatose remittents and intermittents, the exacerbations of which were so violent as to carry off a patient in two or three paroxysms, while others sunk into a low protracted character of fever resembling typhus."

The yellow fever, like the remittents and intermittents of hot climates, would seem to take its origin from terrestrial exhalations—with this further similarity also, that yellow fever may, under certain favourable conditions, be propagated as remittents have, I believe, truly been said to have been, by contagion. This manner of regarding the subject, as stated by Dr. Alison, of Edinburgh, is the only means of reconciling, not the conflicting and valueless opinions, but the apparently contradictory facts presented during the various epidemics of yellow fever, recorded by so many able observers of those calamities. There is, as stated, reason to conclude that remittents may, like the continued fevers of temperate climates, be occasionally propagated by contagion; and of all the accessories to such condition, crowding of the sick

would appear the most efficient.

Another circumstance of preliminary importance and interest in the natural history of yellow fever is, that "whether it spreads by contagion within certain limits, or not, it will in all probability be confined to a tainted district, even in that climate; and that of careful inquiry, in almost every such case, it will appear that there are boundaries at no great distance from the spots first affected beyond which the disease has not formerly spread, and need hardly now be apprehended."\* As examples, the limits set in Philadelphia, half a century ago, and more recently, the Neutra Ground, Gibraltar, are adduced. These facts impress as a necessity, the principle of immediately abandoning a place when yellow fever exists, and not returning to the locality originally occupied until the disease shall have for some time altogether disappeared.

But to continue the general comparison:—Robert Jackso "considers endemic fever, whether it appears in the torrid or th temperate zones of the earth, to be radically one disease."

"The forms of endemic fever are in fact so extremely diversified the symptoms so opposite in their nature to each other on man occasions, as to produce the distinctions of inflammatory, putric nervous, bilious, simple, complicated, mild, malignant, &c., from the operation of a cause that is distinctly and demonstrably one.

. . "The history of differences or distinctions constitute what is called nosology, a study of much importance to medical science; but, in so far as respects endemic fever, a study of more perplexity and loss than utility, as labouring to discriminate where there is no radical difference, or to bring the products of chance under a constituted law of nature."

Of the Gastric or Bilious Remittent Fever, this celebrated write says:—"Bilious remittent fever is one of the most common, and one of the most important of the febrile forms which occur is armies, and even in civil communities. It belongs to all countries and it is endemic in the West Indies at all seasons of the year especially among native subjects, and those who have been so long resident as to be acclimated. It owes its origin to emanation from the soil, more generally or more partially diffused." Beside its own febrile mode of final termination, favourable or fatal a may be, "it is prone to assume a new form and to acquire a new quality so as to constitute a new disease which has its own law of movement. It thus assumes the contagious character, or is readily convertible into contagious fever in protracted campaigns in temperate latitudes."

This doctrine, held so long ago by Jackson and others, in respect of remittent and continued fevers, has recently been verably expounded, as applicable to yellow fever also, by Dr. Alison

<sup>\*</sup> British and Foreign Med. and Chir. Review for April, 1854.

of Edinburgh, in the British and Foreign Medico-Chirurgical Review.

In an article evincing an extended research and much ability, in this review for January, 1848, and in which the author offers a just tribute to the profound knowledge of fever possessed by Robert Jackson, he says:-" It is to be remembered also that very considerable differences in type do not always disprove the existence of a single and common cause. We should have some difficulty in believing that the mild scarlatina simplex had any alliance with the malignant sore throat with little or no eruption, if repeated observation had not demonstrated the transition stages. What more unlike than a mild quotidian and the pernicious intermittent which kills in a single paroxysm; or a slight case of true cholera, and the fatal and uncontrollable disease which devastates India. And if any one had seen only the milder forms of the remittent fever, and had had no opportunity of tracing up the several grades, he might well believe, when he saw suddenly the severest variety, that he had before him a distinct affection. A cause is not unalterable; a morbid poison has not always the same concentration: it is ever varying in its intensity, and in the action it exerts on the system, more or less susceptible of it. And this has long been insisted on by writers on the endemic fevers of the tropics." After quoting Jackson's declaration that a malarious disease "is subject to changes at shorter or longer, but unascertained intervals, without any visible material change in the qualities of the localities or climate," the reviewer justly observes that, "In this absence of a malarious disease for a term of years, a resemblance is perceived with the absence of epidemics for a longer or shorter period."

Another remarkable and important fact is, that while the coloured population and acclimated Europeans suffer only from remittents, or from the very milder forms of yellow fever, the unacclimated European is everywhere and at all times the victim of

the most virulent and fatal form of this epidemic.

It may be said, in addition, that a first attack of yellow fever gives a certain degree of immunity from a second, and likewise that a previous residence in a warm climate greatly lessens the

susceptibility to yellow fever.

Topographically then, yellow fever may be defined—a disease not proved as yet to be a disease, sui generis—endemic only in low districts on the sea-coast, but, under certain circumstances, sporadic in other places; never appearing beyond the 48° of north latitude, or above the elevation of 2,500 feet above the level of the sea, nor without a temperature of at least 72°, or more, promoting its production and propagation; depending, in part, on

causes yet unknown, but in circumstances favourable to its tension, capable of being propagated by contagion.\*

Mason Good prefers the trivial name yellow to that of palue fever, which is too limited to express its source in every instance.

SYMPTOMS AND PROGRESS.—The yellow, like all of fevers, makes its invasion with more or less severity according age, sex, constitution, previous habits of life, and character of prevailing epidemic; but the following symptoms will be found characterize the slighter, the severe, and the more fatal seizures

It is ushered in suddenly and generally, as in cholera, in a night or early morning, by sense of coldness, a rigor, or acts shivering, followed by a vascular reaction proportionate to amount and duration of the previous congestive cold stage; he and dryness of skin follow, with pain of the loins and limbs, her ache, pain of the eye-balls, which are suffused, and of a glos drunken-like aspect; the tongue is loaded, the edges red—there nausea, with sense of rawness of the fauces and æsophagus, a uneasiness at the epigastrium. As in the remittent fever, the symptoms may not extend beyond twenty-four, or thirty-six hor terminating then, under proper means of cure, in the restoration health.

But this fair prospect of recovery too often proves only a decetive lull before the coming storm: nausea, eructations, faintne vomitings, epigastric distress set in, with the ejection from a stomach of a fluid at first clear, and soon to become dirty, brown and eventually the true black vomit. The complexion now becomes, more or less generally, of a lemon yellow, which, as a disease proceeds, extends rapidly over the entire surface of a body. All the symptoms proceed from bad to worse, accompaniby a bloated, desponding, anxious, and inquiring countenance, cessant and distressing jactitation—ending rapidly in exhausting and death.

As with scarlatina, one person will get the fever in its mild while another will, at the same time and place, contract it in most grave form.

When again the epidemic is of a still more severe or fatal te dency, and where the subject is, from constitution, or previous habits of life, peculiarly susceptible to the impression of the mebid cause, then all the symptoms enumerated become fearful urgent; and the danger is imminent from the very onset of t fever.

In youthful and plethoric Europeans the symptoms are now exceeding violence and danger. The heat is of a burning pu

<sup>\*</sup> Vide British and Foreign Med. Chir. Review for April, 1854.

gency, while the skin is dry and constricted—the headache intense -the countenance bloated, oppressed, or agitated by the most fearful apprehensions, or by approaching delirium; the bowels are constipated, the tongue red, clean, and tremulous, indicating increase of gastric and intestinal irritation, and consequent increase of danger-the urine scanty and surcharged-all the secretions are suppressed; there is distension of both hypochondria, with pain, burning heat, and anguish of the epigastrium-eructations, hiccough, and vomiting, at first of a clear, glairy fluid, presently altered to brown and dark matter like coffee-grounds-the quantity of fluid brought up far exceeding any that has been used in drink. In persons of a sanguine complexion especially, the countenance is apt to assume a livid, yellow, and putrid appearance, with black incrusted tongue, while, in most cases, the whole surface of the body becomes more or less yellow; and in this, the most severe and dangerous form of the disease, the patient may be carried off, exhausted in all his functions, so early as the second, but more generally on the third day. The yellowness of the skin may be general or partial; and even the black vomit may be said to be a very frequent, but by no means a constant occurrence.

In some instances, the ejections from the stomach do not commence until the fever has almost subsided, and then the retching and vomiting are uncontrollably urgent, while in others, cozing of black dissolved blood takes place from the mouth, throat, and rectum. The type of yellow fever is continued, though occasionally exhibiting a partial exacerbation towards night; remissions in the ordinary sense, can therefore hardly be said to occur in this disease. But there is a deceptive calm of which the inexperienced should have warning; for, too often, it is followed by a fearful storm of black vomit—ushered in by faintness, sinking of the pulse and temperature. This nearly hopeless condition may appear on any day from the third to the seventh, and sometimes, though rarely,

even later.

Stupor is found to vanish as the hamorrhages increase. This was remarked as universal in the epidemic yellow fever of Peru in 1854. Bleeding of the gums is stated to have attended almost every case of the disease, not excepting the mildest forms of it; and stupor disappeared as the black vomit increased; in other words—cerebral congestion was relieved by the hamorrhages.

But uniformity in the order and character of the symptoms must not be looked for in yellow fever; for all the best writers on the subject, whether recording their experiences in the West Indies, the west coast of Africa, or the south coast of Spain, are uniform and unanimous to the contrary effect; certain symptoms in certain epidemics, varying in their nature and in the time of their accession, while others, common to former visitations, are generally wanting in this. It is, doubtless, this circumstance which has led to scriptions of "two or three, or even four forms of the disease." The French Commissioners state that "the Barcelona fever is yellow fever of America, the same that we have in the Anti and at Cadiz;" but they add,—it is a Proteus clothed v different forms, and which offers such strange anomalies, both the slowness and rapidity of its march, in the combination, succession, and the degree of its phenomena, that it is impossi to assign to it a fixed and invariable course."\* One conditi however, would appear to be nearly uniform in the experience British practitioners; bile is absent, or nearly so, from the intinal and renal secretions.

Dr. Gillkrest states, that where neither yellowness nor hæm rhage supervene, the fatal issue may occur in the shortest tir "The countenance of the patient may not indicate great dang but he is observed to lie with his limbs uncovered, his head having over the bed; he becomes sensible when spoken to, and vassist to arrange the bed-clothes, which soon again become placed." The symptoms in these cases, though wanting in yellow complexion and black vomit, were all of the worst kin often ending in a "low, incessant, and monotonous wailing, tremely distressing to all within hearing." This distinguish medical officer adds, that in the last epidemic in Gibraltar, these cases where copious hæmorrhage took place from any of natural orifices, black vomit and suppression of urine were maless likely to occur."

Another deadly form of the fever, he says, was characterifrom the commencement by a countenance shrunk and decoposed—the face having a mottled ash-coloured appearance—look sullen—the eyes of a dull, red colour—the tongue red raw—the temperature below the natural standard: the scene, adds, is closed not unfrequently within forty-eight hours.

SEQUELÆ.—It is remarkable, considering the violence this disease that so few sequelæ are noted by authors on yel fever; and so trivial do they appear to Dr. Copland, that does not believe in their existence;" while Dr. Gillkrest is sil on his own experiences, and quotes with expressed doubts, observations of others.

THE BLACK VOMIT.—"The vomitings at the beginning says William Fergusson, "are always in themselves clear and cleathe white vomit of some writers;—" presently they become some what ropy; then brown, dark, and darker, till at last the attenuation."

<sup>\*</sup> British and Foreign Med. Chir. Review for January, 1848.

blood, escaping from its vessels, and mixing with the gastric

fluids, comes up under the form of black vomit."

Dr. R. La Roche, of Philadelphia, in his elaborate "Observations on the Black Vomit," says:—"The black vomit, being recognized to be blood acted upon by the acid contents of the stomach, we have no difficulty in perceiving that much of the difference it presents, in regard to its physical appearance, will depend on the manner in which the blood is effused into the stomach, whether drop by drop, or in a stream, and on the degree of acidity of the gastric secretion, or the quantity of serous fluid it meets with in that organ.

It has been found, for example, that whenever blood is exhaled therein in a quantity proportionate to these secretions, it exhibits a black colour, while the aqueous portion is limpid, or

clear green.

3. If there be a slight excess of blood, more than enough to neutralize the acid, instead of black, we have a nut-brown, a chocolate or reddish matter, and the watery portion, when filtered, is of a rum, brandy, or red colour.

4. If the hæmorrhage be great, the fluid presents all the characteristic marks of blood, either with or without admixture of

black vomit."

Dr. Hewson Bache, U. S., gives the following as the results of many microscopical examinations of the matter of black vomit:—

- 1. "Amorphous masses of coagulated mucus or serum, and the debris of blood-corpuscles, which gave the mass a red or brown colour.
- 2. Numerous and irregular masses, sometimes presenting an imperfect crystalline form (probably altered hœmatin), which were of different shades, from a reddish-brown to a deep black colour.
- 3. Red blood-corpuscles, in different stages of alteration, which did not correspond to the degree of acidity of the vomit.
- 4. Various forms of epithelial cells, as the squamous and conoidal. In one specimen, a perfect mucous crypt was observed.

5. Brown oil globules, some of which bristled with fine crystals

of margaric acid.

6. Besides these constant elements, there were some occasional ones, such as the sarcina ventriculi, and the torula, the former more frequently than the latter, and some inflammation corpuscles."

DIAGNOSIS.—The black vomit has been regarded by some authors as the only true diagnostic sign of yellow fever, and in a large proportion of fatal cases it certainly is present; but vomiting of matters, so nearly resembling those ejected from the stomach in

yellow fever, occurs as an occasional symptom in the fever other countries, hence the following conclusions may, I think deduced from the recorded observations of American and o authors:—

1. That yellow fever may, and does occur without blyomit.

2. That black vomit is not so exclusive and universal a feat

of yellow fever as some authors have supposed.

3. That black vomit, so called, has been witnessed occasi ally in the course of fevers of the East Indies, of the Medite nean, and even in fevers and other diseases within the Bri Isles; and it is stated to be "an usual symptom in the yel fevers of America, which are incontestably of marshy origin."\*

4. "That to regard black vomit as an invariable, and altoget exclusive feature of yellow fever, would be almost equivalent asserting that none have been affected with this disease, but so

as died in the stage of actual vomit.

 That in many yellow fever epidemics, in various countr numbers have died of this disease without the occurrence of bla vomit.

6. With rare exceptions, those who have recovered freelow fever have not presented the symptom of black vomit.

7. On the other hand, the matter of black vomit has v generally been found, on dissection in the stomach and bowels subjects, in whom no vomiting of black or other matter occur during life.

8. Black vomit constitutes, in the prevalence of an epider fever, the most unmistakeable sign of its being the yellow feve the black peculiar matter ejected during life, or discovered in stomach and bowels after death, leaving no further doubt as to nature of the disease.

9. On the other hand, black vomit was remarkably absent the fever of the Niger expedition, and in the Gibraltar epidemic

1814 it was seldom seen.

Where, again, the disease is mild in its character and progree the diagnosis becomes one of exceeding difficulty; the reservances between certain forms of the yellow fever and the rentents of tropical climates being numerous. But there are distintions which, on careful observation, mark the yellow fever; and these, the following are held by Drs. Gillkrest and Copland to the most prominent:—

1. Yellow fever makes its invasion more insidiously, but ru its course more rapidly than the remittent, the latter being mo

open and phlogistic.

The shock to the vitality of the body is manifested earlier
 British and Foreign Med. Chir. Review, for January, 1848, page 70.

yellow fever, as is also the contamination of the circulating fluids and soft solids,

3. Jactitation, mental depression, apathy or delirium, appear much earlier in yellow fever.

4. The lemon-yellow of the skin appears much earlier in the

yellow than in the remittent fever.

- 5. Nausea and vomiting appear earlier in yellow fever—the latter symptom being marked by more distress, and by the ejection of a greater quantity of fluid, unmixed with bile, and being composed of some exudation from the stomach, or of the serum of the blood.
- 6. The vomitings in yellow fever are unaccompanied by retchings or apparent effort.

7. The character of the pulse in yellow fever is distinctive, being rapid in its rise, then soft and asthenic, proceeding on to weakness, inequality, irregularity, and lastly, slowness.

8. The suffused, red state of the conjunctive, with more or less brilliancy of the eye, according to constitution and age of the subject, and having sometimes a drunken-like appearance.

9. The lurid redness of the countenance in many cases, and the

depressed and anxious expression in all. 10. The severe deep pain in the orbits.

11. The clean, red, raw, or bloody state of the tongue; the soreness of the throat, pharynx, and along the cesophagus, an acrid burning sensation at the stomach; the sense of constriction in the chest, and anguish at the præcordia.

12. The thick compacted feeling furnished by the skin, its diminished sensibility at the advanced stage, and the dingy tint of

it, with livid or leaden patches at last.

13. The constipation, scanty secretion, complete suppression of

urine, and absence of bile in the dejections.

- 14. The leucophlegmatic, flaccid, swollen, and pallid appearance of the soft solids, without any very evident emaciation in most instances.
- 15. The lemon-yellow of the skin is distinctive, and would seem to depend on a stagnation of dissolved blood in the capillaries, and not upon the colouring matter of bile, as in yellow remittent fever.
- 16. The continued type of the yellow fever in general—its origin, progress, and propagation, as compared to remittent fever.

PROGNOSIS.—The black vomit, although generally a fatal symptom, is far from being universally so—numerous recoveries, after its occurrence, having been observed in every recorded epidemic. But prognosis, in yellow fever, is declared by Robert

Jackson to be difficult and treacherous in the extreme, whatever the locality, or the prevalent character of the disease may have been. Referring to this disease, as last observed by him on the South Coast of Spain, he records the following symptoms as indicating danger:—

1. A sudden invasion by the fever, with intense pain of the head and eye-balls, accompanied by sickness and vomiting.

2. The fever being ushered in by a fit, convulsions, apoplectic stupor, or outrageous delirium.

3. A torpid, heavy, or statue-like aspect of countenance, gave

strong suspicion of danger.

4. A dry, rough, milky-white, or a swollen and red tongue,

indicated danger.

5. Distress and anguish at stomach, with pain at the epigastrium, forcible eructations, or explosions of flatus from the stomach gave impression of much danger.

6. Obscure hiccough marked danger.

7. A ghastly appearance, with a faint nauseous odour from the body, indicated extreme danger.

8. Yellowness of the skin, with turgid veins on the conjunctivæ

in the latter stage, "always decisive of a fatal issue."

9. Torpor of the skin—to such an extent as to be insensible to the stimulation of blisters and sinapisms—is ranked among dangerous signs.

10. Extreme dampness, or extreme dryness of the skin, indi-

cate great danger.

11. Petechiæ are suspicious: streaks, or patches, if livid, green, or violet-colour, are almost certain indications of approaching death.

12. Vomiting of black matter, like the grounds of coffee, is

reported a sign of the highest danger.

13. Vomiting of bitter bile, whether green or yellow, even with straining and severe retching, affords a sign of comparative safety.

14. Black watery stools, with shreds, "are of the worst prognostic."

MORTALITY.—In common with cholera and other epidemics, the mortality from yellow fever has been observed to have generally been greatest at its onset, although exceptions to this rule have been known; insomuch that, even at an advanced period of the epidemic, the disease has made, as it were, a fresh start, acquiring fresh power and virulence.

The violence of the disease, and its power of propagation, whether at sea or on land, have appeared to receive increase from

a certain warmth, humidity, and stagnation of atmosphere—from crowded localities and crowded apartments—from absence of cleanliness and due ventilation—from intemperance, and from dread of the disease. The following statistical details, which might be greatly enlarged, will be sufficient to exhibit the average

mortality of Europeans from this disease :-

1. "In a period of little more than four years," says William Fergusson, "nearly 700 commissioned British Officers, and 30,000 soldiers were swept away by the virulence of this disease." This was at St. Domingo, in 1795, where Fergusson served during the whole time. Referring to the fate of the French army in the same island, he says:—"Witness St. Domingo, where one of the finest armies of France perished, at least for all the purposes of an army, within the year." Sir William Pym, speaking of the epidemic yellow fever, in the several islands in the West Indies in which he served, during 1794, '95, says that there died, "in the course of a few months, not less than 6,000 men."

2. Sir Gilbert Blane states that, "in 1804, out of a population, civil and military, in Gibraltar, consisting of 16,000 persons, there died "a proportion considerably above the usual devastation of the

pestilence of the Levant," or nearly 6,000 persons.

3. "Under the mild, or what has been called the French and Spanish treatment, the mortality of Malaga, in 1804, was, 11,486, out of a population of 36,054. In the epidemic of the previous year, in the same place, 6,684 deaths occurred out of 16,517 persons attacked."

4. Dr. Rush admitted that "in the Civil Hospital of Philadelphia, where bleeding was sparingly used, the physician depending chiefly upon salivation, more than one-half died, of all the

patients who were admitted."

5. The yellow fever, which broke out at Brimston Hill, St. Christophers, in 1812, is described as having been of great severity. "The entire number of cases treated were 422, of which not fewer than 118 died, affording a mortality that treads closely upon the heels of that in the plague."

6. At Barcelona, in 1821, according to Gillkrest, "scarcely a patient survived in the wards given up to the distinguished members of the French Commission; and the mortality, under French and Spanish physicians, in the establishment called Seminaria,

was 1,265, out of 1,739 cases treated."

7. Dr. Townsend, of New York, states that, in 1822, the mortality in that city was in the proportion of three to four of those treated—the virulence of the disease being continued into the month of October.

During five visitations of yellow fever at Gibraltar, the mortality was as follows:—

Years of the epidemics . Military and their families		1810	1813	1814	1828
Civil inhabitants		17	508	132	1170
Total	5733	23	899	246	1677

9. On the island of Boa Vista, the ratio of mortality amongst the various races was as follows:—

Spaniards, Portugues	e, and	11	rei	ich			1	in	2.28
English and America	ns .				2		1	in	1.6
Native population, Sl	aves				4		1	in :	33.4
	ree .						1	in !	14.6

10. It is necessary to add, however, that epidemics are recorded in the West Indies and America as having been of a comparatively mild character; but it has been doubted whether some, at least, of these visitations were not in reality aggravated, or unusual forms of autumnal remittents. But, however this may have been, there were other instances in which the symptoms, and the rate of mortality left no doubt as to the disease having been of the true yellow fever nature.

POST-MORTEM APPEARANCES.—"If ever there was a disease," says William Fergusson, "which the humoral pathologist might claim as his own, it is yellow fever. The crasis of the blood is as much broken down before death, and its vitality destroyed, as it would be by introduction of poison of the serpent's tooth; we may truly say it is killed by the poison, and in the language of John Hunter, that 'fatal yellow fever is the death of the blood.' It swells up in floods from the mucous surface of the stomach, in the form of black vomit; and it escapes from the gums, the nostrils, the eyes, the ears, even from the skin itself, in any and every part, and after death it will be seen to have lost all the character and composition of blood, being found in its vessels like the lees of port wine, and the grounds of coffee. But, as I know well, from dire experience, the affection from the first must be essentially gastric."

Sir Gilbert Blane has "started a doubt whether the yellowness of the skin in this disease was owing to bile, but rather to some error loci, or depraved state of the red globules. 1st. This colour does not appear first in the eyes, as in jaundice. 2nd. Sir Isaac Newton observes, that the blood reduced to their laminæ assumed a yellow colour. 3rd. The colour appears in ecchymosis, some time after a contusion; and the yellowness of the yellow fever

sometimes does not come on till after death."

In common with the bilious remittents of all climates, the force

of the disease under consideration is directed almost exclusively on the abdominal organs, while occasionally, as observed by Robert Jackson, the brain is primarily affected, and the abdominal viscera secondarily. In those who die within forty-eight hours, it is no uncommon circumstance to find the liver softened and friable, while during life its function had been suspended, in common with that of the kidneys—a serous discharge, indicative of anæmia, having taken the place of natural bile.

During the epidemic at Gibraltar, in 1828, portions of liver were washed and bruised by Dr. Gillkrest, but without yielding any colouring matter. The organ he described as being pale, like box-wood. "No trace of bile has been observed in the pores; and little or no blood exuded from the viscus, when deep inci-

sions were made."

The following summary of appearances, after death from yellow fever, is taken from Dr. Copland's great work, from Dr. Gillkrest, and others:—

In the most malignant and rapidly fatal epidemics, the body seldom suffers much diminution of bulk, while the abdomen appears tumid and enlarged—there is a yellow tint of the surface, with livid blotches—some parts becoming of a brown or dark hue. The muscles are of a dirty dusky hue, and flabby in texture, or friable on pressure. The abdominal viscera are more or less congested, indicating a remarkably rapid loss of vital cohesion of the several textures. Where diminution of the bulk of the body is perceptible, on the other hand, owing to previous large loss of blood from the digestive mucous surface, the viscera will be pale, softened, and somewhat shrunk.

The post-mortem appearances in the liver, as in the other viscera, vary in proportion to the violence and duration of the fever,—exhibiting at times congestion, softening, friability—at others, a pale olive hue; while the gall-bladder is generally shrunk, containing no bile, or else there is found in it a little tarlike fluid. The rhubarb colour of the liver has been remarked as characteristic of some of the most severe forms of epidemic yellow fever. The spleen and pancreas have generally been found in a softened condition.

The œsophagus, stomach, and duodenum, present patches of a dark livid hue, with the epithelium of the mucous digestive surface detached in many places, while the mucous membrane is separated from the adjacent tissues. The stomach and small intestines contain more or less of the matter of black vomit:—this matter, as also a black jelly-like substance, is frequently found in the intestines, even where no black vomit existed during life. Less frequently there is found in the intestines a fluid of a reddish tint, more nearly resembling blood.

loping the effects of the cause; and these extrinsic influences have proved powerful in all countries.

ENDEMIC CAUSES.—The question as to the origin and mode of propagation of yellow fever is matter of evidence, and that evidence ought, by this time, to be sufficient to reconcile conflicting testimonies with each other, and with the truth. But, without troubling the reader with a long retrospect, I think it may be fairly said in respect of this much vexed question, that to maintain the disease now under consideration to be at all times and places propagated by contagion only, would be quite as erroneous as to assert that it is never thus propagated, or vice versá.

Yellow fever, then, appears to have two sources of extension— 1st, a peculiar endemic malaria; 2nd, an atmosphere infected by aggregation of the sick. In this order I purpose here to continue

the subject.

The operation of the causes will, according to Mason Good, "depend partly upon the state of the body at the time of attack, but chiefly upon some modification in the power or qualities of the febrile miasm itself, by the varying proportions of the co-operative agents of moisture, heat, stagnant air, and other auxiliaries which have not yet been detected, in their relation to each other

in different places and seasons.

How far the yellow fever is capable of origination from any other cause than febrile miasm from marshy lands, or places subject to like decompositions and plays of chemical affinity, we cannot at present determine. Such places, however, are numerous, as damp unventilated stations, stagnant water, thick impenetrable jungles, and woods that arrest the miasm as it ascends; even high arid hills, after heat and rain; but above all, a foul state of the hold on board ships, whatever be the cause of such impurity."

. . . "Heat alone, however high the temperature, is not a cause of the fever before us: there must be moisture, and, as the result of both, a rapid decomposition and exhalation of organic remains. Provided the air is dry, even tropical climates are found salubrious. The burning province of Cumana, says Humboldt, the coast of Cora, and the plains of the Caraccas prove, that excessive heat alone is not unfavourable to human life.

It has been observed, however, that even high and arid situations, after heat and rain, may also furnish, by chemical decomposition of their soil, the specific miasm of yellow fever; and it may here be added that, if by the violence or redundancy of the rain, the swampy low grounds be at the same time overflowed, the latter will become an arena of health, while the heights are the

seat of disease."

This learned and accomplished physician, on a review of all

the evidences before him, asserts that "the common remote can of this fever is unquestionably marsh miasm." He further st that—"Whatever be the original source of the fever before when once it has established itself and rages with severity, now very generally admitted that the effluvium from the bod the affected 'is loaded with miasm of the same kind, completed deteriorated as it passes off,' and that the disorder is from

time capable of communicating itself by contagion."

"The true scientific point of view," says Dr. Alison, "in w we ought to regard the invisible cause of all such disease simply as a part of the great science of Poisons-of those ago inorganic, vegetable, and animal, which, by the mysterious dis sations of Providence, are destined to exercise an injuriou destructive agency on living structures. Knowing, as we now that many such poisons are continually formed, both by veg bles and animals, the effects of which, taken into the bodie living animals, are very various, some of them truly morbific, exciting a long series of changes, constituting a true disease; that the function of excretion in animals, wherever carried of in fact the provision of Nature for the expulsion of matter wi having lost its vitality, has come under the influence, in all of the body, of 'destructive assimilation,' and, if not expelled, speedily and surely act as a poison; -knowing, also, that it the course of destructive assimilation and excretion that almost animal, perhaps all organic, poisons are formed,—we cannot be prised to find, and it is perfectly in accordance with what we now stated to maintain, that although the chemical process decomposition affords no explanation of febrile disease, ve the course of the decomposition continually going on in li bodies, in dead bodies, and in the excretions from living, un conditions not yet fully ascertained, compounds may be for from time to time which will not only act as poisons, but peculiar morbific powers, and some of which may subseque spread by contagion; and that it is only by inductive study o history of each of the peculiar diseases thus engendered, that can learn, either what are the peculiar conditions essential development or propagation, or what other powers exist in Na by which it may be opposed. Accordingly, we can refer to var facts as to several of such diseases, known only by observa but which constitute of themselves, if we make no attempt resolve them into others, an important and most useful boo science.

The principle we have already stated, as applicable, certa to several, perhaps to all such diseases, has a practical import, which justifies our dwelling on it a little farther,—that the poexciting them may have, even at this day, more modes of or than one; and therefore, that because we see such a disease spreading by contagion, i. e., by intercourse of the sick with the healthy, at one time and place, we are not entitled to infer that it must do so in all, or can have no other mode of propagation, or, vice verså, to draw the opposite inference. Without recurring to the subject of cholera in illustration of this, we have already stated that we have been satisfied by repeated personal observation, that dysentery, commencing only in the usual way, from exposure to cold and wet, and independently of any intercourse with patients already affected with it, may in some seasons assume the character of a specific inflammation, and spread from the sick to the healthy, e. g., from patients to nurses and attendants, long previously healthy, and exposed to no other cause, in an hospital. In such cases, there is nothing unreasonable in supposing that a specific poison is developed under our observation."

These observations—the suggestions of an extended experience, trained in scientific investigations—go far to explain, and to reconcile, many angry disputes and controversies of former times.

After stating his belief that the poison of cholera, although extending itself in ways not yet understood, independently of immediate intercourse of the sick with the healthy, in most parts of the world, does likewise in this country, and in others, spread at times distinctly by contagion, Dr. Alison says :- "In like manner, it is thus only that we can learn whether the malaria arising from the earth's surface, and producing only intermittent or remittent fevers in general, may or may not, in certain circumstances, and at certain times, excite fevers which may take the continued form, and may propagate their kind, as small-pox or measles do. It has been repeatedly stated in this journal, that this seems by far the most probable supposition as to the extension of the Yellow Fever at Boa Vista, in 1845, 46. This doctrine is the very same as has been repeatedly proposed by those who have studied the history of the various epidemics of yellow fever, as the only means of reconciling, not the conflicting opinions (which is a matter of little consequence), but the apparently contradictory facts presented by those calamities."\*

After quoting Sir Gilbert Blane's account of the progress of yellow fever in the Hussar, Dr. Alison concludes:—"In the face of such facts, it would be wrong to assert, on the breaking out of yellow fever, or unusually virulent remittent fever, in any climate known to be fitted for its extension, that it will not spread by contagion; but we are fully justified, by observation of that disease, in asserting on any such occasion, that whether it spreads by contagion within certain limits, or not, it will in all probability be

confined to a tainted district, even in that country."

<sup>\*</sup> British and Foreign Med. Chir. Review, loc. cit. page 331.

"What the yellow fever poison may be," says Dr. Blair, "h never been demonstrated. Whatever it be, it is probable that all localities within the yellow fever zone, it is the same thin The auxiliaries to it, however, may be different, as well as t vehicle of communication; and perhaps it may be generated in variety of soils."

"The cause of yellow fever," says Robert Jackson, "is on the act assumes a variety of aspects according to contingent c cumstances." The great majority of observers in our day seem regard yellow fever as of local or endemic origin, but as being capable also, under certain conditions, of being propagated by

contagion.

Dr. McKinlay, of the navy, speaking of the yellow fever of the Brazil Coast, says:—"My own opinion is, that the disease was not imported into Brazil; on the contrary, that it was of endemorigin. Whether it spread afterwards by contagion, or through the atmosphere, I do not profess to determine, but I feel incline to adopt the latter view. Of this, however, I feel convinced, the people arriving, more especially from a cold country, in a harbolike Rio, Bahia, or Pernambuco, when the disease is prevaled there, will almost to a certainty get the disease, even if they never touch a human being or susceptible agent in those places. I may add that this is infection, or indirect or modified contagion."

"We think we are justified in concluding," says the review already quoted, "that the presumed" hæmogastric pestilence "h no pathognomonic signs, and presents no peculiarities which me not be explained by supposing it to be the result of the most concentrated action of a particular cause. In other words, the seven fevers resulting from all the admitted causes are not distinguish

able by symptoms from each other."

In the Report of the Sanitary Commission of New Orleans, the Origin and Spread of the Epidemic Yellow Fever of 185 we find the following resolutions:—

1. "That there is no just or tenable ground for the opinion th

our fever was imported.

2. That to none of the sources to which general impression ascribes its origin, have we been able to trace even colourable pr

sumption of truth.

3. That from analogous phenomena happening at the san moment of time, and over an unlimited space in the city, and frow hat we certainly know of contagion and its mode of action, the inference is made inevitable that the fever was one of indigeno origin, resulting from local influences developed and intensified by peculiar atmospheric states."

The Reporters, finally, arrived at the following conclusions :-

1. "That the late epidemic has not been derived from abroad,

but is of spontaneous origin.

2. That there existed here, as attested by our records, very peculiar meteorological conditions, known, by general experience, to be capable of producing, in co-operation with local causes, fatal and malignant forms of fever.

3. That these conditions were present in an exaggerated degree, and impressed upon the prevalent type of disease, susceptibilities and habits of assimilating it to another and distinct form

of fever.

4. That this showed in all the localities within the range of the meteorological state or influence, an infectiousness not necessary to, or characteristic of the fever, but purely casual and accidental, the result of physical causes, and which it loses as soon as these causes are changed or disappear."

INFECTION.—To reconcile the extreme and opposite views entertained by so many able and honest writers, on the subject of the propagation, or otherwise, of this disease by effluvia from the sick crowded together, it is necessary, as already stated, to enter on a careful comparison of all the evidences as to the facts in each epidemic, which rest on individual observation and expe-

rience-discarding all opinions.

It results, I think, from such a systematic inquiry, that those authors who, on their experience during a long life of active service, as Robert Jackson, William Fergusson, Gillkrest, and others, declared that they had not been able to discover any instance of propagation of yellow fever by infection, under the order, cleanliness, and freedom of ventilation of military hospitals, were as honest and truthful, and therefore as deserving of our consideration, as those others who, like Lind, Blane, Chisholm, Pym, and others, found that under other and opposite conditions of filth, crowding, and defective ventilation on board ships, and in other unfavourable situations, this fever assumed a different-a contagious character. Here, as it appears to me, each party spoke to the absolute truth, on their observation and experience; and they reported the facts which presented themselves in each epidemic, as they occurred. It is at least but fair to say that we owe something to bygone discussions, however much we may lament their unnecessary acrimony; for from them we learn that vellow fever is sometimes propagated by infection, but more generally

Robert Jackson admits that, in respect of vessels "having a great number of persons ill of the yellow fever, a contagious fever will thus be produced, and carried to whatever port the ship may be destined to go;"—but he adds that, although the im-

Dr. Bryson, while introducing the subject of the fever on board the Bann, states that in 1823, "fever of an epidemic character began to manifest itself in the colony of Sierra Leone, and speedily extended to vessels in the harbour, in the river, or in the adjacent rivers or creeks, and to the different settlements along the coast. In the latter part of March it first assailed the crew of the Bann." Dr. Bryson goes on to say, that Sir W. Burnett was unable to account with certainty for the origin of the fever, either in the colony or in the Bann, "but distinctly proves that it was not, as had been supposed, imported by a vessel named Caroline."

Sir William expressly states, "that it was, in the first instance, merely the common endemic of the country, brought on by hard labour and exposure to the sun, not possessing, under these circumstances, any contagious properties, and continued so until after the middle of February; that it subsequently, by the state of the weather preventing ventilation, and from a great number of sick being confined in a small space, became contagious; and although it became impossible to trace the fever in question directly from the Bann to any individual of the garrison of Ascension, yet there is just reason to believe that the disease was intro-

duced into the island by the ship."

Dr. Bryson, after enumerating the symptoms generally, including the black vomit, in the ships serving on the African coast during 1823—25, says that "in almost every instance where the disease assumed a formidable character, its origin could be traced to one or more of the common well-known predisposing and exciting causes. He concludes that in none of the ships "does it appear to have assumed a contagious form but in the Bann."

In the second instance, her Majesty's steam-vessel Eclair quitted Plymouth on the 2nd of November, 1844, and after touching at Teneriffe, Ascension, and various other places on the African coast, she reached Sierra Leone on the 22nd of February, 1845. Here the crew was employed "variously"—fresh meat, vegetables, and fruits being served to them in abundance. "The general standard of health in the colony, as well as in the Eclair, was considered high; nothing in the form of endemic disease having yet made its appearance amongst the crew of the latter, nor any of that mental depression which subsequently affected them."

Between the 15th of February and the 16th of March the crew were employed along the coast, principally in boats, exploring creeks, and they were thus "exposed much to all the vicissitudes of the weather, to malarial exhalations from the muddy banks of the rivers, and from contagious mangrove swamps: they slept sometimes on shore, but generally in beds; they experienced much fatigue, disappointment, and mental depression. While this duty was being performed, diarrhœa affected many of the

at Goree, stood across to Boa Vista, where she anchored on the 21st of the same month. It being the opinion of Dr. Kenny, a resident practitioner at Boa Vista, as well as of the medical officers of the Eclair, that the two or three cases of fever now on board the ship were "the common African fever," pratique was at once given by order of the authorities on the island.

On the 30th August, sixteen new cases being added to the list, and five deaths, since the ship anchored at Boa Vista, the commander had the crew, sick as well as healthy, removed on shore—the disease being still regarded by Dr. Kenny, the Portuguese colonial surgeon, and the medical officers of the ship, as "common

African fever."

"A report, however, had gone abroad," says the surgeon, "that black vomit had occurred in some of the fever cases, and I mentioned the fact to Captain Estcourt, who assured me that he had

heard nothing to cause foundation for such a report."

Dr. Bryson goes on to state that "from the day of landing the sick, until they were re-embarked on the 13th September, 1837, additional cases of an aggravated type occurred, twenty-five of which proved fatal." It was now that, in the fatal seizure of Dr. M'Clure, who nobly volunteered to serve on board, it began to be suspected "that the disease had acquired contagious properties;" yet, even then, Dr. Bryson suggests that as Dr. M'Clure had but recently left Sierra Leone, he might have imbibed the poison there.

From the departure from Boa Vista to the 12th October at Stangate Creek quarantine station, to which the ignorant fears of the quarantine officers consigned this ill-fated ship, the record is one immense course of suffering and death, including that of the heroic young surgeon, Sidney Bernard.

Dr. Bryson, who considers the fever on board the Eclair to have been "merely of a typhoid character," concludes as follows:

"Having thus traced the fever of the Eclair from its first appearance on board at the island of Sherbro, until it ceased previously to her arrival at Sierra Leone, and from its re-appearance there in a more concentrated form, until it finally disappeared some time after her arrival in England, it is impossible not to be struck with the close similarity it bore to that of the Bann. Both vessels contracted the disease at Sierra Leone, and apparently from the same cause or causes, and under similar circumstances. In both vessels, in the course of a few weeks, it assumed an epidemic character, if it did not acquire contagious properties; the one vessel proceeded to the barren rocky island of Ascension, a few degrees south of the equator, where a disease of the same character made its appearance amongst the inhabitants, and committed great ravages; the other proceeded to the nearly equally

barren island of Boa Vista, a few degrees to the north of the equator, where, in like manner, a disease a short time afterwards broke out and raged with equal severity. This is not the place, however, to trace the progress of the fever on the island of Boa Vista, nor is it, perhaps the time, considering that the sub-

ject is still under investigation."

The investigation here referred to was soon afterwards concluded by Dr. M'William of the navy, in a manner alike honourable to his reputation, and satisfactory to the great body of the medical profession. Two points have been established by Dr M'William; namely, that the fever on board the Eclair had become contagious; and, secondly, that the disease was propagated from the ship to the inhabitants of Boa Vista. We thus find that the similarities between the case of the Bann and that of the Eclair, are highly interesting and important.

1. "In both instances the fever became increased in violence

from the influence of the shore.

2. At Ascension, the mortality in the garrison was over 50 per cent. of those attacked, while it was but 16 per cent. in the crew of the Bann, although they were under the most unfavourable circumstances of ventilation, cleanliness, &c.

3. At Boa Vista, the sick list became rapidly augmented when

the sick were landed.

4. In both cases also the coloured people on board escaped, or

nearly so.

5. If these cases of the Eclair and the Bann be taken together, agreeing as they do in all their main features, it is logically impossible to doubt the existence of a multiplying or contagious virus."

Thus far the able reviewer already quoted, who now arrives at the following conclusions from the whole course and tenor of the argument:—

"1. That yellow fever is decidedly contagious on some occa-

sions.

- 2. That the proofs of the universality of this property are defective. Importation can seldom be proved; inoculation is impossible, and the disease requires a peculiar susceptibility of condition in order to act.
- 3. That it is not safe to generalize from the first observation, and to conclude that yellow fever is at all times contagious, because it is undoubtedly so sometimes; for these reasons:—

a. Because there are fevers undistinguishable by symptoms from contagious yellow fever, which certainly are not contagious.

b. Because we are not sure how far contagion may not be an accidental property impressed on a poison by contingent circumstances, or may be only the development of a property of self-reproduction always possessed by the morbid poison, but generally

in so slight a degree as to be unappreciable. If the former opinion be correct, yellow fever is both contagious and non-contagious; if the latter, it is also in the majority of cases, non-contagious, in the conventional meaning of the term.

4. That contagion is only a property accidentally impressed on

the yellow fever appears probable.

a. Because in no other way can we explain the extraordinary discrepancies and opposing statements of men, whose honesty of purpose is undoubted.

b. Because it is almost a necessary assumption, in order to ex-

plain certain facts in the history of yellow fever.

c. Because it is in accordance with analogous phenomena ma-

nifested by other morbid poisons.

d. Because there is really some direct proof of it in the apparent development of fever on board ships with clean holds, and removed from the influence of the land.

5. That if this conversion of a non-contagious into a contagious poison be denied, there is no alternative but to admit the exist-

ence of a specific contagious yellow fever.

6. That the doctrine of a specific contagious yellow fever is alone supported by the fact, that it destroys the necessity of admitting the convertibility of poisons, a circumstance considered by many observers as in the highest degree unlikely and unphilosophical."

The following is Dr. M'William's summary of the questions relating to the fever on board the Eclair, and that which prevailed

subsequently at Boa Vista:-

"In my humble opinion, the history of the epidemy of Boa Vista comprehends every condition upon which the proofs of the infectiousness of a disease are supposed to rest, namely:—

The healthiness of the island before the arrival of the Eclair,

with Yellow Fever on board.

The outbreak of the same disease among the inhabitants of

the island within a reasonable period afterwards.

The immunity of distant villages for long periods until the arrival of infected persons, and the radiation of the disease in every district from infected foci.

The comparative immunity from the disease obtained by persons who adopted common but partial precautionary measures

against infection.

The absolute immunity from the disease procured by persons who adopted strict measures of isolation and segregation."

On a careful review of this question, we receive the following

additional authority in support :-

On the 31st December, 1850, the Royal College of Physicians of London delivered a Report to the Lords of Her Majesty's most

Honourable Privy Council, respecting the Bulam, or yell which had been drawn up by the President and Fellow College in consequence of a communication from their La

The points investigated by the College were as follow:—
"1. As to the Bulam fever being sui generis, and distiremittent or the marsh fever of warm climates.

As to its being an infectious disease; that is, comble from person to person, and likewise capable of being in
 As to the non-liability of persons to a second attact

disease.

After a very careful consideration of all the facts as ments adduced on both sides, with reference to the first proposed, the College are of opinion, that sufficient have not been laid for stating, that "yellow fever" is a sui generis.

With regard to the second question, it appears to the to be sufficiently proved, that this disease is, under certain stances, infectious, and, consequently, that it may be impo-

The principal circumstances under which the infect of the disease is likely to be developed, would seem to be temperature and moisture of the atmosphere, particularl healthy seasons, and when the influence of these cause gravated by local insalubrity of site, and by the absence ventilation

That the disease has been in some instances import history of the epidemic fever which occurred in her Majes Eclair, and at Boa Vista, in 1845, affords conclusive evidence

The third question proposed respecting the non-lial persons to second attacks, does not admit of being sett decided manner. Strictly speaking, there is no disease of the can be affirmed, absolutely, that one attack renders a posusceptible of a second.

The College are unwilling to conclude their report strongly recommending to their Lordships, that, on an appearance of this formidable malady, persons should out, thoroughly qualified by education and by habits of tion, to collect evidence on the important questions whi now been proposed to the College. This was done in the the Eclair, and most valuable information was thus obtain

(Signed) "Francis Hawkins, M.D., Reg "Royal College of Physicians,

December 31st, 1850."

PREVENTION.—Dr. Alison justly observes that, we more for the prevention of sickness and suffering among n by studying the mode of propagation of epidemic diseas

so far as yet known, and giving that advice by which they may be shunned, than we should do by the discovery of a new remedy, more powerful than any that is known in medicine. The truth of this declaration ought to be deeply impressed on all who enter the medical departments of the public service; for it is with yellow fever, as with cholera, and other concentrated forms of disease;—whatever will prove most effective must rest on measures of prevention; and of these, the immediate removal from the place where the yellow fever may prevail, and the not returning to the locality, or to the ships, originally occupied by troops, inhabitants of towns, or seamen, constitute the first necessity of the case. The most eligible localities to choose will be found in elevated and well drained situations, to windward of the infected spots.

Dr. Alison further states, that the most important principle ascertained in this department of medicine is—"that as to several of the most virulent of the epidemics which have been permitted to afflict mankind, the means of effectually restraining their diffusion are known, but cannot be carried into effect without both trouble and expense to the community that is to benefit by them; and that these means are very considerably different in the case of these different diseases—in the case, particularly of small pox, of erysipelas, of puerperal fever, of plague, of cholera, of continued

fever, and of yellow fever.

As to yellow fever, regarding it as originating in an aggravated form of the remittent from malaria, and therefore affording illustrations of the natural history of that destructive agent in its most virulent form—we can go somewhat farther. Recollecting what has been stated of the very limited origin of the Sausage poison, and of the cadaveric poison exciting erythematic inflammation—not in putrefaction generally, but in a certain stage of the decomposition of certain matters only—we shall be prepared to hope that a somewhat similar limitation may be ascertained for the development of this poison likewise; more particularly as we have already stated, that both in this climate, where it (in general at least) only excites ague, and in hot climates where, under a temperature of 80° or more, of some duration, it excites so much more formidable diseases, all the conditions which we can as yet point out as essential to its development may exist without its showing itself.

On this point the experience of the medical officers of the army and navy, for the reasons already given, being the result of observations on organized masses of men, is the most valuable; and facts are already known which may often be applied to practical use, and promise to lead hereafter to more minute and more uniformly applicable information. Not only is the general measure of removing the population of a district known to be malarious always to be recommended when the yellow fever shows itself, but

precise limits may sometimes be assigned. It seems certain that the poison often originates in parts of the timber of certain ships,

and is long confined in them." \*

That on board ship, as on shore, the limits of putrefaction or other processes of decomposition are confined within narrow limits of space, seems more than probable; and the special investigation of such limits constitutes a branch of inquiry of the utmost importance, and one in which many facilities are afforded to the medical officers of our fleets and armies.

The most speedy means of prevention, in respect to towns and garrisons, will always be found in the removal of both the sick and the healthy to a locality where the temperature is sufficiently low, such as a neighbouring elevated range, or dry well ventilated

ground: the next most ready means is segregation.

1. In Gibraltar, in 1810, the infectious character of the prevailing yellow fever being suspected, "the infected were segregated and removed to an airy locality, communication with them was prevented, the healthy protected, and the mischief was very soon arrested."

2. In 1813, this disease again made its appearance there, when nearly 8,000 persons left the garrison and encamped on the neutral ground. The open nature of the ground, and the strong breezes, arrested the progress of the epidemic, very few cases having occurred after this removal.

3. Upon a review of all the facts and circumstances observed at different times, at Gibraltar, Dr. Hennen concludes, that they were demonstrations of the value of "segregation and moderate

quarantine."

4. "The disease will stop," says Dr. Copland, "as soon as the susceptible are separated from contaminated places, persons, and

things."

- 5. Let houses, barracks, and hospitals be purified by water and air; but no attempt be made, during the prevalence of the epidemic, to cleanse sewers, drains, or privies, for such measures have always proved mischievous, when practised at the time stated.
- 6. Tents, boarded sheds, and temporary hospitals should be so separated from each other as to admit of the most free circulation of air, in every direction.

The prophylactic influence of bark or quinine has often been observed during service in unhealthy countries, but nowhere more

emphatically than on the West Coast of Africa.

"In the North Star," says Dr. Bryson, "twenty men and one officer were employed on boat duties at Sierra Leone; they all took wine and bark with the exception of the officer; he was the

<sup>\*</sup> British and Foreign Med. and Chir. Review, April, 1854.

only person who suffered an attack of fever. Two boats were detached from the Hydra, in 1844, to examine the Sherbro river; the whole of the men were supplied with bark and wine, and not one of them was taken ill, while the whole of the gig's crew, with the exception of the captain, who were similarly exposed for two days only, without being supplied with either, contracted fever of a dangerous character. Facts like these are not to be mistaken."

Dr. Bryson suggests that a daily prophylactic dose be administered, and that its use should be continued for at least a fortnight after the return of the men on board their ship. He gives a

decided preference to quinine over the bark.

Whenever fever of a formidable nature makes its appearance on board ship, she should at once proceed to sea, and into the coolest

atmosphere within reach.

With seamen, as with soldiers, the first and most immediate measures of prevention should obviate direct solar exposure. fatigue, and excesses in the use of spirituous and fermented liquors.

2. Seamen, when employed on unhealthy coasts, should be kept as remote from the shore as is consistent with duty-anchor-

ing some miles out, during the night especially.

3. The duties in boats should be conducted as much as possible in the mornings and evenings—avoiding alike the noon-day heats, and the deadly emanations from the shores common to the night.

4. The labour in boats should be conducted with as much care to the men, and absence from fatigue as possible, avoiding ex-

hausting perspiration.

- 5. The seamen should not land, unless under urgent necessity; and then the more high and dry lands should be carefully selected for encampment.
- 6. The meals should be regularly served, and carefully cooked -no more spirit-ration being issued than is customary.
- 7. Under necessary fatigue, cold, wet, and mental depression, the issue of extra spirit-rations will be proper.
  - 8. Coffee should be given early in the morning, as a habit.
- 9. One or two doses of quinine should be given daily as a prophylactic.
- 10. The rendezvous boat should have an ample tarpaulin for the protection of the men against night dews and noon-day heats.

11. Besides a complete set of blanket-clothing, each man should

have a duck-dress and flannel waistcoat.

12. Where the nature of the service can admit of it, no boats' crews should be absent from the rendezvous boat, or from the ship during the night. This would constitute, of itself, the greatest safeguard of all.

13. Dry-cleaning of the decks—attention to cleaning out the holds—and great attention to the personal cleanliness of the crew should be carefully observed.

14. The free ventilation of every part of the ship should be

carefully maintained.

15. The labour of the men should be sparingly used, and they should, when possible, not be exposed to wet or damp; and they should not be put to work till the sun shall have dissipated the early morning fogs, and till after the serving of coffee.

16. The holds of ships should not be cleansed on the spot where the fever has originated, or during its prevalence, but should be deferred till the arrival of the vessel in a colder latitude.

17. Green wood should not be placed on board ship in hot climates. Wood for ship use should be stripped of the bark, cut into moderately sized pieces, and partly charred. These directions are chiefly taken from the admirable Report of Dr. Bryson, already quoted.

CURE. — Previously to entering on a consideration of the means of cure, the physician should determine, with the utmost care, both the type of the prevailing fever, and the age, constitution, and habits of life of the individual. There are, beyond question, pervading characters in every epidemic—one being of an adynamic, another of a somewhat sthenic, or phlogistic nature; and it is obvious, that on the right determination of these preliminary questions, very much must depend. But, with all the care which we can employ, it must be confessed that, as regards the essential nature, causes, and treatment of this terrible disease, the amount of our real knowledge is lamentably defective. In the yellow fever, as it appeared on the coast of Spain, the brave Robert Jackson speaks hopelessly as to the influences of all means of cure; and he was in all things the very opposite of a despairing physician.

"From the wide range which the symptoms take," says Dr. Gillkrest,—"so wide that but for the black vomit being liable to occur as a connecting link, in the various forms during the prevalence of an epidemic, we should, from the symptoms, as well as from post mortem appearances, often have reason to suppose that different remote causes were giving rise to different impressions. It is obvious that in no disease is it more difficult to lay down rules of practice, and in none can the medical officer's tact and atten-

tion be more needed."

"All medicine," says Rostau, "consists in diagnosis;" but he adds, as justly as emphatically, that "the medicine of symptoms is the worst of all medicines." We must, therefore, beware of a false diagnosis; and recollect also that, as regards the important question of treatment, it too often happens that "What is new is

not true, and that what is true is not new." Holding in recollection these preliminary cautions, we shall be the better prepared to enter on the question of the little that is known, therapeutically, of yellow fever,

The history of the treatment of this disease, like that of the bilious remittents of hot climates generally, is found stamped with the characters of the doctrines taught in our medical schools,

rather than on any ascertained etiology or pathology.

As in the remittents of the East, at different times, so in the Western epidemic yellow fever—emetics and purgatives formerly preceded a systematic course of Peruvian bark, wine and opium; blood-letting being used only where violent reaction, and conse-

quent inflammation was supposed to exist.

But this plan of treatment not proving satisfactory to Dr. Rush, of Philadelphia, he had recourse to repeated blood-lettings, seldom exceeding sixteen ounces at a time, pursued into the fifth and even the seventh day of the disease, or so long as the pulse retained any firmness. Calomel with jalap powder was freely administered throughout, as a purgative and cholagogue; and this course of treatment Dr. Rush declared to be the most successful in his time.

In the West Indies, Robert Jackson abstracted blood even more freely still—to the amount of thirty to forty ounces at a time, and repeated until a powerful impression was made on the disease. Then followed Dr. Chisholm, with "bleeding to the extent necessary, plentiful alvine evacuations, MERCURIAL PTYALISM, and cold affusion."

Different as such means of cure may appear to have been, they were not in reality so different as they seemed; for the yellow fever epidemics, observed and described by so many able men, differed as to time, character, circumstances, and localities, so as very greatly to reconcile certain differences in their treatment. Such influences on health and on disease must ever be held in recollection in passing judgment on the conduct of those who have gone before us; otherwise, we should be doing injustice to their memories.

Referring now only to the experiences of our own day, and of epidemic visitations such as we see them, it must be confessed that it is not here as in remittent fever: we must not have recourse to heroic measures of cure; for we cannot cut short the yellow fever.

In more recent times blood-letting has not proved needful or useful, whether in this epidemic as it prevails on the African coast, or that of Spain, or yet in the West Indies. As in cholera, the shock to the organic nervous system, and the injury to the blood, especially when the subjects of treatment are placed in low, damp,

or crowded localities, induce so great a prostration of the and vascular functions, even in the young, and in those ap of a phlogistic habit, as to render any considerable abstriblood from the arm a more than questionable operation. general or local depletion promises relief, it will be found vantage amongst the newly arrived and plethoric, and placed in favourable, open, and airy localities, and which character of the epidemic fever is more of a phlogistic of than otherwise.

Having duly considered the individual circumstance patient—the stage of the disease, and its degree of seve should be placed in a position which secures a free ver and he should be made to look with a confident hope to sult. The following means should then be put in practice

The warm bath will be found grateful and useful as a nary measure, having in view the objects of purifying and promoting a determination to the surface of the body temperature should be such as to prove graeful to the and gentle frictions with oil should follow the use of the b

Of internal remedies our list is necessarily limited; for so much irritability of stomach, with such dangerous von prospect, that even the most useful and agreeable medic be rejected. With a view to anticipate and allay this di and dangerous disposition, effervescing drinks in mode repeated doses should be given; such as soda or seltz flavoured with champagne; spruce or ginger beer—all w grateful to the stomach.

As in remittents, so in this disease, the bowels should evacuated by means of jalap powder with calomel, or by followed by senna and salts mixture. Throughout the prothe fever, indeed, and whatever the degree of its severity purative functions must be kept in activity by alterative mercury, purgatives, and diuretics. In this manner, means, and with the aid of farinaceous diet and demulcen the severer forms of the disease may be somewhat mitigate the milder degrees of it will in a few days be advanced it valescence:—but, too often, we have to deal with more gr

If, in the severe cases now contemplated, blood-letting recourse to, it should be abstracted from the epigastrium ously fomented, and by means of leeching or cupping—avoiding the induction of vascular and nervous depression have now to regard black vomit and the death of the paprobable contingencies in the progress and termination case; and we must, with such means as we know, proceed ticipate and prevent them. Of these, turpentine exhibite stomach and in the form of enema, and applied to the w

domen as an epithem, has been for many years past recommended by Dr. Copland, as holding forth a favourable prospect—assisting its operation by wine, brandy, camphor, and opium, as diffusible stimulants.

In the epidemic yellow fever of Peru, described by Dr. Archibald Smith, and which took three years to attain to its full development, one attack formed no security against a second. From the very gradual manner in which this epidemic acquired maturity, so as at length to constitute unequivocal yellow fever, the Spanish physicians compared it to the tender cub which, as it

grew up, became the voracious beast of prey.

The treatment found successful in this epidemic, after the failure of other remedies, was turpentine in one drachm doses by the mouth, or half-ounces in the form of enema "several times daily"—using it also as an epithem to the abdomen. In this manner fifty per cent. of the Indian race are stated to have been saved, even "in the last stage with black vomit." This course of treatment was not so successful with the other coloured and mixed races, and it was least successful with the European. Dr. Smith "can only recal to mind three decided instances of a white man having reached the stage of full homogeneous black vomit, and yet to have survived that danger."

On the decline of the febrile stage, the infusion of valerian with ether, camphor, and musk, was generally used, as were the carbonate of ammonia, opium with creasote, camphor and opium, and capsicum with opium, in form of pill. Two minims of creosote and half a grain of opium in pill, and administered every four or six hours, were very effective in preventing the accession of

black vomit.

Very recently also this practice has been very favourably reported upon by Mr. Laird, of her Majesty's ship Medea, who exhibited the turpentine in doses of fifteen minims, with a little nitric ether and camphor mixture, every three or four hours, which course of treatment was pursued until a remission of symptoms ensued; when quinine and a nutritious diet concluded the treatment.

As adjuvants, occasional doses of blue pill followed by castoroil—sinapisms—cupping—blisters—evaporating lotions to the head, &c., were employed; and, out of sixty cases treated, only four deaths occurred. This was indeed a remarkable, and I believe an unique amount of success; and should it prove in other epidemic visitations of yellow fever equally true, we shall no longer approach this disease with hesitation, the result of dread.

Blood-letting Mr. Laird found to be "extremely hazardous," and an operation which, "except in some particular cases, might be safely dispensed with;" but local depletion by cupping, from

the epigastrium and nape of the neck, appears to have been occ

sionally useful.

The action of the turpentine on the skin and kidneys is rate highly by Mr. Laird, who found the medicine seldom to disagre with the stomach, or induce strangury. Turpentine exhibited in ternally did not excite the pulse; and applied to the abdomen: friction, along with olive-oil, "seemed to act better than if it ha been taken by the mouth," in the instance of "a sailor-lad, with black vomit and suppression of urine, who would not take th draught." The report of this interesting case goes on to say the "three hours after the frictions were applied, this patient made pint of healthy-looking urine, and, after an interval of eight hour he passed a pint and a half more urine of the same characte Early next morning black vomit, which had been present for about thirty-six hours, entirely ceased, and the lad appeare otherwise much improved." The frictions were continued; bu while struggling against the administration of a nutritive enema he was seized with convulsion, and died.

"On dissection, no trace of black vomit could be detected unless a little dark green matter found in the ileum might be considered as such. The mucous lining of the stomach was muc congested, and discoloured dark-red; but its epithelium was perfectly intact; and although the liver presented the usual faw colour, the skin was not tinged yellow, as was the case with the other patients who died with black vomit. This case, I might be excused saying, ought to have been a recovery from black vomit.

had not the unlooked-for head affection come on.

In another case of black vomit, I gave turpentine in doses of forty minims, combined with hydrocyanic acid, with the eviden effect of controlling this fatal symptom, although the patient wa carried off soon afterwards by profuse and uncontrollable hæmorr hage of fluid-looking blood from the mouth and nose."\*

Dr. Blair, of British Guiana, another recent writer on yellow fever, was sparing in the use of blood-letting, even as a local mod of abstraction, and which he had recourse to only for the relief of

severe and early cerebral and epigastric distress.

His treatment consisted in twenty-grain doses of calomel, wit twenty-four of quinine, followed by castor oil. In severe case he repeated the calomel and quinine every six hours; and never had occasion to exhibit more than six such doses to one patient and ptyalism did not result from this treatment. "Emesis was frequently moderated or checked by the exhibition of drop dose of creasote made into a light emulsion with mucilage and sugar."

This gentleman used cold ablutions, sinapisms, and blisters while he exhibited as adjuvants-Rhenish wine, chalk mixture

<sup>\*</sup> Medical Gazette, April 14, 1855.

with essential oil, creasote, liquor potassæ, ammoniacal paste, musk, carbonate of ammonia, spirit of Mindererus, magnesia, laxative enemata, lemonade. The diet consisted in barley-water, sago, arrowroot, tea, chicken-broth, beef-tea, and toasted bread.

The treatment of the epidemic yellow fever, as described and

treated by Dr. McKinlay, was as follows :-

Purgatives, calomel and antimony every three or four hours, cooling saline diaphoretics, sponging with vinegar and water, quinine and bitter infusions during convalescence, with barley and rice-water for drink. To alleviate and overcome vomiting, he used effervescing draughts, with tincture of opium, enemata containing turpentine, sinapisms and blisters to the epigastrium.

Such is a fair exposition of the most recent means of cure adopted in the treatment of yellow fever. That all or any of them may prove very generally effective to the cure of other epidemic visitations, is more than present experience will warrant us to conclude; but that they present subject-matter worthy the attention

of the tropical practitioner, may, I think, be admitted.

Desiring to attract and to concentrate the attention of the reader to the little that appears to be practically ascertained respecting the modes of treating yellow fever, I have purposely avoided the entanglement which might result from an enumeration of all the adjuvants which have at different times found a place in the lists of authors, and which are common to the treatment of yellow fever as to the remittents of tropical climates. The truth is, that many remedies have been recommended for the cure of both diseases—not on the ground of proved experience, but simply as unreasoned opinions. We have been oftener told what they were, than what they signified.

## DELIRIUM TREMENS.

Without calling in question the occasional occurrence of acut delirium, as an affection connected with inflammatory irritation of the arachnoid membrane—the conjoined result of "a drunker bout," and of insolation—I purpose here to notice only the tru delirium tremens, as it generally appears amongst European soldiers in the East Indies. It is usually seen as the indirect result of the temporary withdrawal of the accustomed stimulating drink or it arises, in a more or less subdued or masked form, when the soldier has for some days been subjected to the abstinence of the hospital system of discipline, on account of a wound or accident or for the treatment of acute disease—as fever, dysentery, o hepatitis.

The importance and interest attaching to delirium tremens, or the brain fever of drunkards, will be rendered apparent by a reference to the numerical table at the end of this article, and with which have been favoured by Colonel Tulloch. The disease, as already suggested, is important also, as forming a very frequent complication with many of the acute diseases of tropical climates—a face

not sufficiently noticed by the generality of authors.

In a report on the statistics of Delirium Tremens among the troops in Canada, during 30 years, by the Inspector General Henry, we find the following general results:—

Number	r of cases	of Delin	ium Tremens in	30	years			. 1	1769
"	deaths		"		-				143
Ratio o	f cases to	strengtl	n, first 15 years			as	1	to	175
"	"	"	second 15 year	S		as	1	to	75
>>	,,	"	whole 30 years	3		as	1	to	93
"	deaths to	cases,	first 15 years			as	1	to	16
99	"	,,	second 15 years			as	1	to	12
"	"	,,	whole 30 years			as	1	to	12

SYMPTOMS AND PROGRESS.—Delirium tremens appear in forms more or less acute and complicated, as the disease ma occur in the newly-arrived or in the old seasoned soldier. In the cases of young soldiers, we find that occasional courses of drinking, with insolation, will produce great perturbation and excitement of the nervous and vascular functions, amounting, in some instances, to maniacal fury—the most strange illusions, violence, and hurry of manner, with incessant muttering and general irritability, the face being flushed, and the head hot. There is great anxiety, with oppression of the præcordia, vomiting, spasmodic pains, and occasional headache; the urinary and intestinal excretions being scanty and deprayed, the tongue foul and tremulous. While this earlier stage continues, the skin is harsh and dry, the pulse frequent; but if the disease proceed, symptoms of exhausted nervous and vascular energies take the place of the previous tumult and excitement, even in the youthful and newly-arrived soldier.

In the old soldier, again, habituated to the abuse of intoxicating liquors, every debilitating cause—even the withdrawal of the accustomed bane, is followed by collapse of the vital powers, as indicated by nervous depression, excessive gastric and intestinal irritation, a feeble pulse, sometimes frequent, but more generally slow, a cold and clammy skin, a hurried and imperfect respiration. The manner, though wild, is more subdued, and the temper melancholy, the mind being in a perpetual anguish—haunted with imaginary alarms and illusions; but when the attention is aroused by a question, the answer is generally rational. The wretched sufferer, deprived of half his reason, is yet miserably conscious of his condition; in short, the moral and physical ills produced by habitual inebriety, are here concentrated.

After a time sleep is entirely banished, with aggravation of all the symptoms; until at length, as the disease advances, and the energies sink, a complete exhaustion of the vital powers announces the immediate approach of danger. I have seen men walking about in a frenzied state, muttering and chattering, whose death took place within a few minutes, instantaneously, and as if struck by lightning; while in other instances, death was preceded by aggravated tremor, ending in general enervation and convulsion.

Mr. Playfair, of Bengal, describes a soldier as insisting on being discharged from the hospital, because the guard had conspired against his life; or, he saw his comrades attempting to shoot him through the windows. He then collected his bedding, and everything he could find, and piled them against the door, to guard against the supposed attack, crying aloud for justice or revenge. With the fury and strength of a madman, he then threw everything about the room; after which he seated himself in the corner, and with eyes fixed on vacancy, he listened to and conversed with imaginary beings. In this latter mood he would injure no one; indeed, such is not the general desire, though there is usually some

person of whom the patient complains. As in ordina so in this disease, personal fear predominates.

The patient often calms on finding himself under to fithe strait-waistcoat, and begins to relate his griev then only can he be prevailed on to take medicine; at restraint only seems to increase the fury of the patient at the mouth with rage, blaspheming in the most horri

A patient supposed himself a messenger sent from save the souls of those who sought his protection, and the heavenly hosts on the wall, with "the devil lookin corner, afraid to advance." Another soldier imagin charged with the most unnatural crimes, but asserte cence in the most pathetic manner; while a third belie condemned to be shot for a crime he never commithough he was to suffer innocently, he called for instant and, placing his back to the wall, requested that a might be ordered to despatch him. A fourth complair wife's honour was in danger from the designs of vilifting up the corner of the quilt, pointed to Mr. Playfa of her imagined concealment.

A soldier of the Bengal Artillery is mentioned by who required six men to secure him in the strait-jacke stripped himself naked during the night, burst from into the adjoining dead-house, barricading the door on which the bodies of four deceased comrades were ly an appalling scene was presented—the raving drunka close conflict with the dead, uttering at the same tindreadful imprecations that language can afford. Suc of the effects of spirit drinking on the European soldie

CAUSES.—With soldiers habituated to excesses, privation of the accustomed drinks—the absence of nourishing food—the debility caused by acute disease dysentery, diarrhœa, and cholera—active measures of these are all frequent causes; as are, the abuse of tob ing—the use of opium, and of drugged beverages.

But while it is admitted that, in young men, occasion of drunkenness, with or without solar exposure, commost frequent causes of the more acute forms of delirius there can be no doubt that, with the older soldiers of residence in tropical climates, the habitual tippling ar smoking, especially when, of a sudden, one or all of the stimuli have been withdrawn—forms by far the most ge of the disease in its true and more frequent examples.

The truth of this is seen in the relative statistics of and navy. In the latter, though delirium tremens is f

sult from excessive debauches in harbour, the disease is neither of such frequent occurrence, nor so fatal in its results by many degrees, as in the army, and especially amongst soldiers who are paid daily; for with these last, in place of the periodical drunken bout following upon the receipt of the month's pay, after the olden custom, there is now prevalent a far more injurious course of excitement in the daily tippling of ardent spirits.

COMPLICATIONS.—The complications of delirium tremens would alone constitute this disease one of much importance in military practice; for not only is it, in this form, very prevalent, but it is both insidious and dangerous in the hands of the inexperienced.

In the soldier habituated to the use of ardent spirits, nervous symptoms arise as frequent complications in course of treating him for other diseases, and such symptoms will always be found, varying in degree, in persons of better condition who indulge freely in wine, spirituous liquors, and tobacco. After the free use of depletory means for the removal of fever, dysentery, or hepatitis, how often do we find the patient exhibit at first a latent or subdued nervousness, soon to be unmasked in all the illusory and real horrors of delirium tremens: \* in other words, when febrile or inflammatory symptoms have yielded to the use of blood-letting, purgatives, sudorifics, and low diet, how often do we see the patient become low, desponding, tremulous, agitated, and sleepless; symptoms which, if disregarded, become sources of aggravated suffering and danger in the sequel: in short, life may here be sacrificed, if the enervated condition under consideration be mistaken for, or treated as, cerebral inflammation.

It is here then, when, under the depressing influences of the measures of cure indicated, room has been afforded for the display of nervous irritability and exhaustion, that other quieting means of a special nature become really necessary to the successful treatment of the case; and many a physician has found the safety and advantage of such means, under such circumstances, without having been sufficiently alive to the reason of his success.

PATHOLOGY.—Congestion of all the secreting organs, and consequent arrest of all the depurative functions, would appear to constitute the most ordinary results of excessive spirit-drinking. When delirium tremens has actually set in, the urine is declared by Dr. Bence Jones to present an extraordinary diminution in the total amount of phosphatic salts excreted, as compared with the vast increase of the same in inflammation of the brain.

<sup>\*</sup> Delirium tremens is always designated by the European soldiers as "the horrors."

"Alcohol," says Dr. Bence Jones, "hardens the skin applied locally. It hardens and thickens the stomach. long continued it thickens the cellular tissues of the liver. It off by the lungs and kidneys, and affects both organs. Ultimetremoval of the products of decomposition by the urine. It elst the composition of the blood, and affects the nutrition of organ and structure in the body. It has its own peculiar pous action on the muscles and nerves. As a poison it has placed by its effect on nerves and muscles between ergot and arsenic."

"Alcohol is the chief poisonous ingredient of unadulterat mented liquids. Like all poisons, when not taken in too le quantity, or too long continued, it has a medicinal acti which its dietetic value depends. This action may be sumn

in one word, stimulant."

Alcohol, according to Dr. Bocker, diminishes the exc both of the solid and fluid constituents of the urine.

2. Alcohol does not increase the cutaneous perspiration.

3. Alcohol does not augment the fæcal excretion.

 Alcohol diminishes not only the absolute quantity of ea acid exhaled by the lungs, but also the relative proportion

in the products of respiration.

Dr. Ogston of Aberdeen, in his pathological observations bodies of known drunkards, gives the following results of spection of the bodies of 44 persons:—

1. Cranium.—Abnormal appearances within the cranit

43 cases, 97.7 per cent. of the whole.

 Chest—Abnormal appearances in the respiratory orga all, in 33 cases, or in 75 per cent. of the whole. Simulta abnormal appearances in the head and respiratory organs cases.

Abnormal appearances in the pericardium, heart, aorta, o monary artery, in all, in 26 cases, or in 59 per cent of the

Abnormal appearances within the chest, in all, in 38 cas in 86.3 per cent. of the whole.

3. Abdomen.—Abnormal appearances in the stomach, in

24 cases, or in 54 5 per cent. of the whole.

Intestines.—Abnormal appearances in the intestines, in

8 cases, or in 18:18 per cent. of the whole.

Liver.—Abnormal appearances in the liver, in all, in 36 or in 81.8 per cent. of the whole.

Spleen.—Abnormal appearances in the spleen, in all,

cases, or in 40.9 per cent. of the whole.

Kidneys.—Abnormal appearances in the kidneys, in all cases, or in 63.6 per cent. of the whole.

Entire absence of morbid appearances existed in none of the cases.—(British and Foreign Medico-Chirurgical Review for

October, 1854.)

Such have been the post-mortem appearances. But, during life, besides the contamination of the blood from the generally pent up and vitiated state of the secretions—the result of universal congestion—there is reason to conclude that, as a direct consequence of spirit-drinking, the circulating fluid is dissolved, and otherwise deprayed. The result altogether is—a severe irritation of the entire nervous system—followed by nervous and vascular collapse.

PREVENTION.—Doses of two grains of emetic-tartar, or half-drachm doses of powdered ipecacuanha, have been exhibited as means of prevention of habits of drunkenness; some practitioners recommending that the emetic be given at the moment that craving for strong drinks comes on, while others add the medicine to the intoxicating potation with a view to excite nausea, or cause vomit-

ing, and thus produce a feeling of disgust.

Dr. Gilbert, of Lurgan, administers tartar emetic "secretly, in occasional doses of from five to ten grains, as soon as convenient after the person commences drinking strong liquors in tea or in any other proper vehicle. It generally produces such nausea, vomiting, and fear of death, as completely to frighten the person from going on in his drunken course of life, for that time at least; and, much to the joy of friends, ultimately produces, in many instances, what nothing else would effect, a complete change of life, one from drunkenness to temperance."

Mr. Higginbottom, of Nottingham, considers the ipecacuanha more safe, and altogether preferable to tartar-emetic; and that from its "extraordinary power of stimulating the whole system, equalizing the circulation, promoting the various secretions, and indeed of assisting each organ of the body to perform its function, and to restore it to its normal state"—ipecacuanha should always have the preference. He gives the powder in half-drachm doses.

The inspector-general, Henry, on an extensive observation of delirium tremens as it appeared in Canada, says:—"As the diarrhoa, which so often warns of malignant cholera, may be readily cured by proper medicine, and the dangerous attack averted; so, in the premonitory stage of this malady, a manly effort, aided by a little medicine, may save both the reason and the life. When the tippling of the sot, or the outright intemperance of the drunkard, are about to end in this disgraceful disease, the poor slave of evil habits will sometimes be induced to make great efforts to break his chain; but at first he will feel himself most miserable: relaxation and debility have succeeded artificial tension and strength, whilst

the stomach craves its customary stimulus, and some s must be found.

According to my experience, gentian, as below, is the dicine under these circumstances:—

R Infus. Gentianæ comp. 3xij. Tincturæ ejusdem, 3v. Magnesiæ sulphatis, 3iv.

A common wine-glassful the first thing in the morning,

mid-day, and in the evening.

If the patient can be persuaded to take it regularly for his morbid craving for stimulating drink will be abated, the his stomach improved, his moral dejection lessened, and feel himself a new man. Of course this applies only to in whom organic mischief has not yet taken place."

CURATIVE TREATMENT.—Before entering on the ment of this dangerous disease, the medical officer shoutute a rigid scrutiny into all the antecedent circumstances to the soldier's health;—such as, his constitution—age—lesidence in a tropical climate—previous diseases, and e if the present be the first seizure of delirium—habits whether he is at the moment a sufferer from an actual bout, or whether the intoxicating drink has been with two or more days—whether the patient is a habitual and edrunkard. From a careful consideration of these circum together with an examination of the skin—pulse—tongoldition of the nervous system—of the secretions—and of the and degree of the delirium—the observer is enabled redetermine whether the condition of the patient be one of and vascular excitement, or of nervous irritation and exhaust

When the case is not acute delirium tremens from a bout, the patient being young, plethoric, and recently arrived to climate, cold should be applied to the head and fathe body is immersed in a warm bath. On removal is bath, a full dose of calomel and opium should be admit followed in two hours by a powerful cathartic, composed pound decoction of aloes, infusion of senna, sulphate of m and tincture of senna. After the free action of this pure of croton-oil, which many prefer to every other form, it wis cessary to quiet the nervous agitation; and this is best the acute form of the disease, by a combination of tin opium with antimonial wine, exhibited every two hours tranquil sleep is secured.

When the symptoms, as here supposed, are the direct

mediate results of a fit of drinking—no inflammatory action being present within the cerebral cavity—the course of treatment suggested will generally prove of speedy effect. In the true delirium tremens, however, or that resulting from the withdrawal of an accustomed stimulus, with exhaustion of the nervous and vascular systems, the treatment becomes at once more complex and difficult.

Here, as in the former instance, we should commence by the exhibition of a full dose of calomel and opium, followed by a powerful aromatic and stimulating cathartic, with a view to remove retained and vitiated intestinal accumulations, and to act as an energetic cholagogue. After the free operation of the cathartic, opium in moderate doses conjoined with camphor or ammonia, should be given every two hours, until evidence of abated nervous irritability and nervous exhaustion be made apparent, in a tranquil sleep. For the purpose of allaying nervous agitation and sleeplessness, very large, and very unwarrantable doses of opium have been exhibited; but it should at all times be the object of the attentive physician to obtain his objects by a moderate use of direct narcotics; and happily, in the use of chloroform, internally exhibited, we possess a powerful and harmless means of commanding nervous perturbation and depression. Chloroform, in very severe cases, may be conjoined with small doses of laudanum, and so as to command sleep. Chloroform is thus valuable in itself, and as enabling us to dispense with large and repeated doses of opium.

As stated in the article on fever, Doctors Corrigan and Gordon, of Dublin, have, for some time past, been in the habit of using chloroform in delirium tremens. One drachm is given by the stomach at first, and it is often increased to two drachms, or even larger doses. The dose is repeated every two or three hours, until sleep results; and, unlike opium, this medicine does not lock up the secretions.

But to aid the actions of opium and chloroform, the use of active cholagogue purgatives, of warm baths, of spiced negus, and of well made soups, will materially advance the solution of the symptoms; by such means we emulge the depurative organs, while we support

the nervous energy.

Personal restraint and coercive measures are seldom called for, and should not be used without a necessity. The disposition of the patient is generally harmless; and by careful and constant attention on the part of the attendants, and by judicious management, he can be induced to comply with all the requisite orders of the hospital.

In the hands of exclusive practitioners, I have often seen means, apparently opposite in their nature and influences, followed by

successful results:—for instance, a late friend of mine, a of the Bengal army, and one of the most able men whom I have known, entertained a positive dislike of opium, from known quality of locking up the secretions. He treated of tremens with very large doses of calomel, followed by pury and his success was, according to his impression, equal to his brother officers. But this instance only proves how m use of cholagogue purgatives has to do with the treatment disease, and how necessary it is that they should always tute a prominent part in its treatment.

It is here, as in other complicated diseases; it is not favourite remedy, or fancied specific, that disease is to be come, but by the just applications of all the known remedi

The following table exhibits the relative prevalence as tality from delirium tremens amongst the European to Her Majesty's army in the several Commands mentioned:

Delirium Tremens.	Aggregate Strength.	Admitted.	Died.	Prop of d	
West Indies	86661	1426	175	15	
Jamaica	51567	191	42	11	
Cape of Good Hope	22714	13	3	1 i	
Mauritius	30515	514	50	l i	
Cevlon	42978	35	7 9	1 i	
Bermudas	11721	102	9	1 i	
Nova Scotia and New Bruns-	46442	207	18	1 i	
Upper and Lower Canada .	64280	296	18	1 i	
Gibraltar	60269	44	5	1 i	
Malta	40826	38	5	1 i	
Ionian Islands	70293	192	30	1 i	
United Kingdom	44611	27	4	1 i	
United Kingdom Bengal, 1836 to 1840	36286	672	14	1 i	
Madras, 1832 to 1836	31267	496	21	li	
Bombay, 1836 to 1840	18073	113	15	1 i	
Tenasserim Provinces, 1836-37	7000	34	5	1 i	

The above table suggests many inquiries, but which car answered without entering into a variety of circumstances a the moral and physical condition of the soldier.

Why the mortality by delirium tremens in Madras she double that of Bengal, for instance, and why that of Tens

and of Bombay should so very far exceed that of both the sister presidencies, are points of much interest. Again, so far as ordinary tropical influences are concerned, the colonies of the Cape of Good Hope, Mauritius, and the provinces of Tenasserim, enjoy a remarkable exemption, the deaths by fever, dysentery, and hepatitis being few in these colonies, as compared to the other British possessions; yet in these most favoured climates, delirium tremens proves most fatal.

Even within the British islands, we find that, though delirium tremens is comparatively a rare disease, the ratio of deaths to cases treated is nearly eight times as great as in the unhealthy

countries under the Bengal presidency.

It may be that, as in the instance of mercury, a certain amount of injury is prevented by so much of the alcohol being removed from the system by the copious perspirations prevalent throughout India generally, and in Bengal especially, during so many months out of every year. That the excessive transudations which occur in the hot and rainy seasons have such a tendency, appears to me more than probable; for, in the instance of mercury, we have a demonstration of the fact.

Through the same channel of the skin, but by means of active exercise in the open air, the "three-bottle-men," the fox-hunting squires of old, "carried off" their port wine. The hunting during three days of every week kept the depurative functions in activity, and thus they lived on to a lusty old age, while, without such active habits, they must have died in youth or in middle life from visceral disease and delirium tremens—the inevitable fate of the sedentary lot.

## HABITS OF THE BRITISH SOLDIERY, AND THEIR INFLUENCE ON HEALTH.

THE helpless, listless condition of organized bodies of men, as compared to the active, independent condition; the free agency and private enterprise, of civil communities, is a subject which, with its moral and physical results, has not yet been appreciated either by official bodies, or by the public, which pays for all the

labour of naval and military organization.

In civil life, men are very much masters of their actions, and of their habits of life; while in the army and navy a man is neither master of his diet, exercise, clothing, fuel, hours of rest, or of anything else, in short. The government that employs the seamen and soldier is the sole arbiter of his habits and customs, and of everything that relates to his condition. In all this the governing power incurs an obvious responsibility; and it would prove a mournful as well as unprofitable labour to trace, since the days of Charles the Second, how this responsibility has been used for the soldier's and seamen's moral and physical welfare.

The barrack life of the British soldier;—what a world of wretchedness and misery, moral and physical, does that term bring to the memories of all who have witnessed it in our tropical possessions! Here we find men at the mean age of twenty-six, "at the healthiest period of life, and selected from the healthiest of the population," yet "fallen below the health level of their native country, where all the chances are in their favour;" and this, amongst other causes, owing "to the reckless dissipation incident to all military bodies," as we have managed them.

The late Inspector-General of Hospitals, William Fergusson, says:—"I never saw a barrack that was not too crowded, or in other words, where, if the men had been lodged in separate cottages, or even hovels, they would not have been healthier."

"Of the soldiers' life within these barracks there is much to be said, and much to be amended. To take his guards, to cleanse his arms, and attend parade, seems to comprehend the sum total

of his existence; amusement, instruction, beyond the drill, military labour, and extension of exercises, would appear to be unthought of; as it is impossible that the above duties can fully occupy his time, the irksomeness of idleness, that most intolerable of all miseries, must soon overtake him, and he will be driven

to the canteen, or the gin-shop, for relief.

Labour in every shape seems to have been strictly interdicted to the soldier, as water for his drink. All, or nearly all, must have been bred to some trade or other before they became soldiers, but they are to work at them no longer. Labour, the labour of field works and fortifications, strengthens the limbs and hardens the constitution, but that is never thought of in our military life at home; so thought not the ancient Romans, whose military highways still exist, and who never permitted their soldiers to grow enervated in idleness during peace. Better, surely, would it be, that every one should work at his own craft, or be employed on the public works, in regulated wholesome labour, than thus to spend his time in sloth and drunkenness.

But his exercises, without even going beyond the barrack premises, may be made manifold;—running, wrestling, gymnastic games of every kind, swimming, leaping, pitching the bar, the sword exercise, that of the artillery, all that hardens the muscles and strengthens the limbs, should be encouraged; and when the weather forbids out-door pastimes, the healthful exercise of single-stick, in giving balance and power to the body, quickness to the eye, and vigour to the arm, may properly be taken as a substitute for the drill, which, after the soldier has been perfected in his exercise, is always felt to be a punishment. So is the unmeaning

evening parade, and perpetual roll-calling.

Surely, if the soldier present himself once every morning, correctly equipped and in order, the most teasing martinet ought to be satisfied, and then no more should be required than to see that the men are all in their quarters on the beating of the tattoo. Surely the use of the sword has been too much frowned down, as if it had been a forbidden thing. In the night attack, the musket is worse than useless, its fire leading to every kind of confusion; and at the breach it is little better, for it can only be presented against stone walls and ramparts that conceal the defenders; but it would cover the swordsman advancing to the breach, and a couple of chests of ship's cutlasses furnished to every regiment as regimental baggage—a single horse-load—provided the men had been taught to use them, would generally supply all that could be wanted for the exigency of the service.

Let any one reflect on the fearful expenditure of life at the breaches at Badajos and St. Sebastian, and say if some means should not, if possible, be devised, to render it less costly hereafter. One is almost tempted to regret the times "when," according to the old song, "our leaders marched with fuzees, and we with hand-grenades:" and could the good grenadier have carried a sword by his side, to use after he had tossed the ball, he would, I believe, have done much more execution than with a musket and bayonet: and why should the artillery be to him a closed book, as if in the course of his service he was never destined to handle or to suffer from it? A couple of guns, even if wooden ones, in every barrack-yard, with an old invalid bombardier to teach the use of the rammer, and the sponge, and the match, would fill up many a vacant dreary hour, and open his mind to a most useful professional lesson.

The lesson, moreover, would be as useful to the infantry officers as to the privates. He would then, should he ever prove the captor of a prize gun, at least know what it was, and be able to turn upon the enemy the engine that had just been used for the purpose of destroying himself. Every sailor, even on board a merchant ship, where there are no idlers, must become more or less an artilleryman, and why should not the too often idle soldier?

Foot-racing too, the art of running, so little practised, and so supremely useful, should be held amongst the qualities that constitute military excellence. It was so held at the Isthmian games of ancient Greece, and deserves a better place than has hitherto been assigned to it in the military pastimes of modern Britain. In our school books we are told that the youth of ancient Persia were taught to launch the javelin, to ride the war-horse, and to speak the truth. Let the young British warrior be taught to use his limbs, to fire ball-cartridge, to cook his provisions, and to drink water. The tuition may be less classical, but it will stand him in far better stead during every service, whether at home or abroad.

Barracks, from time to time, should be evacuated for purification. The evils and dangers of accumulation will otherwise beset them, inducing disease; and to obviate this, it would be well, whenever practicable, to march out their inhabitants, in the summer season, to the nearest heath or common—always, however, without tents—and there make them hut themselves. No military lesson could be more useful than this. Every man so hutted would be advanced in soldiering to the full instruction of the campaign. The change, breaking the monotony of barrack-life—the novelty, would animate; he would be taught how to live in a camp, how to cook and to forage, to use the mattock, the shovel and the axe.

Tents, when the soldier lies upon the cold ground, with a crowd

of comrades enclosed within a superficially heated atmosphere, loaded with animal exhalations, can only be considered hot-beds for the generation of dysentery. On their return to barracks, they will find everything healthy and refreshed, and they will know

that they have been made better soldiers.

Some have strenuously recommended barrack libraries; and surely, when we think of the dismal monotony that hangs over the soldier in barrack-life, no one with good feelings could object to them. Still, I must confess that I never knew or heard of a reading army. The military exercises and pastimes would seem better adapted to the soldier's character; and I acknowledge I would rather see him a cook than a student, for on that art his very existence may depend; but if he feel disposed to read, let him have every advantage and opportunity that the rules of the service can admit.

Music would seem far better adapted than even books to fire the soldier's mind, for when played in national airs, it awakens a chord which has often electrified armies; and amongst all nations, at some period or other of their history, it has been the accompaniment and incentive to war. The highly civilized English soldier now fights, and can fight, without it; but if taught to feel its powers, would he not fight better with it? To the Irish and the Scotch soldier it still speaks the language of the heart, and the Highlander, when he hears the gathering of his clan blown from the mountain war-pipe, becomes elevated and transported beyond himself; he will then encounter anything in human shape, unappalled by all the forms of death that the engines of war can inflict." This admirable military physician concludes:- "Regular bodily pleasurable exercise has been said to be worth a host of physicians for preserving military health; and occupation, without distress or fatigue, is happiness. The philosopher can make no more of it; and every idle hour is an hour of irksomeness, and every idle man is, and must be, a vicious man, and to a certain extent an unhealthy one; for the mind preys upon the body, and either deranges its functions in a direct manner, or drives the possessor to seek resources incompatible with health."

Thus spoke William Fergusson, after an experience of actual war of twenty years, and thus had spoken before him Robert Jackson; but in everything relating to the soldiers' welfare, Jackson, Fergusson, Hennen, Henry Marshall, and Ballingall, will be found to have been half a century, or more, in advance of all opinion, and especially in advance of military authority. That the warnings of such men should for so long have been unheeded is but another instance out of many of the utter disregard of the subject by the authorities of the state, both civil and military, who for ages regarded the soldier as but a red-coated parial. In France it is

well understood that "the degree of perfection of military medical science is the true measure of the importance attached to the preservation of the soldier."

It was with a rare sagacity and penetration of the subject in all its bearings, belonging only to an officer of the highest order of talent, that Sir John Moore-the intimate personal friend of Robert Jackson-declared, in addressing Sir Ralph Abercromby, that "a Roman army would have gone through their military exercises in the West Indies and have been healthy." The same distinguished commander added :- "The troops, I observe, which have been most active, are the most healthy; a proof that the sun is not the cause of sickness. But in general the greater part of the sickness proceeds from want of interior discipline and economy in regiments.

Great attention should be paid in this country to the cleanliness and even neatness of the soldier's person and the regularity of his diet, an addition to the eating part of his ration instead of rum; sea or river bathing, constant activity and movement. In short, General, (excuse the pedantry of the expression,) but with a Roman instead of a modern exercise and discipline, the troops in

the West Indies might, I am convinced, be kept healthy.

A parade twice a day, consisting merely of an inspection and exercise of arms, is easy for officers; it leaves them what they call more time, but it leaves the soldier also to lounge the whole day in barracks, where the air cannot be good, and where, from indolence, his body becomes enervated and liable to disorder." How much that is preventive of disease do these short sentences enjoin, and how applicable is the entire reasoning to the condition of the European soldier in the East Indies as well. No wonder that, living in the miserable manner here described, condemned formerly to a residence of twenty years in the East Indies-moving from bad stations to worse-hope should shed but a feeble influence over the soldier's enjoyment or future prospect, and that he should thus yield to the temptation of momentary gratification.

Robert Jackson states, that "Planters, who may be said to work hard, experience good health comparatively; even soldiers are healthy when actively employed; they suffer when they remain immured in barracks in ease and apathy." Again :- "The writer ventures to say, by fair induction from fact, that if the soldier in the West Indies, instead of being restricted from labour, were permitted to do for himself whatever he is capable of doing, his health would suffer less than it now does; the mind would be occupied, there would be satisfaction, and fewer of those causes of temptation which in idleness lead him into error."

The most complete army of modern times—in health, discipline, and equipment, was that mustered by Napoleon in 1805, on the shores opposite to England, and which a popular historian declares to have "acquired a degree of perfection, in point of discipline, organization, and military habits, unprecedented since the days of the Roman Legions."

This notable army amounted to one hundred and fourteen thousand combatants; and it is highly deserving of remark, that its exemption from disease was ascribed by its incomparable leader, to habitual light and cheerful occupation of mind and body,

short of fatigue.

"Constant employment," says one of his Generals, "was the true secret both of their good health and docile habits; neither officers nor soldiers were ever allowed to remain any time idle; when not employed in military evolutions, they were constantly engaged either in raising or strengthening field-works, or in levelling down eminences, draining marshes, or filling up hollows, to form agreeable esplanades in front of their habitations, and where their exercises were performed." The same army, according to Tardien, marched from the sea-coast without halt to the field of Austerlitz, over nearly 400 French leagues, leaving hardly any sick behind.

What the "interior discipline and economy" of Marlborough, England's greatest commander, may have been, I have no means of knowing; but there must have been something great in that order which, according to one who served under him, made his camp resemble "a well governed city. Cursing and swearing were seldom heard among the officers; a sot and a drunkard were the object of scorn; and the poor soldiers, many of them the refuse and dregs of the nation, became, at the close of one or two campaigns, tractable, civil, sensible, and clean, and had an air and spirit above the vulgar." How melancholy to look on the reverse of this picture, and to reflect that, in the middle of the nineteenth century, so much should remain to be done to render the British soldier healthy and happy. On the abuse of ardent spirits in our various colonies it were almost in vain to speak. Before that terrible vice can be overcome, something far more powerful than medical reasoning on facts, or the warnings of experience founded on them, must be brought into active operation. Discipline must still further alter its direction—in place of being active only to punish wrong, it ought and must be exerted further and further in the encouragement to good conduct.

Henry Marshall, speaking of the British soldier in the East Indies in his day, says:—"By the daily custom of imbibing spirituous potations, a new want is created, intemperance is established as a habit, and frequent intoxication is the consequence. The wretched drunkard must now have a large supply of liquor in the morning to recover him from the effects of the quantity drunk on

the previous night. He perhaps has neither money nor and his clothes are then sold at a small portion of their Some do not stop here; for, after having sold all their of they will rob their comrades, and with the proceeds of the honesty, provide the means of intoxication. Confinement upon confinement, court-martial upon court-martial, and present upon punishment, until the worn-out wretch dies in hof the "horrors," fever, or dysentery; or if he should for resist the fatal effects of disease, his constitution becomes down by the combined influence of the poison of spirits, hausting climate, and repeated attacks of illness, so that in years he is found unfit for further service in India, and he home to be invalided.

Death is the last, but perhaps not the worst conseque intemperance. This description of the life of a British in India is by no means highly-coloured. But the evil defall on the heads of the unfortunate sufferers; military dis in all its branches becomes deeply affected by habits of perance. To the generally prevailing vice of drinking be attributed almost every misdemeanor and crime con by British soldiers in India.

The catalogue of these, unhappily, is not a scanty one; rapid steps, first from petty, and then more serious neglectinattentions, slovenliness at, and absence from parades, disobedience of orders, riots, and quarrels in barracks, a from guards and other duties, affrays with the natives, the selling of their own and their comrades' necessaries, roll abusive language, and violence to non-commissioned office last of all desertion, mutiny, and murder may be traced source. This frightful picture is not exaggerated. I hat thirty-two punished men in a regimental hospital at one perhaps not a single individual of that number suffered for which was not a direct consequence of the immoderate spirits.

I recollect attending to the punishment of seven men same regiment, who received among them four thousar hundred lashes. They had been all tried for crimes arisin habits of intemperance. Since the institution of the Recond Supreme Court, at Madras, no less than thirty-four soldiers have forfeited their lives for murders, and most of were committed in their intoxicated moments."

How, it may be asked, can courts-martial and officers i mand, altogether reject the excuse of intemperance, so lo soldier is furnished by Government with the means of de him of his reason as part of his daily subsistence, all of w is directed to use? Becaria is of opinion that the punish

crime cannot be just, or, in other words, necessary, if the laws have not endeavoured to prevent that crime, by the best means which times and circumstances would allow.

We instil the moral and physical poison with one hand, and hold out the lash with the other, as the antidote against indulgence. Coercive measures have at all times been found very ineffectual for preventing the evils arising from intemperate habits, partly because the soldiers do not generally regard drunkenness as an ignominious vice. Where punishment has no influence in checking a man from repeating a breach of discipline, or in preventing others from committing a like offence, it is worse than useless.

Where endemic fevers prevail, the temperate, it is true, are attacked as well as the intemperate; but it is universally admitted by medical men, that the former has a much better prospect of recovery than the latter. Desgenettes, in his medical history of the French army in Egypt, observes that daily experience demonstrates that all the soldiers who indulge in intemperate habits, and that are attacked with fevers, die; nay, we may go still further and say, that they have been more liable to an attack of disease. Sir James McGrigor repeats the same observations even more emphatically.

The obvious purport of Henry Marshall's able report is, to recommend an abolition in the East Indies, and everywhere, of the indiscriminate and uniform issue of spirit rations to the troops, with a view of abating the extent of crime, of lessening the proportion of sick, of reducing the rate of mortality, diminishing the numbers discharged on account of disabilities, and of promoting

the welfare, efficiency, and discipline of corps.

But there is another habit respecting which I shall venture to say a few words, because it is both a bad one, and a comparatively new one,—I mean the immoderate use of tobacco—a habit brought amongst us from the continent of Europe, on the cessation of the French revolutionary war. Young military men are apt to regard the habit as a manly one, until severe dyspepsia, shattered nerves, sallow complexion, disturbed action of the heart, and other symptoms, show themselves, and then it is frequently too late to stop. "The sallow complexions, black, broken and unsound teeth" of the Germans, are matters of notoriety to all travellers. "You may," says one of them, "smell a German in any part of the room, or scent him at a quarter of a mile's distance in the open air, if the wind be favourable."

Much is talked of the good effects of tobacco-smoking in damp localities, by persons who, in defiance of geographical differences, carry the habit wherever they go—from the marshes of Aracan to the arid plains of Delhi; but I think there is good reason to ques-

tion the benefits of this habit of smoking even in the fa fog and damp, or that tobacco ever acts as a preventa disease, and least of all to fever.

The truth is that many persons puff themselves into graces of snobs and spoonies like themselves, and use the score now, as Lord Chesterfield drank and smoked notwithstanding his aversion to wine and tobacco—"I thought such practices very genteel, and made him I man." How his lordship may have looked, under the fluence of wine and tobacco, his biographers have failed but we all know how our modern "spoonies" and "snot thoroughfares look, after a course of cigar-smoking alor

Damp and cold climates may confer a greater toler partial immunity from, the evil effects of the drug, but haps all that can justly be admitted. Dr. A. T. Thoms ing of the medicinal effects of tobacco, says:—"I power, when it is employed in excess, weakens the organs, obtunds the nervous sensibility, and depresses vital energy."

Of the empyreumatic oil of tobacco, which is priminute quantities in the ordinary process of smoking author says that, when introduced into a wound, "it can death." Of the other active principle, the nicotina, he it is a direct sedative; and that, medicinally, under a stences, tobacco must be employed with the utmost caut

If such checks and cautions are necessarily put on this powerful drug, in the hands of the physician, who added on the continuous and unlimited use of it by the and ignorant soldier.

Of hookah-smoking I need say nothing, as happily nearly gone; but I have seen many cases of severe con and cardiac disturbance from its abuse, with perfect r health on the discontinuance of the habit; the digestive those of the heart and nerves, having been seriously the most inveterate smokers. Of the miseries, mental a which I have witnessed in the persons of young officers abuse of cigars, I will only say that they very far exception of the confessions of an Opium-eater."

Many persons flatter themselves that by long use s become a harmless second nature—the truth being that never become a second nature, for they have nothing the first, as has been said of graver matters.

Lallemand has reported, on an extensive personal of that the use of tobacco deadens the generative function convinced," he says, "that this must be much more free is generally believed, if I may judge by the torpor into organs of generation fall, as soon as the narcotic action of the tobacco is experienced, and by the habitual indifference evinced by

confirmed smokers for the society of women."

Professor Lizars of Edinburgh, after enumerating the modes in which syphilitic ulcerations of the lips, mouth, and throat are propagated by the short-pipe and cigar, and how carcinomatous ulceration of those parts is produced by the same cause, states that the constitutional effects of tobacco are numerous and varied, and occasionally truly deplorable. Amongst these he enumerates; dyspepsia, vitiated taste, a loose condition of the bowels, congestion of the brain, loss of memory, amaurosis, deafness, nervousness, palsy, emasculation, and cowardice, or want of

moral courage.

Dr. Prout states that "tobacco disorders the assimilating functions in general, but more particularly, as I believe, the assimilation of the saccharine principle. I have never, indeed, been able to trace the development of oxalic acid to the use of tobacco; but that some analogous and equally poisonous principle, probably of an acid nature, is generated in certain individuals by its abuse, is evident from their cachectic looks; and from the dark and often greenish yellow tint of their blood. The severe and peculiar dyspeptic symptoms sometimes produced by inveterate snuff taking are well known; and I have more than once seen such cases terminate fatally with malignant disease of the stomach and liver. Great smokers also, especially those who employ short pipes and cigars, are said to be liable to cancerous affections of the lips." Dr. J. P. Murphy states most truly that there is nothing more injurious to a flaceid heart than smoking, many cases being traceable to this cause alone.

Were I to relate but a small portion of the results of my personal observation, as to the effects of the abuse of tobacco, I might be suspected of exaggeration, I therefore prefer to place before the

reader the results of other men's observations.

It must not be supposed from what is here stated that I would place the stout British soldier on a diet of vegetables, treacle, and tea: far from it, I would take nothing from him which constitutes a liberal, nutritious, and wholesome diet; not even a moderate spirit ration to the seasoned soldier—especially when in work or in full exercise. Cooped up in barracks, however, the present plan of dieting in the East Indies is too uniform and unvarying, and much too highly stimulating; and the spirit-ration constitutes "a pernicious bounty" to the men. The entire of the dieting of the army requires revision and regulation so as to adapt it to the wants of the soldier under the various climates in which he is called upon to serve.

William Fergusson justly observes, that a great physiological

principle seems always to have been overlooked, and that natural appetite for change and variety. "It is ever the and no man, even if he will, can be satisfied with this stomach and digestive organs will be heard in their own and if they be not attended to, their owner will fly to in solace of his disappointment. There is a mistake her if we wish to wean the soldier from drunkenness, we shou careful to place within his reach more wholesome indulgence which a diet suited to his taste (and it cannot be suited w variety), must ever be the first. Were he allowed to cater own mess (always, however, under due superintendence) can be little doubt of his better relish for his meal, besides ing that most useful of all lessons, the art of subsisting hi In the case of the soldier, to cook is to live; for if he cannot pare his food, he will be poorly fed, even with flocks and he his command." . . . "In all this the soldier our have been trained and supported by the wisdom of the co But has it been so?"

These most excellent because practical suggestions, if evince less learning than some chemical rules of diet which seen, nevertheless contain more knowledge. They are also important and interesting in another sense, as showing he the author was in advance of what is called professional or While naval and military men, scouting and scorning all or but their own unauthorized dicta, and asking the old worn-out tion—" not being a naval or military man, what can he know of or military affairs," we have here a most eminent and experi army surgeon calling upon "the wisdom of the country" to and support" the soldier. The miseries and horrors Crimea, with their disastrous results, have at length rouse people to a sense of the national danger; and the common of the country, guided by the common interests of self-pre tion,-in other words, "the wisdom of the country"-is br at length to bear on naval and military organization-not, how before it had become an urgent state necessity.

Referring, in conclusion, to the more especial subject of article, we find that Dr. William Fergusson, whose "Note Recollections of Professional Life" were published in 1846, — "The exceeding vulgarity of the prejudice that ardent simpart strength and vigour to the human frame, is disgrace educated men; yet true it is that many of our most experi commanders of the army and navy still attempt to justify continue the practice."

ON THE SELECTION AND IMPROVEMENT OF LOCALITIES FOR THE EUROPEAN TROOPS IN OUR INTERTROPICAL POSSESSIONS: LOCALITIES TO CHOOSE—LOCALITIES TO AVOID, &c.

It has been justly observed by an able military surgeon of old, that "where the hygiene of an army is judiciously regulated, the soldier may be kept in health and vigour; but allow an ignorant general to encamp on a marsh, let filth stagnate, fatigue excessively the men, crowd them in low damp rooms, and, despite of drugs, they will fall as unripe and blasted fruit, not by the sword, but by the fever." Frederick of Prussia has stated that fever cost him as many men as seven battles. What between the fever and the battles together, it must have fared badly with "the tyrant's counters" in those days. It were well if we could be assured that in our time the external causes of fever have become less rife in our camps, and that our commanders by sea and land have become more vigilant and better informed as to the means for preventing them; but I fear that neither can be affirmed with any confidence. As to commanders, we would appear often to have clung to men ignorant of, and disinclined from learning the most simple rules affecting health, as to a prescriptive right.

We have lost an army in every quarter of the globe, and precedent has consoled us: it has always been so in our campaigns, and in our colonies. We have often appointed unproved men, and men of proved insufficiency; and we have retained them in their places, because it has always been so with us. But a better order of things is at hand: in truth matters could not continue

much longer as they have been.

To the neglect of healthy localities for camps, cantonments, and stations—to the absence of proper structural arrangements in barracks and hospitals—to inattention to the suitable feeding and clothing of the men—the neglect in promoting and cultivating better habits of life—the absence of instruction in cooking, hutting, exercising, and bathing—to the neglect of change of air and of climate in the remote stages of the diseases of seamen and soldiers;—to these neglects, which it must at all times rest with

governments and commanders to remedy, we must refer the stant and terrible losses of the British army, in our intertropossessions.

I venture to say that, from the causes here briefly enum we have in every score of years during the last century as wards, lost unnecessarily, in our fleets and armies, more me have fallen in our battles from the landing of Julius Cæsar

present time.

Colonel Tulloch has stated to me, as the result of his invitions in the War Office, that of British soldiers of the Queer Company's armies, in the East Indies alone, there perished cessarily, from 1815 to 1855 inclusive, about one hundred the men. "During the years," he says, "between 1815 and there died of Europeans belonging to Her Majesty's and the India Company's army in India, very nearly 100,000 me greater portion of whose lives might have been saved, had localities been selected for military occupation in that countries."

Nor are the causes of such destruction of life difficult of discovered and prevented. Robert Jackson says that, "The instances where the sick-list in armies amounts to one-third total force, and others, where it does not exceed one-fiftieth one-hundredth part. The causes of such difference are som visible and obvious. To discover their sources, and to p their operations, is important; and it is frequently a work great difficulty."

Samuel Johnson, referring to the contests of his time with F and Spain, states, that "the wars of civilized nations make slow changes in the system of empire. The people perceive seany alteration but in an increase of debt; and the few individuals who are benefitted are not supposed to have the clearest ri

their advantages.

The life of a British soldier, he adds, is ill represented by fiction. War has means of destruction more formidable the cannon and the sword. Of the thousands and tens of thou that perished in our late contests with France and Spain, small part ever felt the stroke of an enemy; the rest languin tents and ships, amidst damps and putrefactions: pale, to spiritless, and helpless; gasping and groaning, unpitied a men, made obdurate by long continuance of helpless misery were at last whelmed in pits, or heaved into the ocean, we notice and without remembrance.

By incommodious encampments and unwholesome star where courage is useless, and enterprise impracticable, flee silently dispeopled, and armies sluggishly melted away. T a people gradually exhausted, for the most part, with little e . . . In the last war Havanna was taken; and at what ex is too well remembered. May my country be never cursed with

such another conquest.

The attack on Carthagena is yet remembered, where "the Spaniards, from their ramparts, saw their invaders destroyed by the hostility of the elements; poisoned by the air, and crippled by the dews; where every hour swept away battalions; and in the three days that passed between the descent and re-embarkation,

half an army perished."-POLITICAL TRACTS, 1771.

Nor can these events be regarded as matters of a bygone history; for Walcheren, the Peninsula, Arakan, Rangoon, China, Sindh, Varna, and Sebastopol, have brought them back to our very doors, and they have occurred always from the same causes. Sir Gilbert Blane, speaking of the neglect of the seamen's health in his earlier experience, says:—"In former times they had not the attention paid to them which would have been due even to inanimate machines of equal utility; for there seemed to be much more anxiety about preserving arms from rusting, and cordage from rotting, than about maintaining men in an effective state of health."

We have, in the instances cited, witnessed our armies scourged and wasted away as of old, through the ignorances and incapacities of authorities at home and abroad; but a higher order of intelligence is drawing upon us, having its origin in the sanitary investigation of our naval and military surgeons. "The wisdom of the country," too, invoked to her soldiers' aid by men like Robert Jackson, William Fergusson, and Henry Marshall, is being manifested through a free press, and a Parliamentary inquiry;—sharpened, perhaps, by a sense of the common danger resulting from abuses and ignorances without end. We are said to have acted hitherto as if armies had been made for generals, and not generals for armies; but let us hope that the time is not distant when soldiers and seamen shall cease to perish without necessity, and without results.

Let us hope also that Englishmen may not again witness, as in the instance of Walcheren, the degradation of the sovereign's grant of "approbation" for "perseverance and celerity—for the promptitude with which he had commenced, and the vigour with which he conducted" and consummated the destruction of an army. One such humiliation—one such shameless laudation of a commander who ought in justice to have been cashiered by sentence of a court-martial—ought to be enough for one century.

The pestilential island of Walcheren received the largest expedition which had ever, up to that time, left the shores of England, amounting in seamen and soldiers to seventy thousand

The climate into which this most powerful and admirably disciplined force was thrust—during the months of its utmost insaludious encampments, and unwholesome stations, where courage is

useless and enterprise impracticable."

Dr. William Maclean, of the medical department of the Madras Army, writing in 1854, says in reference to India generally: "Barracks have been built, and costly repairs wasted on others that for half-a-century have been mere pest-houses. Our cantonments, for the most part, are to this day in a most disgraceful condition, well-known causes of disease being rife and abounding on every hand; and all for want of some person, or body of persons, who can speak out on the subject so as to compel a respectful attention." But to return to our narrative:—

It had long been subject of complaint and remonstrance on the part of Robert Jackson and other military surgeons, that, in the West Indies, facilities of accommodation for trade—the polar star of British policy—had condemned the soldier to the scourge of disease without a necessity; and whoever looks at the distribution of the European military force in the East Indies will come to the conclusion that the soldiers have been exposed to disease there on grounds even less justifiable than have held sway with

the authorities in the West.

"It is positively true," says Robert Jackson, "proved to a demonstration, in numerous instances, but proved by accident, not by avowed experiment—that European troops may be so stationed in the Islands of the West Indies as to retain health nearly as well as they could be expected to retain it in their native country. The air of the interior and mountainous parts of the larger of the intertropical Islands is comparatively cool and pleasant, and not unfriendly to the European constitution.

The expense of filling up the military ranks, thinned by sicknesses which arise from bad position and badly constructed quarters, actually amounts, in course of a few years, to a greater sum of money than would be required to erect barracks of the best form of construction at the most eligible sites in the country, even

at the extravagant price of government contract."

The truth of what was here, many years ago, advanced, Jackson proved in a statistical summary of the Inspection Reports of Military Positions, Barracks, and Hospitals, submitted to the Commander of the Forces on the West India station. His representations though so true and so easy of comprehension, met with no attention whatever; and after a public life of the highest honour and of the most noble labour and usefulness, extending over halfacentury—every day of which, during peace and during three great wars, was devoted to the moral and physical good of the soldier, Jackson concludes with a prophetic appeal to "the legislative branch of the government;" there being "a presumption amounting almost to proof that it has no chance of

obtaining it from the executive;" that is, from military or

authority.\*

Referring to the fact that in England "much opposition was by the subordinate authorities," to his suggestions for comstatistical reports on the sickness, mortality, and invaliding a the troops in the colonies, Henry Marshall adds that, rance of the principles of military hygiene, and an inve prejudice in favour of former usages, seemed to be the le cause of the opposition to the measures recommended, the tion of which has proved so singularly beneficial."

Very different was the conduct of the generals of Napoleon the landing of the British forces on Walcheren, and on the ca of Flushing, there was found a Memoir by the French comma General Monnet, on the preservation of the soldier's health was a document full of interest, and of much scientific and p

cal value.

But the removal of the British troops from the pestile plains to the healthy mountain regions of Jamaica was at 1 ordered by a man who united the most active and enlarged volence to the most invincible firmness and resolution in car out every measure that was right-Sir Charles, afterwards Metcalfe. And, strange to say, the result has been to the letter as Jackson foretold; the British soldiery now retaining health in the mountains, "nearly as well as could be expect their native country."

The result as regards the mortality of British troops is, in ple figures, as follows:-From 1803 to 1816, and for how lon fore we do not know, the soldiers perished in Jamaica at the of 130 per thousand per annum, while by the simple and

\* I have already suggested that if, by the usages of war, the punishment of be applicable to a disgraceful capitulation, on the part of a commander, some punishment ought in justice to await the man who, through wilful neglect,

wanton ignorance, causes his army to perish. In these days of canting speculative humanity, and maudlin morality, death m be so much mentioned for the holocaust even of a fine army; yet death ha awarded, and suffered too, for very minor crimes comparatively. Certain am I, the very first Sir Michael or Sir Peregrine who shall be made to suffer the mild sente cashiering for such a crime, under the award of a court-martial, composed of and military officers in equal numbers, will prove the last of his class who will on the same course of criminal neglect in time to come. From the day that this common justice shall be done, we may hope never thenceforth to hear again of being planted in marshes, there to die, while the surgeous had for fifty years urge-removal to the hills, where they would have lived well and happily; of soldiers thrust into barracks and hospitals unfit for the reception of convicts. Let this be and we shall no more be told by a statesman in Parliament that " nearly one half lives that used to be sacrificed in the West Indies, was sacrificed through neglect that "this mortality might have been prevented by the most ordinary precautions "while the soldiers, who were unstained by crime, were condemned to five day provisions weekly, the convicts were allowed five days' fresh meat; and that wh troops were suffering, the convicts were in good health." I say, let the right jud be brought to bear on the right man, and we shall hear no more of such atrocitie arranged measure ordered by Sir Charles Metcalfe, the mortality has, since 1842, been reduced to 35 per thousand per annum.

This beneficent act, so becoming the generous and noble nature of the Governor of Jamaica, was "characteristic of Metcalfe. He made the first movement on his own responsibility, and offered himself to pay the expense. He saw that the troops under his command were being sacrificed to ignorance, negligence, and false economy; and he exerted himself, not without success, to establish a new order of things, by locating on the high healthy grounds of the island, the European regiments which perished miserably on the plains."—Kaye's Life of Lord Metcalfe.

Detachments of British troops were stationed at Maroon Town at different and uncertain times between 1795 and 1839, by direction of Lord Grey and others; but nothing fixed or determinate was arranged until Lord Metcalfe gave the final order. When will any of our great foreign possessions fall under the charge of such

another governor?

Robert Jackson speaks also of the value of military labour in securing military health—a subject not yet taken into consideration by our authorities. Fort King George, he says, in 1803, stood in lee of a swamp, and the exhalations from it being considered by Colonel Macdonald, commanding the Royal Scots Regiment, to be injurious to the health of his men, he ordered them to drain it. "The planters lent the tools—the soldiers of the Royals drained the bog;" and Fort King George, which formerly stood noted for unhealthiness, is now "one of the healthiest quarters in the Windward and Leeward Island Station, as appears by a comparative view of the sick returns of the army."

In Jackson's various writings there are numerous instances of similar results from the gratuitous labour of the soldier; and, what is very important also, this most eminent military physician always concludes by declaring that "the thing was done—with obvious effect upon health, morals, and happiness, while the work was

under execution."

With reference to the purposes for which British soldiers are sent to the East Indies, and to the efficient exercise of their powers when called into action, it has always surprised me that so large a portion of the European army has been retained in the plains of Hindustan. With urgent political reasons or state necessities I do not pretend to deal; but if these do not exist, I know of no defence that can be set up, on the score of humanity or of expense in plea of the cruel and unnecessary sickness and mortality—of "the serious loss in the balance of national accounts," consequent on the loss of men.

"The case is important, and the sovereign power will be false to its own interests, if it do not apply a remedy when furnished with proof that a remedy is attainable, and particularly when formed that it is not difficult of attainment."

But the evils resulting from neglect in choosing proper for our colonial emporiums do not end here. The choice of sites having, as stated, reference to trade only, and not to he the sickness which is ever present is not referred to its true of the inherent unhealthiness of the place, but to the arrival of trading vessel from some other equally insalubrious locality; hence quarantine, with all its evils.

Sickness is always prevalent in such places—ships are al arriving;—and people always dislike to believe that they living in a pestilential locality;—so, disease is readily asce to importation. With the unreasoning many, the supposi foreign ill is always more dreaded than the real home-bred pe

In the East, as in the West Indies, the loss of military life endemic and epidemic diseases has always been unnecess great—sometimes enormous. "If the subject be viewed corr and without prepossession, the loss sustained will be foun have been principally owing to mistake, that is, to inatter to truths furnished by experience for the choice of healthy sitions."

These just and manly sentiments are quoted from the celeb work of Robert Jackson on the "Formation, Discipline, Economy of Armies," and especially from that portion which is devoted to a general view of service in Tropical Clim wherein, though he speaks in special reference to the West In Colonies, there is necessarily much which applies with equal tice to all hot countries. This admirable work ought to be in hands of every officer, whether military or medical.\*

I may here mention that I prepared, by desire of the late Metcalfe, without previous notice, a memorandum relating to arrament for constructing barracks and hospitals, and which he has to Lieutenant General Sir George Murray at a meeting, hel desire of Lord Stanley at the Colonial Office, on the 16th of 1842. This note was of a purely suggestive character, no having been allowed me for entering into details; and the foling quotations are here presented:—

1. "The author of this note has satisfied himself, from an tensive range of observation in the East Indies, that great lo

<sup>\*</sup> The great military work of Robert Jackson, is not sufficiently known to the I or to the Indian army; and as I am desirous that military men should have a warrant for its excellence than my opinion, I subjoin that of one of the highes brightest of military critics—that of Sir John Moore:—

<sup>&</sup>quot; My dear Jackson,

I have perused your military book, with which I am much delighted. are none of us, even the most experienced, who may not derive instruction from it I only regret that you, who possess a soldier's mind, had not been a soldier b fession."

life, injury to health, and sacrifice of the public money, have resulted from the present modes of locating and constructing barracks and hospitals for the European troops there; and his information, derived from printed and other sources, leads him to conclude that similarly injurious results are also produced in various of her Majesty's colonies, from similar causes.

4. To obviate for the future such extensive waste of life and money, and with a view likewise to improve such existing buildings as may admit of amendment, it is proposed that standard or model plans be prepared for barrack and hospital accommodation, in scale proportioned to certain given numbers of men, and suited to the various climates occupied by British troops. The most cursory observation will satisfy us, that of certain given plans one only will prove excellent; whereas now, all over the British Possessions, we find in existence every conceivable plan, ending in no approved or fixed plan whatever; so that, in fact, that which ought to be settled, and is very easily capable of being settled by regulation is, up to this day left, as I have stated, to the fancy of individuals.

5. It is not apprehended that either difficulty or the expenditure of much time would be encountered in carrying out the proposed plan, provided it were undertaken by persons competent to the task; for the same plan of barrack and hospital accommodation would suit the climates of the East and West Indies, the Cape of Good Hope, Mauritius, Ceylon, China, and Australia—the Stations of Gibraltar, Malta, and the Ionian Isles—while plans for the other colonies would be easy of settlement.

6. To arrange and complete so desirable an object, it is respectfully suggested that a committee of engineers and medical officers be assembled in London, and that power be given, in furtherance of the purposes in view, to call in evidence such persons as may be supposed to possess experience and a knowledge of the subject."

Sir George Ballingall, in the last edition of his admirable "Outlines of Military Surgery," says of the memorandum just quoted:—"This excellent suggestion has not been as yet acted upon, and consequently no established plan has been adopted for the construction of barracks either at home or abroad."

It is impossible to overrate the influence of structural arrangements, of an abundant supply of pure air and water, whether in military or civil communities. Dr. Farr has the following table of mortality in town and country throughout the changing seasons of the year. It is derived from the returns of the ten years 1843—52; and contains comparative estimates of deaths in towns and in the country, concluding with statements of the numbers destroyed in towns by matters which are poisons in houses, streets, and streams:—

	Average annual number of D				
in the months of	Deaths to every 10,000 persons living in Towns.	Deaths to every 10,000 persons living in the Country.	Lives the m are house strea fertilis		
January, February, March	69	56			
April, May, June	62	52			
July, August, September	63	46			
October, November, December .	64	49			
The Year	258	203	1		

"The population of England is, there is reason to be lectively healthier than any equal amount of population other kingdom; but the rapid increase in the proport town population,-in which the mortality is 27 per ce than it is in the country, and the sickness, the suffering lity, the physical degeneracy of race are in an equal makes this question of the health of towns and the ferti the surrounding fields one of the great questions of th manding immediate solution. It is difficult for the in to conceive all the beneficent effects that would flow possible diminution of the mortality which the above i press."

The practical question may now be asked,—have we our Eastern Possessions, almost everywhere, mountain veniently situated wherein the European soldier mig health and vigour through the advantages of a better clima proper application of his own labour; and, if so, why ar made use of? That such favourable localities abound t India, no one can doubt; and I am disposed to hope calling the attention of authority to their sanitary advathe benefit of all classes, civil and military, will prove the many valuable results which may accrue from the de systematic plan of requiring from military surgeons India, Reports on the Medical Topography of the Co nerally, and of its Districts, Stations, and Cantonmen ticular.\*

<sup>\*</sup> The plan here referred to was a measure submitted by the author, of March, 1835, to the Governor-General of India, Sir Charles Metcal sanctioned and ordered by him for the three Presidencies of India and the

Such courses of investigation can hardly fail of important sanitary results. It is probable that some of the lower ranges of hills may prove available for military stations, and that medium degrees of elevation may be found which, while not so lofty as to cause bowel and other disorders, may yet prove of sufficient height to place the soldier above fever-level. This would be an important topographical discovery. Maroon Town, Jamaica, is elevated but little more than two thousand feet above the level of the sea; yet this has proved a sufficient protection against the deadly fevers of the country; and even against the yellow fever, a rise of 2,500 feet is everywhere found to be a sufficient protection.

The same security is obtained in Constantinople and in Cairo, in the instance of plague, by the elevation of Alem-Daghe in the

former, and of the citadel in the latter.

Nor are the moral influences of the mountain climates to be disregarded by the medical philosopher. Montesquieu and Malte-Brun have assigned topographical reasons why the inhabitants of the plains should of necessity be weak, effeminate, idle, and timid, while the elevated and cold regions are inhabited by their conquerors;—regions wherein the severer forms of fever are unknown. It would seem, indeed, that the diseases of tropical climates, like certain vegetable productions, are restricted to certain altitudes and particular degrees of temperature.

"If British Colonies," says Horace Wilson, "be ever formed in the East, with a chance of preserving the moral and physical energies of the parent country, it is to the vales and mountains of

the Indian Alps that we must look for their existence."

The removal of troops even to short distances from their stations and encampments has, I am happy to learn, been acted on systematically in various of our colonies since the publication of the Army Medical Statistics, and in every instance with a most satisfactory result. I trust this important subject will continue to be present to the minds of both military and medical authority, and that no narrow contracted notions of economy will anywhere be allowed to stand in the way of a duty imposed alike by considerations of humanity, and of the best interests of the State.

I had continually to treat the ailments of men of business in Calcutta, which I often found to be of an anomalous and perplexing nature, affecting both mind and body. In such cases mere medical treatment very often proved vain, until the sufferer was released from the cares of official duty, by a trip to sea (which I

to the Eastward. The order bears date 23rd November, 1835, and was a *direct* act of the Government—the medical authorities of Bengal having afforded the measure a very equivocal support, or rather no support at all.

This great question is now in the hands of a highly-intelligent body of medical officers;—the field is almost untouched before them, and we must not doubt the eventual results, whatever obstruction may be offered by the vis inertiae of old men.

always preferred), or by a removal to the Upper Provindeed, where the circumstances of the patient admit of it resource of medicine should never be neglected. It general tonic, invigorating the whole frame, improving to the blood, and stimulating every organ to a more her formance of its office.

Of the advantages of an apparently trivial change of p health we have a very interesting example in the ins Augusta in the southern division of the United States. 1829 this arsenal was situated on the Savannah, and vailed to such an extent that, in the third quarter of 18 garrison except two men suffered from it; while, by a 1 Sand Hills, three miles from Augusta and two from the River, where the soil was hard, dry, and sandy, with m in the vicinity, the ratio of cases of intermittent fever was to 150, and of remittent fever to 156 per 1,000 annually and Foreign Medical Review, July, 1843.

The selections of localities for barracks and hospital generally faulty throughout the East Indies, while the conference of the Company for the accommodation of their troops," nant, "have cost several millions, and may be regarded a instances of their economical principles giving way to their army. The sums thus expended, laid out at pound interest of this country, would, at a determinate very remote, have equalled the national debt."

The buildings for the European troops at Berhampore lately abandoned on account of the unhealthiness of t are stated to have cost, including capital and interest, mous sum of £16,891,206. The details are from the of Accountant-General, and that of the Military Board, Ca

Happy the condition of the European soldier in the the West Indies, had a tithe of the sums wasted in ere racks and hospitals in low and marshy sites, been appliposes of preparing less costly but yet suitable accomm positions of ascertained salubrity; but, instead of that everywhere throughout our intertropical possessions, occasionally of good construction, erected at enormous of

* The military buildings of Berhampor year 1757, at £302,278, or Rupees	e were est	imate	d, in	the	3,8
Compound interest on this sum for 77 y Average annual charges for repairs, &c.		year	rs, at	the	1,3
	Total				168,9

Exchange at 2s. per

are eventually abandoned because, from their improper positions, and proved insalubrity, their inmates are found to die in

a proportion to shock even the humanity of Governments.

By a table furnished me by the late Inspector-General Macleod. the average annual admissions into hospital during thirteen years, at Berhampore, was 2,196 per 1,000; while the deaths averaged 106 per 1,000 per annum. The proportion of deaths to cases treated in hospital was, during five years, as follows :-

In Fever, one in . . . 21 and a fraction. , Dysentery, one in . . . 10 , Hepatitis, one in . . . 9

Secunderabad, Madras Presidency, now also in course of abandonment, has long been noted for its bad locality-bad barracks and hospitals-all which have long operated destructively on the health of British soldiers; yet it is surrounded by stations which are noted for their healthiness. The barracks are described by Dr. Burke as "composed of two squares enclosing one another, so as to make assurance doubly sure against the possibility of perfect ventilation;" and added to all this we have defective drainage and sewerage.

In no station in the south of India, the old abandoned one of Massulipatam excepted, has the mortality been so great; and it is produced, as might be anticipated, by fevers and dysenteries of formidable characters, principally :- in short, it would be difficult to determine which is most to be lamented in this unhappy station-the defects of nature or those of art. There died at Secunderabad, between 1804 and 1835, the large number of 2,620 men and officers, women and children, inhabiting the bad

locality and buildings described.

According to a return made by the Madras Medical Board, the average annual deaths amounted to 84.89 per 1,000 of strength, or 96.89, according to other corrections. These are melancholy reckonings; but they will continue to be the rule of the British army-"until," according to Robert Jackson, "physicians have the place in the councils of military commanders that is due to science. The health history of the late wars in Europe is demonstrative in proof of the important fact that military life has been sacrificed in an enormous proportion to ignorance; that is, to the unwillingness of commanders to be advised on subjects which they could not themselves be supposed to know."

So serious a charge as this against the commanders of his conntry, by one so far-seeing-so just-so learned and experiencedso humane and virtuous as Robert Jackson, is deserving the most serious consideration. The declaration of this celebrated army physician was not only true when it was made, some fifty years ago; but it remains emphatically true to the present hour, not-

withstanding the unremitting labours of such men Blane, Hennen, Trotter, Fergusson, Ballingall, and shall, and their endeavours to bring about a more

just, and humane treatment.

While all the horrors related by Earl Grey, in his speech in the House of Lords, on the Administration 8th April, 1854, were in full and destructive existen of Wellington was all-powerful in this country:—he General of the Ordnance, Commander-in-Chief, Pri&c.;—but if he had held no official station during the of peace—the time for perfecting military institution improving the condition of the soldier—his power to that he been so minded), was without limit. That move more actively in the cause of improvement, or can do so, must always be subject of regret to those who ciate the Duke of Wellington's military services in according to the soldier.

During the period of his active field career in I Europe, he rendered vast services to his country; bu long years of peace which followed, and when the mil tions of the empire, as well as the locating, housi feeding, and arming of the soldier, might have been most appropriately perfected, "the Duke" did not suff sider those easy, obvious, and incumbent duties, for to which his previous course of life and experience in have been supposed to qualify and prepare him.

In place of pursuing a course so becoming, and on so natural, his Grace took to politics, in which he co ceed:—he not only did nothing, or but very little, can discover, to improve the moral and physical cond soldier to whom he owed so much, but he is said to h many vitally important suggestions for the improve military organization and administration.

Here we perceive, for the hundredth time, how much to the soldier's welfare—how much for good or for ev

on the character of the commander.

If our commanders will not be instructed betimes—not be made to look before them and pay a just rewarnings of the medical staff of the army—it is quit range of possibility that the soldier may some day be take the question into his own hands—for this is esoldier's question—and thus bring it to a summary One month's arrest of the voluntary enlistment, and marrive at a most unpleasant issue.

If we would look well to our recruiting, and to the r of our old superiority by sea and land, we must bring t

question to "an immediate solution."

The disasters of the Crimea have at length opene

eyes of the public to some of the necessities of the case. "The worst waste of all," says a popular writer, "the waste that can least be afforded—the waste the most extravagant—is the waste of the soldier's life; and no means which can contribute to keep him in health and condition should be spared. The best economy now is to make the most of our men, and to furnish freely whatever will contribute to their health, their strength, and their efficiency. For this they will not fail to make a return in action which must conduce to the restoration of peace."

The greatest of all commanders was most careful of military health; and being Emperor as well as General, he ordered so that all the arrangements of the military hospitals were excellent throughout the armies of France; the historian, Napier, declares that Napoleon "furnished his hospitals with all things requisite,

even with luxuries."

He also knew how to appreciate and reward the difficult and dangerous services of the military surgeon: he acted on the principle urged by Robert Jackson, that as "the medical staff shares the fatigues and dangers of war, in just reason it is entitled to a

share of advantages."

Napoleon designed that there should be rewards for every rank—from the common soldier to the field marshal; and he bestowed the highest grade of the Legion of Honour on the field of battle, on Barons Larrey, Desgenettes, and Percy. The commanders of the French army considered that the Legion of Honour derived lustre and renown from the enrolments of such names in its ranks; while, in the British army and navy, neither mental sympathy, nor material aid, has ever come from our admirals or generals—always excepting the honoured names of Abercromby and Moore, the friends of Robert Jackson, and of Sir Howard Douglas, and Sir De Lacy Evans, in our own day.

It would seem to have resulted as a matter of course that, from the systematic neglects of the army and navy, the friends of the seamen and the soldier—the Trotters and Jacksons of our fleets and armies—should also have met with ill-usage:—persecution was the persevering lot of Jackson, while the equally long and meritorious services of Trotter, on his element, were treated with a cruel neglect. It was not thus with their contemporary, Larrey, whose memory received in France the highest honours but the

other day.\*

<sup>\*</sup> MILITARY SURGEONS IN FRANCE.—"In last week's 'Lancet' we gave some details touching the inauguration of the statue erected in honour of Baron Larrey, in the great entrance court of the Val de Grâce, in Paris. We subjoin the speech of M. Dupin, President of the National Assembly. The President spoke as follows, in a very impressive tone of voice:—

I have the honour of addressing the military surgeons of the French army, are you—if any one were bold enough still to dispute your right to proportionate

If in France, "the degree of perfection of military in science, is held to be the true measure of the importance at to the preservation of the soldier," so, in England, the neglithe authorities, of the science of military medicine, and of who, under such great discouragements, so ably and disint edly professed and cultivated it, may be taken as the true most the disregard shown in this country for the soldier's heal comfort. Such it has been for ages with us: our soldiers on too many occasions, been condemned to the worst mise defeat in the very midst of their triumphs and glories. olden times it was "famine and the ague;" at Rangoon, in and the Crimea, it was scorbutic dysentery—a term which to be expunged from the registers of our fleets and armies.

But let us hope that a better order of things is at hand the soldier is about to receive the cares and attentions that a to him, and that harshness may not in future be the reward army surgeon.

"Harshness," says the historian Napier, "is the essence

aristocratic principle of government."

The "Examiner," remarking upon Lord Raglan's censure medical officers, says:—"We should be glad to see some as well as blame, and that the meritorious services of mofficers do not escape notice more than their faults. Was a of honour given to the memory of Dr. Thomson, for a not

the army, proudly answer by pointing to this statue, and by citing the life of th represents—the life of the worthy Larrey.

'I have said it elsewhere, and I will repeat it whenever an opportunity the military surgeon, fearless in epidemics, fearless in the field of battle, p all kinds of courage. He has military courage, because he faces death off all sides by fire and sword; and another courage, far superior to this, preserves his calm coolness and presence of mind when his life is in the greate The blow which is aimed at him, and which he sees threatening, cannot, and even able, would not, be returned by him. He knows his hazardous situation, a not hesitate to fulfil his dangerous duties.

'Kneeling by the side of the disabled, with as firm a hand as when he is so Nature in the anatomical rooms, he dresses their wounds. But with these two courage he reaps two kinds of glory; and Larrey, who has shown his courage equation both, now deserves to be honoured with double glory. He has proved, who wounded, that the dangers which the military surgeon runs, are not imaginar was wounded once in Egypt, in times of glorious memory, and another time at

loo, on that mournful day for France.

You heard, from those who addressed you before me, what the life of Lanbeen, and what services he has rendered to science! It is not for me to enter idetails of this noble career. I am, besides, not prepared for it; I speak, carrie by momentary impulse, and by the admirable speeches which have just been utte judge this learned man,—this defender of mankind,—by considering, as a who useful life, marked by the most enlightened and noble services, and I bow before the which so worthily represents him. Yes, I greet Larrey! the virtuous, man, whose self-denial and devotion triumphed even over the elements, and when among us as an incarnation of genius and humane feelings. He deser thanks of Science, the Army, France, and the whole civilized world!"

What would they have said in France to the memory of "that hero of the mo

Alma, the Surgeon Thomson ?"

ample, who volunteered the charge of 700 wounded Russians on the field of the Alma, assisted only by a single servant as devoted to duty as himself? This hero, in the cause of humanity, died of cholera, immediately referable to his exertions and exposure among putrefying bodies. Had he lived, would he have obtained the praise which is bestowed on a staff officer for galloping with a message on the field of battle, or for helping the Commander-in-Chief in doing nothing in a soldier's battle and victory? The smallest services of the men with names and positions are over-weeningly prized—the noblest services of the unconnected and humble are unnoticed and unhonoured."—Medical Gazette.

Napoleon, on the other hand, in a manner for ever memorable, and while in the agonies of a disease which he knew to be incurable. remembered with kindness the surgeons of the Grand Army. He bequeathed "To Baron Larrey, Surgeon-in-Chief, one hundred thousand francs. He is the most virtuous man I have known." Again, he bequeathed "To Surgeon Emery, one hundred thousand francs." Nor should the conduct of this illustrious patient to Dr. Archibald Arnott, of His Majesty's 20th Regiment, be forgotten. "The Emperor, on his death-bed, desired that a valuable gold snuff-box might be brought to him; and having, with his dying hand, and last effort of departing strength, engraved upon its lid with a pen-knife the letter 'N,' he presented it to his kind and valued friend, as a parting memorial of his deep esteem and heartfelt gratitude." Not satisfied with this mark of personal kindness and regard, the dving Emperor bequeathed to Dr. Arnott a sum of money also.

The treatment of the army surgeon by this incomparable commander and statesman, is not only important and interesting in itself, but also for the memorable reasons which he urged in support of his conduct—reasons which ought to be known to, and

appreciated by, the medical profession.

Napoleon deemed the honours conferred on such men to be called for alike by justice and necessity, in any plan which should pretend to give "consistence to the system of recompenses" granted by the State to its servants. He considered their exclusion from honorary distinctions as founded on the "injustice of feudalism." In the awarding of honorary distinctions, he acted promptly, and upon the standard of true merit:—"Returned to Paris, Napoleon gave the gold cross he had worn through the Russian campaign to his physician, Corvisart." With him the cross of the Legion of Honour was but a symbol which expressed the gratitude of the country—the outward sign by which the State recognized equality of merit.

In the Council of State, held to frame the code Napoleon, it was

proposed by Count Mathieu Dumas, that the orders of the Legi

of Honour should be confined to military men.

"Such ideas," said Napoleon, "might be well adapted to feudal ages, when the chevaliers combated each other man to ma and the bulk of the nation was in a state of slavery; but when military system changed, masses of infantry and phalanxes, co structed after the Macedonian model, were introduced, and after that, it was not individual prowess, but science and skill, whi determined the fate of nations." . . "What is it now which co stitutes a great general? It is not the mere strength of a man of feet high, but the coup-d'œil, the habit of foresight, the power thought and calculation; in a word, pacific qualities, not such we find in a lawyer, but such as are founded on a knowledge human nature, and are suited to the government of armies. T general, who can now achieve great things, is he who is possess of shining civil qualities; it is the perception of the strength his talents which makes the soldiers obey him." . . "We must r reason from ages of barbarity to these times. France consists 30,000,000 of men, united by intelligence, property, and commer-Three or four hundred thousand soldiers are nothing in sucl mass; not only does the general preserve his ascendancy over soldiers, by civil qualities, but when his command ceases, he comes merely a private individual. The soldiers themselves but the children of citizens. The tendency of military men is carry every thing by force; the enlightened civilian, on the oth hand, elevates his views to a perception of the general good. T first would rule only by despotic authority, the last subject ever thing to the test of discussion, truth and reason. I have no he tation, therefore, in saying, that if a preference was to be award to one or the other, it belongs to the civilian. If you divi society into soldiers and citizens, you establish two orders in wh should be one nation. If you confine honours to military me you do what is still worse, for you sink the people into nothing. "Moved by these profound observations, the council agre that the proposed honours should be extended indiscriminately

civil and military distinctions;" \*- while, in our country, such m as the Earl of Chatham, Whitelock, Murray, Keane, England, a Prevost, have been deemed more deserving of honours, than pub benefactors like Pringle, Blane, Robert Jackson, Macgrigor, Bu nett, James Johnson, Wm. Fergusson, Ballingall, Hennen, Guthr and a host of distinguished officers, who have bravely encounter every kind of danger, in all climates and situations, and who ha conferred incalculable benefits on the fleets and armies of Englar

as well as on mankind at large.

After a digression, which I trust may not appear altogether in · Alison's History of Europe.

levant, but rather as excused by the occasion, I return to the concluding subject of this article:

LOCALITIES TO CHOOSE.—Where military circumstances do not imperatively forbid it, an elevated and dry soil should always be selected for bivouacs, camps, and cantonments, on undulating ground or the declivity of a hill, the most healthy sites being such as do not, from the natural fall, or from the quality of the soil, retain moisture.

2. Open downs, so called, are healthy.

3. Such grass-lands as have a sufficient fall, or inclination, are healthy.

4. The elevated banks of rivers that have a sufficient fall either

way.

- 5. Tongues of land, or slender promontories jutting well into
- It is said that wherever pure spring water is procurable, there
  the ground is generally fit for encamping or cantoning troops.

7. When marching: to halt, during wet weather, on ground somewhat elevated and sloping, and sheltered from the wind.

8. In such case to increase the number of fires, and to keep

them up to the starting time.

To make frequent halts, selecting places where the water is good, and to take care that the men do not drink cold water when heated by the march.

10. In marshy localities, to encamp on the highest ground pro-

curable.

LOCALITIES TO AVOID.—All damp ground to be carefully avoided; and even where the surface may appear parched up and destitute of vegetation, if it be moist underneath, as by previous rains, it may be found charged with the worst exhalations.

2. Half-dried beds of rivers, of canals, tanks, or ditches; narrow

gorges.

3. Marshy grounds, and such as are elevated immediately above marshes, and grounds which are exposed to winds and currents passing over marshes.

4. Ground covered with underwood, or the vicinity of such

ground.

5. The low, jungly, or marshy banks of rivers and lakes.

6. The best ground, if long occupied by masses, becomes contaminated: it is therefore an established rule to change the ground of encampment frequently.

Though not in perfect relevancy to the subject of this article, I shall conclude with the following directions for the guidance of medical officers, and of such as may be placed in command of

troops about to embark on board ship, and who should attend the following circumstances:-

1. The condition of the transport vessel—the state of health the crew. When ships of war are used, a frigate to be preferre

to a ship of the line.

2. That all old and foul ballasting be changed, and no fresh green wood be allowed in the ship's hold, or any matter capable ready decomposition.

3. The state of health of the troops to be carefully examined and cases of contagious disease, or of old or extensive ulcers,

be left behind.

4. The crowding of the men to be especially guarded agains and care taken that berths are secured for the sick and married.

The most vigilant care to be taken that the aggregate number men do not exceed the proportion of one man to two tons of mea surement.

5. That the men be not allowed to guit the ship, or to be of deck after sunset, when navigating rivers, or proceeding alon

level marshy sea-coasts.

6. Careful and special examination of the provisions should be made, and it should be seen that the salted meat and biscuit are no old: the same care in regard to the water which, when possible should be taken from springs, and kept in iron tanks. Quick-limshould be provided for purifying the water.

7. For distant expeditions, there should be hospital ships, well provided and cleansed, to allow of the separation of the sick from

the healthy.

8. Cold and damp in the ship's hold are to be guarded against and the pumps to be frequently worked, so as to leave but a few inches in the well.

9. Ventilation and cleanliness to be promoted by every means and in bad weather, when wind-sails cannot be used, portable stoves or fire-pots should be carried between decks.

10. In fair weather, the men should be much on deck, and the

should be made to bathe daily.

11. Moderate exercise and amusement of every kind should be encouraged by every means during a voyage; and an union o discipline, with humanity and indulgence towards the men should

characterize the conduct of all classes of officers.

To conclude: - were it desirable even, which it is not, to dis pense with our Native Indian army, it is financially impossible with the revenues of England and of India united, to maintain in Hindustan an European force adequate to the support of the British rule there.

"Our force," says Lord Metcalfe, "does not operate so much b its actual strength as by the impression which it produces, and that impression is the opinion by which we hold India." If, again, the declaration of Colonel Sutherland be correct, that the groundwork of our power in India consists in our substantial British soldiers, then are they deserving our most serious care, in everything which relates to their moral and material welfare.

Their highest qualities may at any moment be subjected to the severest trials; and it behoves the ruling power to see that these men are at all times and places maintained in the most complete readiness for action.

## ON THE TERM OF EFFICIENT SERVICE IN IN

During a long and varied course of service in India, extended opportunities of observing the effects of clim well on the moral as on the physical nature of Europeans slow blight of the constitutional power"—but especially a the military classes; and the result, so far as I can judge funusually wide range of observation at home as well as in is, that the term of thirty years of actual residence in the the very utmost during which persons under ordinary heals of an average constitution, may be expected to retain their

vigour of thought and action.

Of the precocity described by Doctors Moseley and Ai have never seen any evidence: they would seem to have for every other British youth within the tropics a second Gro Reliqui viri tandem fuere; Grotius vir natus est. (speaks of but one form of precocity in hot countries—that age (vieillesse precoce),—and I fear this last is nearer the tru if the age of forty-eight be the stage of highest intellectual in Europe, that period would appear to be anticipated in telimates; and, after the maximum has been attained the decline of the mental and corporeal powers is both sile rapid. Of hot climates it may be said in an especial degre "We may stop when we are rising, but never when we are downwards."

Certainly, all of my profession, and I think I may say Indian public services generally, who have attained to eminthe East, have done so long *before* the term of thirty years's and few indeed have retained much of it for long *after* that

We are glad to see in general society occasional except this melancholy rule; but how rarely? One may, perchar the veteran of forty years' service, in all the mental fresh youth, and beside him the officer of thirty years' service, so muated and enfeebled in all respects; but the first is the except the last is the rule that must ever govern the mass, or, more perly speaking, the average of Europeans.

In India we are too much attached to the absolute

seniority in military promotion, a rule which does not work well for command. It may serve the interested purposes of individuals, but it ill serves the interests of the State.

The disposition to rise by the rule of the muster-roll, rather than by the exercise of knowledge, diligence, and discernment, is injurious in many ways :- it brings men who never were fit to command into the list of generals; while it causes others, who were once efficient, to linger in the army, and to look to the rewards which seniority may bring them-after their energies have faded. Any system, therefore, which should induce young officers to look to the acquisition of rank rather than rupees, would prove an advantage to themselves and to the service. The Marquis Wellesley said, forty years ago: "The army in India rested too much on the footing of profit, without reference to honours at home;" and this owing to circumstances over which the army had no control: its officers "were not held forward to public view as much as other officers in other parts of the world"-persons whose "services were not more meritorious or important than their own." Rank has been withheld from the officers of the Indian army until too late in life to be of use to the recipient, or to the State. Nor can education and experience united counterbalance the deteriorating influences of age. The officers of the Indian army are, as a body, remarkably well educated. Lord De Ros, in his work for the guidance of young officers, states that the superior education of the officers of the Indian army has become an object of remark during the late campaigns in India, where the talent and services of many of the younger ranks have frequently attracted the notice and praise of the most distinguished authorities.

Here, again, we perceive that it is in "the younger ranks" that the benefits of education produce the most efficient results; while Brigadier Hodgson, of the Bengal army, assures us that the "average age of the Major-Generals of the Indian army is somewhat above sixty-five years, and the average length of service alone of the Lieut.-General is about sixty years. This is altogether a state of affairs most unjust to the Indian army, and politically criminal as regards the great national interests at stake, and which are indeed perilled by a great standing army being kept in so palpably inefficient a state with respect to its senior grades, the members of which are already unfit from physical decrepitude

to exercise energetic command in the field."

It is evident, from all that is here set forth, that a superannuation

list has become a necessity to the Indian army.

The promotion by absolute rule of seniority—so injurious in so many ways—often causes our older Indian officers to be superseded in command; and thus their experience even is made to give way, and is set at nought, under an arrangement grounded on a hard necessity.

FF 2

Nor is this important subject to be regarded as foreign to cal investigation; for the moral qualities, age, and physical bilities of the commander do not involve military and politic siderations and results only, but also the moral and physic fare of the army under his orders—its health, comfort, and effit—everything, in short, depends on the character of the comm

Governments may confer material power on a general, but do not wield moral power, he is as nothing. When Lord Cl read over the list of British generals, with a view to emploin the first American war, he observed:—"I know not what the reading of those names may have upon the enemy, but fown part, I confess it makes me tremble." Seniority had a time done its worst for our commanders.

"The arguments in favour of purchase," says the "Times assume as their basis the pre-existence of a system of ser and derive their force from the necessity of remedying the rable abuses to which the unmitigated system of seniority, have it in the Artillery and Engineer Corps, gives rise. Se has in it a certain degree of fairness, since it is the fate of a noble or plebeian, rich or poor, to grow old, but it buys this advantage at the expense of every other defect to which a crof promotion can be liable.

For the first twenty years of military service, increased as be some indication of increased efficiency, but after that this congruity gradually ceases, till every year that is added age of the officer detracts from his utility, and seniority dehim fit for the highest command just when the inexorable him has stripped him of his last qualification for any. Moseniority, by annexing the highest rewards to the longest strengthens the inducement to continue in the army after strength, and faculties have failed, thus keeping down rising and applying the strongest incentives to old men to unduties which they cannot fulfil.

It is a vivid perception of these enormous evils which induced mankind to betake themselves to the system of pur as involving, with all its absurdity and injustice, a less promischief than seniority; and certainly, if the alternative lies we between the two, we shall not be disposed to doubt the wisd the selection. . . . . We believe that the assumption there is no alternative between purchase and seniority is cious; and that in war with much ease, and in peace without surmountable difficulties, a system of promotion by merit mig devised which would supersede both the one and the other."

In the French army I believe the system of promotion co in giving one-third to seniority, one-third at the recommend of the officer in command, and the remaining third under dire of the Minister-at-War. This system gives satisfaction to all classes; and it gives to the armies of France Commanders-in-Chief of the age of five-and-forty.

The retiring pensions of public servants in India have been

ordered as follows:-

For the Civil Service, Pension on 22 years actual service.

22	22	Military Medical	"	27	"	22	37	"	22
"	"		"	22	"	17	22	"	"
22	99	Clerical	22	22	27	15	77	22	"

In reference to the respective ages on entering the several public services, this scale must be admitted to be fairly fixed: but all the annuity funds, whether civil or military, are founded on the general expectation of a longer course of service than that which earns the government pension; and such expectation is verified in the result—but few indeed being in a position to retire on completion of the government term. It therefore would seem as if some term—intermediate between twenty-five and thirty years' service, had been in contemplation both of the government and of the founders of the annuity funds, in respect of the civil and military servants at least; and such an idea is justified by the actual influence of the climate of India on the European constitution.

This article formed part of a Report on the Medical Topography of Calcutta, presented by me to the government of Bengal, and published by its order at the public cost in Calcutta, in 1839. I am now gratified in the support afforded to the opinions I have so long entertained, by so eminent a public functionary as Mr. Welby Jackson, of the Bengal Civil Service, who, in a Memoir submitted to the Government of India, states as follows:—

"In appointing persons to offices of high trust and responsibility, though no positive and distinct reservation be made as to the time they are to hold the appointments, it is tacitly understood that they are not to occupy them longer than they are able to dis-

charge the duty efficiently.

This is evidently a part of the contract between the State and the nominee; and where public opinion exists, and can make itself heard through the press, or the delegates to a national assembly, the condition is sufficiently enforced. No one, under such circumstances, can long continue to hold an office for which he has become unfit by reason of old age, or other fortuitous circumstance incapacitating him, or impairing in a material degree his ability to perform the duty. In India there is no public voice. The people, as a body, know little or nothing of the administration of the government; they take no part in it, either directly or by deputation. It is therefore incumbent on the government, and

more especially on the Honourable Court of Directors (wh some measure supply the place of the public by acting as a cl on the Executive), to frame and adopt such rules as may pre incapable and inefficient persons from continuing to occ

offices of public trust.

In regard to the chief offices in India, the principle is acknowledged and acted upon. The appointments to the off Governor-general, of Governor, of Member of Council, as as of Brigadier, and General of Division in the army, are made a limited period of five years. There is no rule against a sec appointment, though it is of rare occurrence. Apparently, considered probable that at the age when these dignities are ferred, the efficiency of the nominee will not last much lon and, further, it is perhaps thought advisable to have period opportunities of correcting any error of judgment which may be taken place in the selection.

The principle is applicable to all offices, of every descript high or low, ministerial, judicial, fiscal, or representative; more particularly to offices held by an exclusive service, members of which rise by seniority, and become eligible to motion as they grow older. It is true, that the government w appoints has always the authority to dismiss; but, in practice, has been found insufficient, and hence the custom of appoin for a definite period to the high offices above mentioned. To is one standing reason why the period should be defined; quite certain that, in every instance, official capacity is impart after a certain time of life; the period may vary with individe but an average may be fixed beyond which the services of

The truth and justice of what is here so ably set forth canno gainsaid; yet, happily, the evils of the seniority system do press with such severity on the civil service as might, a priori expected—safety being found in selection for all the higher of throughout the country.

But the unmitigated application of the absolute rule of senio

promotion, staff and regimental, to a scientific corps, like the medical department of the army of Bengal is, as might be anticipated, attended with the worst and most disastrous results of all—the necessary detention of the whole administrative staff till altogether worn out in mind and body; till long after the term of efficient service in India has passed away.

The three senior members of the medical service of Bengal, forming what is called the medical board of that presidency, number between them about 120 years of service, or an average of 42 years each; while the next ten officers on the list, constituting the administrative staff of the service, are not far behind in

this groove of seniority.

To speak of such a system as this is to waste time; for a board so constituted must be superannuated by science as much as by years—the ebb of opinion having everywhere left this pernicious rule of seniority high and dry on the drift of old custom. An institution which is not respected has lost its salt, and in these days it cannot last. A slavish adherence to the principle of seniority is, in truth, only a pretence for avoiding a modern

reality.

Henry Marshall observes that-" Under certain circumstances, competition is as necessary to promote intellectual exertion as it is to effect mechanical production. Where persons are remunerated by time, instead of according to the beneficial results of application, or where there is a uniformity of reward without reference to that which is produced, competition may take a wrong course, and he who labours least thinks he has gained the prize. Establishments in which the enrolment and advancement in rank are chiefly awarded in accordance with length of service, or where the advantages are professedly administered on that principle, involve a spirit of COMMUNISM which is very unfavourable to zealous intellectual industry beyond those routine duties which cannot be evaded; and without the energetic exertions of individuals, we are not warranted in expecting much progress towards the cultivation of science, and the diffusion of information; a result which rarely occurs except from competition, honestly, impartially, and widely administered."

Robert Jackson, speaking on this subject, says, that "If reward or emolument be the portion of a man borne so many years on the muster-roll, rather than of a character distinguished for activity, diligence, and discernment, the purpose of bounty is mistaken."

The sedative effect of a fixed situation, with a fixed salary, is nowhere so thoroughly exemplified as in the Bengal medical service; it surpasses the hop-pillow in its results.

The subject of this article, I repeat, is proper for the consideration of the medical officer, and it has not been sufficiently before authority; but it demands the most serious attention, so lowe may be expected to hold India by what, in Europe, is the power of opinion. There is no country in the world who much depends on the persons of the functionaries, and cospirit which animates them, as in India, and consequently, the right selection of persons is of so much import Asiatics always look to the persons, and never to the depart. The Sepoy, for instance, whose own "idea of loyalty," accepted Lord Metcalfe, "is loyalty to his 'salt,' a sense of what is to the hand that feeds him," knows right well who and who Colonel is—who the General of Division, and who the Comma in-Chief; but he has only a very vague notion of what our dispersion.

Thirlwall observes, that "the character of every people is or less closely connected with that of its land;" and we stake but a limited view of the effects of climate, and especial localities abounding in marsh poison, if we considered on more familiar effects of those agents on the European constituents as the various forms of fever, anæmia, &c. The states air affect the mental energies and moral feelings of many industry to a degree inconceivable to those who have not thus subject to their influences, or who have not watched their effects. Even in Europe, the long application of such cause in Tuscany and Switzerland, affects the mind to very idiocy. I have seen a complete, but temporary prostration of the powers result from a residence in our tiraces and jungly distribution, as in Gondwana and Aracan, but especially after the of such districts.

Finally, and independently of the influences of climate, I be with Mr. Macaulay, that official aptitude is everywhere acquit the expense of the general powers, and that in all countries a limit in years should be ordered, beyond which public s should not extend.

## ON THE MORTALITY AND PHYSICAL MANAGEMENT OF EUROPEAN CHILDREN IN BENGAL.

For children of strumous habit, or for the offspring of parents that are what is termed weak-chested, the climate of Bengal, and of India generally, is favourable. Under ordinary precautions, they may be freely exposed to the open air at all seasons, during the early morning and evening, without risk of being chilled; an inestimable advantage, and one for the absence of which hardly any other circumstance will compensate.

The diseases of childhood run their courses also very mildly in Bengal; and, upon the whole, it cannot be said that under proper management of diet, and with precautions against extremes of temperature, the climate of Bengal is unfavourable to European infant life, up to the age of five or six years, when they begin to exhibit the necessity for change of climate by emaciating, and by

out-growing their strength.

The climate of Italy has in like manner been found beneficial in the instances of weakly children, who in England suffered from swellings of the lymphatic glands, gastric irritation and catarrhs; and there can be no doubt that at the tender age here spoken of, and also at the more advanced age of puberty, the removal for a few winters to the south of Europe would tend powerfully to the prevention of pulmonary diseases in weakly children, and those in whom a hareditary disposition to them is suspected.

whom a hereditary disposition to them is suspected.

Of accurate or statistical record, bearing on this interesting question, we have as yet little in India; but I believe that the results of a close observation would afford corresponding facts to those obtained in England and France, as to the greater mortality among infants under the extremes of temperature—the very cold and the hotter months. The winter and spring seasons have been observed at the Military Asylum, Chelsea, to be very productive of tuberculous disease in the children born in the East and West Indies especially.

Dr. Duncan Stewart, of Calcutta, states, that amongst the European inhabitants of that city, the ratio of mortality of adults is greater than that of infants in the proportion of five to one;

while, with the poor degenerate Portuguese, the mortality of dren exceeds that of adults in the proportion of four to one. late Inspector-General, Burke, states Fort William, Calcutta. "one of the worst, if not the very worst, of the military statis India for children," the average annual mortality per cent. 16:29. Here we perceive the difference in the fate of the I soldier's child, in the Garrison of Fort William, and that of better classes of his countrymen, residing within half-a-m him.

The observations of Dr. Edwards on the differences of contion at different periods of life are of the highest important relating to children; and if more generally understood, and carefully put in practice, they would considerably reduce one greatest sources of mortality. The power of generating heat feeble in the infant system, probably from a smaller consum of oxygen in proportion to the bulk, that it is impossible to guarded in the important article of clothing, during the cold sin Bengal, where scarcely a year passed in which I had not sion to witness the most serious, and sometimes fatal results the exposure of infants, imperfectly clothed, to the piercing wind. It may be concluded that the younger the infant, the necessary does caution become on this head.

But where the opposite system is adopted, or that of hard by means of half-clothing, or the equally unreasonable systemathening by means of a double diet, the results are ever speedily destructive than in Europe, and they are brought by the supervention of gastric fever, with cerebral determinates severe diarrhea, or the more slow process of mesenteric. In respect of the management of infant health, and referr the theories and empiricisms brought to bear on it, I have where observed, that even the fatal results of mismanagement rarely cures the mother of her theory or her quackery—so stronger are ignorance and prejudice than death.

It may be said that, under ordinary care in diet and clot the operation of the teething process proceeds kindly it climate of India; and speaking of my personal experies should say that severe teething irritation is seldom a profection, but that, on the contrary, it generally follows upon viously existing gastric, intestinal, or febrile disorder; and not too much to say that, in eighteen cases out of twenty, last are but the results of mismanagement in diet and cloth errors of ignorance and weakness more common to the civilized than to the most barbarous communities. To read books, and to hear some people talk, one might be led to su that the teething process of infancy is a morbid one from ning to end.

I believe that the seasons in which the greatest mortality occurs amongst European children in Calcutta are—the cold season of December and January—April and May of the hot season. In the former, congestive and catarrhal fevers prevail, and in the latter, remittent fever, occasionally of an ardent character, and diarrhœa. Altogether, the deaths are not numerous as compared to the births; and this circumstance is mainly due to the generally rational plan of rearing the children of the better classes, where the wet-nurses, the aliment, clothing, cleanliness, air, and exercise, are carefully attended to.

I did not see one case of small-pox amongst the children of the better classes of Europeans during my residence in India;—a fact which speaks powerfully in favour of the protecting influence of vaccination; for not a year passed that small-pox did not prevail

in the native portion of the city.

It is stated by Lind that "European women in general enjoy a much better state of health than the men;" but that "child-bearing is peculiarly fatal to them in Calcutta; on which account it would be advisable for them to retire to a more healthy situation at the approach of their delivery. Dr. Clark observes that, on the coast of Coromandel, parturition "is not attended with such great danger as in Bengal; neither is the puerperal fever of such a

putrid nature."

From whatever cause it arise, the very contrary of the above holds at present, parturition being unattended with even so much risk as in Europe. Puerperal fever I never saw in Calcutta; and during sixteen years in which I was familiar with the state of health among the better class of Europeans there, I heard of but one death connected with child-bearing, independently of previous disease; and I remember but very few instances of children being born dead. Some three or four instances of fever or of dysentery, inducing premature delivery and death, I heard of during the time specified; and that was all. I would therefore ascribe the fatality spoken of by Drs. Lind and Clark to the circumstance that, in the olden times, the management of parturition was left to ignorant native nurses; indeed, in Hickey's Journal, published in Calcutta in 1780, the public is frequently warned against permitting what the editor terms, the improper interference of European physicians!

On the subject of rearing children in Bengal I would observe that all experience is against it—that it is altogether a cruel and an impracticable endeavour—and that I have seen several deaths to result from the unnecessary attempt to wage war with the

climate.

The late Mr. Twining states that, after much and careful inquiry, he was not able anywhere to find a sample of the third ge-

neration from unmixed European stock. I believe that so scarcely to be found in any part of India, and least of Bengal Proper:—so much for the question of European coltion, respecting which much has been said and written—re less of the fact that Nature has set her ban—a blighting int—upon it.

Volney, the eloquent historian of Egypt, states "that n the Mamelooks, who were a *Caucasian* race, nor the Turks are Mongolians, unless they married native women, whice Mamelooks never did, could continue their race in Egypt their offspring perishing in the first or second generation."

Without an accurate census, and correct tables of mortalic turns such as the following can be of little use; but I could one better information on the spot:—

Burials in Calcutta of Protestant Christians.

Years.	Adults.	Children under 5 years.	Total.
1820	239	43	282
1821	196	50	246
1822	275	47	322
1823	215	55	270
1824	209	63	272
1825	240	54	294
1826	234	36	270
1827	190	45	235
1828	157	77	234
1829	128	32	160
1830	180	28	208

## PART II,

PRACTICAL OBSERVATIONS

ON THE

NATURE AND TREATMENT

OF

THE DISEASES OF EUROPEANS
ON THEIR BETURN FROM TROPICAL CLIMATES.



In the following attempt to explain the physiological is under which the balance of European health is disturb climates, as well as the counter-effects of the climate of there are certain inevitable repetitions of matters wh already been set forth in the chapter on medical climates many persons will peruse this second part of the present whom the first will not possess the same interest;—and it to repeat than to be misunderstood.



## INTRODUCTORY CHAPTER.

In the foregoing portion of this work, I have endeavoured to explain the origin of those formidable diseases which, under tropical influences, affect the stranger European, and to trace them to the altered physiological actions induced by an unnaturally high range of temperature and other causes which, during a residence generally protracted, disturb the various functions on whose

just balance health must everywhere depend.

I purpose now to examine the converse of all this—the reverse of everything that holds under tropical influences—to describe the flow and ebb of excitement—the inertia, approaching to a collapse of the various functions of the body, which follows as a necessary consequence of the previous tumultuous action—and to describe, both the condition and the treatment of the tropical invalid on his return to his native climate. I design, in short, to trace the peculiar physiological circumstances that have led to the state of ill health common to tropical invalids, and also to explain the real or essential nature of their diseases; and, finally, I shall detail the special conditions to be regarded in their treatment. As in the previous portion of this work, I venture with much diffidence to approach the subject by the high but difficult road of physiology.

The courses of investigation here spoken of are surrounded with the difficulties inherent in everything that relates to climate as a science; but fully impressed with these great disadvantages, I am nevertheless more and more persuaded, by every-day experience, of the necessity of prosecuting such inquiries in the manner here proposed. I believe it to promise a superior usefulness in result, whether we regard the advancement of the preventive or

the curative departments of our profession.

A familiar acquaintance with the most concentrated forms of

disease, as they prevail in tropical climates, assists mater wards a just appreciation of those that lie between, or of degrees of disease, as they prevail in more temperate. Observation of the nature and power of external causes production of disease within the tropics assists us also to more just estimate of the power of other and different stances of climate, whether morbific or sanative, in our country. In other words—the antecedent circumstance known, the consequents become more easy of being app. This counter-experience, acquired in the observation of on their return to Europe from hot climates, ought to be evalue by extending the circle of our acquaintance with the ditions of health resulting from the influences of tropical of

Without a knowledge of the antecedent circumstance under tropical heats, disturbed the functions of the strange pean, as well as a knowledge of the converse present co which, under the cold and damp of Europe, disturb his fi in a different and opposite manner, it is impossible to just or comprehensive view of the history of his health preciate his actual condition, or to regulate his medical ment. "To know the cause of disease is sometimes to to cure, often to be able to prevent it." \* The subject is and of increasing importance also, seeing that the numi annually increasing who, from choice or necessity, betak selves to the intertropical possessions of this empire. The sideration, as well as the great difficulties above alluded to, it a matter of serious regret to me, that in such an under am not aided by the experience and observations of ot more competent authorities.

With the exception of the admirable "Observations" of friend, Dr. James Johnson, the subject as here propose treated is almost, if not altogether, new. It is incumbent therefore, to explain the grounds on which I pretend to un the discussion of it.

During a residence of two-and-twenty years in various India, and in some of its worst climates, I had extended tunities, in peace and in war, during seasons of prosper adversity, in hospital and in private practice, of witnessing fluences that affect the health of both Europeans and nativ many years I was a severe sufferer from fevers, remittent termittent. In Gondwana, in Upper Ava, and in Cal nearly fell under the violence of malignant remittents. seen how the masses and how the individuals were affect they lived and prospered, how they sickened, recovered, c—the habits of life and the causes of death, in short,

<sup>\*</sup> Lectures on the Principles and Practice of Physic. By Thomas Watson,

Asiatic and European races. I have had the further advantage of witnessing the influences that tend to produce like results in the native climate of the European on his return home, and it is this last circumstance that specially encourages me to make the present attempt, trusting that, ere long, I may be followed by other experienced and more competent observers from amongst my brother officers of the medical departments of the Indian army.

In proceeding to treat an European invalid just returned from India, it will be found that a consideration of all the antecedents, within the tropics, is quite as necessary to a just appreciation of his condition, as that of the present influences of an English climate. Both circumstances of climate have in fact so strict a relation, one to the other, that a consideration of either, examined apart, must fail to lead to a thorough understanding of those important agencies and changes which modify and stamp the habit and constitution of the individual, so as to render him a special and peculiar subject of medical care. These truths, which can admit of no denial, will, it is hoped, be received, in excuse of the following details. And let no one consider these subjects tedious, or unimportant, but rather that the manner of their exposition

may be so.\*

It is necessary to hold in recollection, in reference to the subject-matter of the following Essays, that of the sufferers from the diseases treated of, a large proportion always consists of young officers belonging to every branch of the Indian services, civil, military, and naval. They generally present in their persons, on their arrival in Europe, examples of the severer ravages of acute tropical disease, or of those which from their original violence, from their frequent recurrence, or from the consequence of severe treatment, have baffled alike the endeavours at immediate cure, and those intended to promote convalescence by recourse to the many changes of climate attainable by a removal to the Hill Ranges and the Seas of the East. None of the servants of the Government are sent home but such as have resisted all the known means of cure; for none can, under existing regulations, obtain sick leave but through the most strict medical scrutiny, and through a succession of different checks; there must be exhibited in every case a proved and certified necessity for so protracted an absence from duty as is involved in a return to England.

Of the other class of sufferers, or those who have retired from

<sup>\*</sup> The object of these cursory remarks is, to point out the injurious results following upon exposure to extremes of climate. Were it proposed here, to speak of the remedial powers of climate, I should quote largely from the valuable scientific and practical work of Sir James Clark, on "The Sanative Influence of Climate"—a work deserving the careful consideration of all classes of the community.

the public services of India, in middle or more advanced life, may be said generally that, from their longer residence in India and from their comparatively increased years, their ailments ar of a more chronic character. The peculiar circumstances her mentioned render it evident that, in treating persons of either class in England, the duties and cares of the physician are increase and complicated in no ordinary degree.

The return of the tropical sojourner to the land of his fathers strange as it may seem, is not unaccompanied by serious risks to his health, and by many moral considerations of a painful and

distressing character.

Since the day when, as a mere youth, he bade adieu to his sor rowing relatives, quitted the protection of the paternal roof, and his native climate, he has undergone changes in his moral and physical nature but little considered or understood by his kinsmer and countrymen in general, amongst whom he now finds himself in middle, or more advanced life, differing in habits, associations and pursuits, from those around him—his nearest relatives de parted, and he an invalid and a stranger in the land of his birth This is a great and sad revolution in life, whether we regard it in a moral or a physical sense, and the medical officer who has observed its violently acute causes abroad, and its chronic and protracted consequences at home, can bear testimony to severe and complicated sufferings on the part of a highly intellectual portion of our community.

The youth, on his arrival in India, is at once placed in a society wherein the social affections and the domestic habits of our native country are cherished and cultivated with a luxuriance characteristic of the climate, and of which no other country, perhaps, has

offered so distinguished an example.

At a distance of twelve thousand miles from the British shores and separated for more than half our lives from all those who, it our early youth, were most cherished, it is but natural that in India we should cling together the closer that all feel in common—the old Indian and the new comer—the utter uncertainty of again revisiting the homes of our youth. Such is the state of feeling in the East; and throughout the whole period of an Indian's career, whatever his line of service, he lives in a society whose generous kindness in prosperity, and whose soothing attentions in adversity, or in sickness, are unequalled in any other quarter of the world; and it may be said with truth, that no one who deserved its consideration was ever otherwise treated by the community of the Indian presidencies.

If we look to the physical condition of the youth, it is at first under favourable circumstances, no less cheering. All the functions of life receive an impulse from the agency of heat, while the susceptibility to impression is increased in the like proportion

The comforts, and even luxuries, of life, are within his reach if he will be but prudent; for want of any kind in India is unknown, except to the heedless or the profligate. Disease, tropical disease, too, is as yet a stranger to him; and should he be in the army, and that he has not campaigned, or served in unhealthy countries, or during the most unhealthy seasons, the equable determination to the surface of the body, produced by an elevated range of temperature, tends, along with the increased excitability of the nervous and vascular systems, to raise the animal spirits, and impart a feeling of invigoration and health delightful to the senses. But time passes on, and even if, through the contingencies of service in various regions, or through his own imprudence, disease has not ensued, the lengthened application of these operations of climate, which at first appeared beneficial, are sure to impair the health, and that, too often, with a termination so suddenly dangerous or fatal, as at once to shock and terrify all who behold it. Loss of time in the climates of the East, implies loss of opportunity to save life; for an omission or a mistake in the treatment admits of no future correction or remedy. As an example of the amount of disease which may occur in India within a brief space of time, I will quote the report on Ensign F., of the Bengal Army :- " Has been in India only three years and a few months, and has already had four separate attacks of hepatitis, two of dysentery, one of continued fever, and two severe tertian intermittent fevers, besides numerous other illnesses." This officer bore the marks of his stormy career of disease, being greatly shattered; but he was young, and he recovered his health. In tropical climates the attentive observation of the early stages of disease is of the utmost importance, for then we can both prevent and cure with ease; we now can cure by mild remedies, while by such simple and easily applied means of cure, we prevent severe ill consequences. In the East, diseases of every kind, when once established, proceed with a tumultuously rapid course to the destruction of certain organs; and exactly as may be the judgment and power with which remedies are applied, so will the result prove a complete cure, with speedy restoration of vigour, or a partial and imperfect cure, ending in chronic disease.

The distinctive value of the two courses here indicated, is most important to the welfare of fleets and armies; for by early and just measures of cure, the men are restored to service, while, by feeble and inert courses, they are consigned to the hospitals, there

to linger and die, as mentioned, of chronic disease.

It will now be proper to look in some detail to the operation of those causes which tend, by their protracted application, to deteriorate or destroy the best of European constitutions; because a practical acquaintance with these causes, and with their modes of

operation, is absolutely necessary to a right understanding effects. In India, we find that no amount of talent, or of and professional knowledge, can compensate for the ablocal experience in the treatment of its diseases; and it is necessary to say that the same kind of experience is necessary to the just management of their sequelæ in Eur careful consideration of all the influential circumstance and physical, under which the European lived within the ("Whilst he did climate there," as Shakspeare expresses go far, however, to supply the want here mentioned; and the hope of contributing in some degree to this end that I to submit the following observations.

The most immediate of the tropical influences on E constitutions are evidenced in the excited conditions of vous and vascular systems, while the most important araffecting the functions of the skin, liver, lungs, and kidne owing to the extraordinarily high range and rapid change temperature, humidity, and other qualities of the atmospithese important influences are exemplified in extreme degrees.

The transmission of the nervous influence, according hard, is more rapid when the temperature is higher.

During the hot and rainy seasons, which in Bengal exte the beginning of March to the end of October, the determ of the fluids generally is to the surface, accompanied by crease of the nervous excitability and animal heat; by an ration of pulse, and a prodigious increase of the pulmon cutaneous exhalations, the latter amounting, in the rainy to an exhausting discharge, as if the very serum of the blo exuded. At this latter season, through the saturation atmosphere, its vivifying properties are diminished, while cessive decomposition of animal and vegetable matter give air in some of the most unhealthy localities the properti tainted vapour bath. The perspiration is not carried off poration, while the transudation is enormously increased heat accumulates in the body, and the local action of co duces serious consequences, rendering the system suscer the least impression from cold, or malarious influences, strong tendency to congestion in all the abdominal organs.

When the atmosphere is saturated with moisture, there no cooling relief to the body by evaporation. The air, according Professor Smith, of Edinburgh, expanding with the heat, which was remain of the same capacity, they must take in a quantity by weight, though the same by measure, of oxygoupporter of life. This, he adds, is one of the reasons of the tude felt in warm climates: "but, if in addition to the air ratified, it be also still further distended by the vapour of

being mixed with it, it is evident that a certain number of cubic inches by measure, or lungsful, will contain a less weight of oxygen than ever; so little, indeed, that life can hardly be supported, and we need not wonder at persons lying down almost powerless in the hot and damp atmosphere, and gasping for breath." It has been supposed by some observers, indeed, that in hot climates both the volume and the weight of the lungs are diminished, as compared to those properties in Europe.

The Natives of India, of the higher classes, avoid all exertion during the rainy season; while the working classes, at all seasons, are sparing of extra labour, and when compatible with the work in

hand, sitting is ever the posture of the artisan in the East.

The excessive watery discharge from the skin during the rainy season must of necessity have the effect of rendering the venous blood unnaturally dense, and of thus causing the European to be more liable to congestive forms of disease. Dr. C. J. B. Williams refers the disposition to liver complaints, dysentery, and cholera, to the stimulating properties of the blood, deprived, as we have seen, of more than its usual amount of water, and of less of its hydro-carbon.

Such, briefly, are the hot and rainy seasons of tropical climates, and such are some of the reasons for their proverbial unhealthiness. The mind even seems to partake in the general relaxation and lassitude, being unfitted for vigorous or sustained efforts of any kind. During the eight months here under consideration, the sleep is seriously disturbed—a circumstance in itself sufficient to

account for many of the attendant evils.

The "cutaneo-hepatic sympathy" of Dr. James Johnson is well known, and so is the theory which considers that, while respiration is less perfectly carried on, owing to the rarefaction of the air, a vicarious decarbonization of the blood is established by a great increase of the biliary secretion. If, as supposed by Liebig, the function of the bile is "to support respiration and produce animal heat, by presenting carbon and hydrogen in a very subtile form to the oxygen of arterial blood," how greatly must the condition of the vital functions-respiration, the generation of heat, circulation, and secretion—be influenced by the palpably increased secretion of bile in tropical climates, followed as that excited action of the hot and rainy season is by a proportionately diminished action, and consequent tendency to congestive disease in the cold season? Indeed, we are everywhere, in hot climates, thrown back on the observation of that acute and accurate pathological observer, Dr. John Clark, that of all the viscera in the human body the liver is the most subject to disease.

The condition of the atmosphere, during the hot and rainy seasons, produces in reality many of the results of an insufficient

disposition to morbid affections of the chest, I recommend such persons to proceed to India, as offering the best prospect of escape from a condition of disease which, in England, terminates too frequently in hopeless pulmonary consumption. Such, in a few words, are the principles on which I have been in the habit of resting this important question.

During the hot and rainy seasons here referred to, the function of the kidneys is always diminished, while at the same time the

urine is surcharged with saline and other impregnations.

Unnatural and over-excited in all his functions as the European subject has hitherto been viewed, we soon find him, under the influence of the cold seasons, affected in a manner completely the reverse. The blood, owing to the larger proportion of oxygen used in respiration, now becomes more arterialized, while its consistence is rendered more viscid. In the cold season, a larger quantity of oxygen is consumed in maintaining the temperature of the body. The now altered balance of the circulation and neryous function; the entire drying up of the surface, and the corresponding congestion of all the viscera; the consequent diminution of the biliary, and the enormous increase of the renal secretions; the oppressed state of the cerebral, thoracic, and abdominal functions, leading to the most formidable and complicated diseases, and to the suppression of chronic eruptions in a previous state of development; all these circumstances render the cold season of Bengal the reverse of healthy, except to those of sound constitution, or the newly-arrived European, or such as have not suffered from the previous unhealthy seasons.

The fevers of this season, unlike those of the hot weather and rains, approach gradually, insidiously, but not less dangerously, than those more ardent and concentrated types of fever last named. It will be seen that congestion has repeatedly been mentioned as participating in the most dangerous of our diseases. That climate has a principal share in producing this unfavourable state there can be no doubt; but admitting this, we must consider also that the absence of all exhilarating exercise of mind and body, with their animating, varied, and healthful influences, on all the functions, predisposes much, when aided by a too full and stimulating diet, to this end; unhappily, too, the European resident in tropical climates, has no sufficient remedy against the evils of this double inaction, except moderation in diet, which many will not adopt; for during the hot and rainy seasons, the amount of exercise necessary to health in temperate regions would here be impossible, and would be hurtful even if possible. Those who would preserve their health, therefore, must be temperate-use such bodily exercise as each season admits of, and relieve their minds from the monotony of routine official duty by the inexhaustible resources of European literature, science, and elegant accomplishments. is only thus that health, happiness, and reputation can be insu under the accumulated disadvantages of our position in Inc where too commonly the whole time is given up to business.

I shall now enumerate the diseases peculiar to each season, principal that it is to a long continuance of exposure to the varicauses stated, more than to their intensity, that we must refer greater portion of the injurious influences of tropical climate

European constitutions.

It has been seen that in hot climates the lungs are relieved of portion of carbonic acid through the increased activity of cutaneous exhalations, while the kidneys and mucous surfaces alike relieved of uric acid, urea, salts, and water. This state activity, which holds during eight months of every year, will plain how it is that in such climates diseases of the air-passaglungs, and kidneys, are of but rare occurrence, while, on return to Europe, dangerous diseases of these organs are liable to occ

In many persons, on their return home, the disturbed balance function between the skin and kidneys gives much trouble, a requires an attentive medical care; for, owing to the more or partial arrest of the cuticular discharge, during the English win and spring seasons especially, the urea and other salts, carried of part through the skin in India, can only pass off vicariously through the kidneys in our native climate. The processes of accommodat and adjustment, under the cold of Europe, are of much difficult some constitutions, requiring alterative and sudorific medicame aperients, and warm baths, for their eventual accomplishment.

Fever is by far the most common result of the high temperat of our hot season in the East, complicated with disturbance of nervous centres and occasional determination to the liver. I fever is generally of the continued, occasionally of the remitt form, and of an ardent character. It demands experience a care in the treatment. Cholera is most prevalent and fatal in

hot months of April, May, and June.

In the succeeding rainy months, fever assumes a remittent of racter, with diminished vital action generally, and a danger abdominal congestion. In place of the burning dry skin racking headache of the fever of the hot season, we have her moist, cool surface, indicating a want of action in the sudat vessels, an oppressed pulse, and a tendency to the most formida collapse. As the season advances, dysenteries as well as few become more frequent, severe, and complicated, the former important the whole of the abdominal organs.

The diseases peculiar to the cold season are, congestive fer of the continued form; intermittents, with the sequelæ of tumid indurated spleen or liver; insidious, subacute, hepatic inflam tion, terminating rapidly in abscess; dysenteries, frequently complicated with hepatic disease,—all more or less acute, according to individual habit, or length of service in India. Apoplexies of the most fatal form known in India occur at this season, as do the paralytic affections so commonly associated with them. Chronic eruptions, in active development during the two previous seasons, are now suddenly repelled, and are followed by congestion of some of the viscera.

Having exhibited, in the rapid manner here sketched, the diseases to which the European stranger is liable in the East, and the powerfully deteriorating influences to which he is exposed during each year of residence, we shall now suppose him, after a sojourn of from twenty-five to forty years, returned to his native country. Here he enters at once, and without preparation, on a season of eight months' cold and moisture, in exchange for the same period of excessive tropical heat, and excessive moisture conjoined. I am here speaking of the English season, from the beginning of October to the end of May. This is a great revolution in climate, and it at once induces a corresponding physiological change in the system, stamping the individual with a peculiar physical character. The exhalent organs of the skin, which for so long a succession of years had been called into the most violent action, now become sealed, as if with a coating of varnish; from the extreme of transudation he comes at once, under the rigour and vicissitude of an English winter, to constriction and dryness of the entire surface of the body. The result of such changes it is easy to perceive.

From the previous extensive, equable, and sometimes violent determination to the surface of the body, he is suddenly subjected to an opposite action—an extreme dryness of the surface, amounting, occasionally, to a furfuraceous desquamation of the cutiele, and at other times to a miliary eruption, accompanied by determination of the entire mass of the blood inwards; a loaded condition of the great venous trunks of the abdominal, thoracie, and cerebral cavities, with diminished power in the heart and arteries.

This is by far the most common result of exposure to the cold of an European winter and spring; but we occasionally find its opposite—a relaxation of the surface, inducing a copious and clammy sweat on the least exertion, accompanied by extreme languor of the circulation, with cold, damp extremities. This state, which is generally confined to persons of leucophlegmatic habit, is only to be removed by a complete restoration to health.

The nervous function, highly exalted during the earlier period of residence in the East, becomes now proportionately obtunded and depressed. These are great changes, and it is next to impossible that any man shall go through them with impunity; indeed,

More unfounded still is the prevalent notion amongst i rienced and thoughtless lay persons, that on returning to E from hot climates they are to take leave of all their ailments same time that they need take no care whatever as to their This is only another out of many examples of the facili which the mass of men receive mere opinions as established it is but the "It is true, they say it" of the Spaniards, and enough for the unreflecting many. "A moderate acquai with mankind," says Dr. James Johnson, "is sufficient to the truth of the remark, that experience seldom instructs the This is quite as true as the observation of the older doctor same name, the moralist and lexicographer—viz., that the some men who are incapable of acquiring experience. In rare both remarks so just as in all that popularly relates to whether private or public.

A gentleman of the civil service came home after twenty residence in India, on private leave, not being considered si arrived in October, and early in the following month he r journey of four hours on a railway, with the window open lung next the window became that night completely eng and owing to the impoverished condition of the blood, and deficient powers of the system, the substance of the lung

filled with serum. He died in a few days.

Engorgement of the lungs of a fatal character is no infi result of exposure to cold, in feeble and anæmic persons, o arrival from hot climates, during the English winter especial three such instances have lately come under my observation modes of treatment proving vain. "It is really lamentable Dr. James Johnson, "to see men returned from a tropical c walking about the streets of London, or going to places of ment, in the cold raw evenings of winter, while the hacking emaciated figure, and variegated countenance, proclaim a tion of the lungs which ill comports with this exposure to th situdes of a northern climate." Sir James Clark justly obse the natives of tropical climates, and of children in this c that owing to their habit of body being disposed to tube affections—the most nearly allied to tuberculous cachexy nary diseases are prevalent and rapid with them in our "co humid atmosphere." The same may be said with truth of pical invalid in general, but of the anæmic class more esp Catarrhal affections, more or less severe, are almost univers tropical invalids on first coming to England; and many 1 thus affected on their passage through Egypt and the Med nean, weeks before their arrival at home.

A captain of cavalry, while in India, had discharged hepatic abscess through the bowels. He consulted me on a

liarly baneful influence. My experience here in the last twelve years, would lead me to conclude that, if there really be any such immunity from cold during the first year of residence in England, as we hear spoken of so generally in India, it is enjoyed only by the healthy, the youthful, and the robust. Numberless examples have satisfied me as to the truth of this observation. A dry, or even frosty cold is well borne, comparatively, even by the enfeebled tropical invalid; but the damp cold produces sensations of indescribable distress and depression in persons possessed of considerable powers of resistance. Many invalids, again, arriving in England in an enfeebled state, seek what they call "the bracing air" of Brighton and other such places during the winter and spring months, in forgetfulness, or in ignorance, that without a previous restoration to health, this said bracing is impossible of attainment. Many valuable lives are annually sacrificed in this vain endeavour. The truth is, that in general, the cold which a tropical invalid can bear without injury for a year or two after his return to Europe, is not of a low degree; and in extreme cases of disease, the in-door temperature, carefully regulated during the winter and spring seasons, is that alone which is safe to him. He must now be treated with a climate very different from such as may in a year or two, and when health is restored, be suitable to him. Common sense points out that the delicate and the sick must be prepared for invigoration, and not forced into it. We cannot "comfort" such a person "with cold," as the Americans do their English customers with ice.

The truth is that it depends on the state of the health and constitution at the time whether cold is to prove salutary or injurious. Certain moist winter and spring seasons are followed in temperate climates by increase of sickness and mortality in summer. "The proverb, 'a green yule makes a fat churchyard,' says Dr. Farr, probably expresses a scientific fact, notwithstanding the evident fatality of cold; and we may hope that the population will escape some of the epidemics which germinate in mild winters, and burst out suddenly in summer." Referring, however, to the dangers arising from cold, the authority quoted adds that, "as age increases the system rapidly loses the power of producing internal heat; and the resulting fatality by reduction of temperature is doubled every nine years after the age of twenty to forty."

In India all our numerous and excellent appliances are directed towards the counteracting of heat, while here, in England, they ought to be directed, especially in the case of the tropical invalid, so as to defy cold. The applications of the former to health and comfort are well known in the East; but amongst the returned Indians, the necessity for the last is not so generally or so well

understood.

suffered from acute inflammation of the liver, but came ho afterwards in what he deemed perfect health. On getting English Channel he was exposed to a cold easterly wind caused violent congestion of the liver and right lung, from he nearly lost his life.

A friend of mine who had resided twenty-six years in I being asked how he got through the winter and spring set home, replied: "A tight band drawn round the head, and round the belly, with indescribable sensations in the skir whole body, represent my feelings during the cold weather

There is another influence, though seldom of a serious which often proves injurious to the returned Indian; I me exertion of the muscular powers. Men who for years other exercise in India than an evening and morning ride miles, at a gentle pace, will, under the counter-influence make great and prolonged exertions at home in riding aring. The exhaustion of the nervous and muscular eresulting from such imprudence, is very injurious, and weasons do not recover from it for months subsequently. I hapermanent disturbance of the heart's action follow upon su exertion.

An Indian officer of a stout habit, but of previously good ran, he said, for a quarter of a mile only, to overtake a train. From the time of making this exertion he never enj hour's comfort, both the circulation and respiration bei permanently disturbed. His death which took place twafterwards, was one of unusual suffering.

We have seen that in hot climates the lungs are relieved portion of carbonic acid through the increased activity of the neous exhalations, while the kidneys and mucous surfatalike relieved of uric acid, urea, salts, and water. These tions, which hold during eight months, or more, of every Bengal, will explain how it is that, in such climates, distinct the lungs, air-passages, and kidneys, are of but rare occurand how, on returning to Europe when the complete contall the above-named physiological actions is taking place been established, dangerous diseases of the lungs, bronch kidneys, and mucous surfaces generally, are so liable to occur

It is but natural to conclude, also, that both in India Europe material and opposite changes take place in the c constituents of the blood, as well as in those of all the e fluids. I entertain no doubt on this head, but we must animal chemistry for the demonstration of the facts.

If in hot climates the animal heat rises from 2°.7 to 3° that of temperate climates: if for years the functions system are adapted to this increased temperature, what

must be produced when circumstances render the stimulus of this temperature impossible—when the tropical invalid, or the old

Indian, returns to his native climate?

The process of respiration under the tropical rarefaction of the air, and under the condensing cold of an English winter and spring, must also be materially affected, so as to cause corresponding changes in the amount of oxygen absorbed, and in that of carbonic acid expired. The increase of the animal temperature in hot climates is dependent, to a certain extent, on the more rapid circulation and metamorphosis of the blood, for less oxygen is inspired in hot than in temperate regions, owing to the rarefied state of the air in the former.

The relation between the functions of the lungs and liver, unnaturally disturbed in hot climates, becomes again on a sudden disturbed in a direction towards the restoration of the balance of health, on our return to temperate climates. This is a difficult process to the tropical invalid, and one that is attended with risk to both health and life. In addition to these hazardous circumstances, the dangers from excesses in diet, from close and overheated rooms, and from subsequent exposure to night-cold, are not

to be overlooked.

Anæmia is common to fifteen out of twenty of tropical invalids on their return to Europe. In this morbid state, the paucity of blood corpuscles causes a diminished absorption of oxygen, a quicker circulation, and consequently a diminished animal heat. I have no doubt, that by long-continued exposure to tropical heats, and to malaria, the nervous and muscular systems of the heart and uterus are relaxed and weakened, becoming thus irritable, and defective in contractile power. I believe that to these circumstances we must, in a great measure, refer the diminished power of the forces that circulate the blood, and the intermitting character of the pulse so common to old Indians; also the frequency of uterine hæmorrhage in females who have been long in India.

We have now viewed the European from his arrival in India, during his career of active service in the tropics, to the termination of his public life, and his return home. How vast the range of the physical and moral influences to which he has been exposed, and how changed does he find himself on his return, which is, in fact, his first settlement in life; how altered are those of his nearest relatives who remain to greet him, and how many more of

them have been gathered to their fathers!

The depressing influence of the great and long-continued flux and reflux of the functions—of all the causes, in short, just enumerated—will go far to account for the hypochondriasis and other

disorders, not to speak of the graver ills to which our count are prone on their return to Europe. If the returned Ind the father of a family, he has seen his children removed fr care of a mother and his own control, at the tender age years or earlier, their education, moral and intellectual, consigned to relatives in England, perhaps to utter stranger thus we occasionally find that the children of the absent g without that deep instinct of filial reverence, those family tions, and that self-respect, which are so necessary to p youth in the trying passages and ordeals of life. He has aga these children return to India, the land of their birth, at of eighteen, knowing their parents only by name, their affections being fixed on those at home who had the care of the domestic ties having been broken in infancy, never to paired. Truly has the Hindu poet said, "There is no for a soul wounded by the sharp sword of separation." Th have been rendered strangers in the house of their fathdaughters have never known the care of a mother; and it happen, that before either has been allowed time to cultiv acquaintance even of their children, they are again door separate, the father being driven home to repair his shatter stitution, while the sons and married daughters remain in No wonder that, under such unnatural circumstances, the h his youth should be to him cheerless.

It has been seen that all the conditions of life, in the in of the European resident in tropical climates, tend to u depression of the constitutional powers, whether we regainfluences of an unnatural climate, aggravated by an uns regimen, or the moral circumstances. In warm regions, over, there is a precoce development and maturation of the pean constitution, while, unhappily, the tendency to decay proportion, unnaturally and prematurely accelerated.

Curtis says that "persons returning from either India bring home with them a constitution wasted and debilitated diseases of those climates; from which they recover with

culty, or not at all."

Let it not be supposed that I here present an exaggerate of those disorders and diseases, or of the social and moral vantages to which an European is subject, during a resimore or less protracted, in a tropical climate; or yet of smay visit him at home. I do not relate a mere description have personally, largely, and repeatedly shared in the suf I detail—both in India and in this country. Mine is not ture drawn from the imagination.

It must be recollected that I am not speaking of the healt

Indian-a select life in any country-nor of the youthful and healthy officer on furlough; for to both the cold of an English winter may prove tonic and agreeable. Such men may control, through their superior power of resistance and of accommodation, the influences of one law of nature, by the operations of another. These observations refer to such only as have suffered from protracted residence in hot countries, and to those who have laboured under the diseases incident thereto. And here I must bear testimony to the manly fortitude and cheerful resignation of the Indian officer under suffering. We find in him no peevish, fretful, or presumptuous intolerance of sickness. He does not expect an instant cure of all his diseases: and failing of this, he does not flee in ignorant fancy to the empiric. He has generally seen and suffered so much as to render him considerate at once of what is rightly due to himself and to others. He is therefore, and under all circumstances, an object of interest to his physician and surgeon. The highest professional authority in this country has spoken to me emphatically on this remarkable characteristic of the Indian officers, and on this account, as well as on that of their great intelligence, he designated them a "peculiar class."

I know not a better test of good sense, or of a well-regulated mind, than that furnished by our conduct in sickness. Under suffering are exhibited in the most marked manner the results of a sound education, religious, moral, and intellectual. The practised physician detects at once the sensible, firm, and well-trained mind; and with the same speed he discovers the senseless, frivolous, and ill-regulated tendencies of his patient's disposition. I have everywhere observed that the same absence of principle which leads weak minds into extravagancies in religion, renders the same persons also prone to follow quackeries of every other

We come now, in conclusion, to the practical issue as to the medical treatment of the European in the Eastern and Western hemispheres. In the first, we have had to overcome, by the most powerful remedial means known in medicine, the most acute and dangerous forms of disease, and that under the continuous disadvantages of a high range of temperature, and other injurious influences; while, in the latter instance, we have to deal with the sequelæ of these formidable diseases, in a temperate climate, and under the sedative influence of cold.

kind.

It is obvious that under these opposite circumstances of climate and of disease, our means of cure must have corresponding differences. Inasmuch, however, as the injurious influences of tropical climates, in the great majority of instances, have an exact relation to their duration, so also do those favourable operations of temperate climates require time for the demonstration of their three per thousand annually, during the first three years after their return from India! If this be so with the youthful and comparatively healthy soldier, what must be the influence of the variable climate of England on the exhausted and more aged invalid, during what may be termed his probationary residence in it, and how insufficient must be the ordinary sick furlough of two years for the restoration of health in all cases of tropical diseases, and how dangerous must be the cold of our winter and spring to the

broken-down tropical invalid?

British soldiers who have served in India entertain an extreme aversion from home service, especially during the winter and spring seasons; and pensioners who have served in hot climates volunteer cheerfully to settle in the Australian possessions. The very deprivation of the accustomed sensations derivable from heat, will account for many of the discomforts and personal sufferings complained of by every healthy European, during their probationary acclimation on return to their native climate; but the Indian invalid finds in the homeward voyage round the Cape of Good Hope, and nowhere else, that equable temperature, to quote the words of Lord Halifax, between "the climate in which men are roasted and the climate in which they are frozen," which is alike agreeable and necessary to him.

Among the older soldiers who return home, the excess of mortality beyond that of the same class in the United Kingdom of the same age, does not much exceed three or four per thousand annually, notwithstanding their longer service in the East; but they are the survivors of hundreds who have sunk under the same course of service, and must obviously have possessed uncommon vigour of constitution originally, to have stood the trial of tropical

service for the greater part of twenty-one years.

Referring once more to the influence of protracted residence in tropical climates, it has always been remarked in the East Indies that Europeans are frequently carried off in what is there termed —" the one year more"—the last year of residence. It is a peculiarly melancholy event; for the deceased has all but completed his term of years, and all but accomplished the object of his hopes, and of his long-cherished ambition. With a view to insure comfort at home, he remains a year or two longer in India; and, with the goal in sight, he is cut off, and all his fond anticipations end but in death. All statistical evidence goes to prove the increase both of sickness and mortality in relation to length of residence; yet, with these proofs before them, we see in India that every season brings with it instances of Europeans, broken down by long residence, and who fall victims to the unnatural struggle against the fatal "one year more."

The Dutch, who are not given to jesting, appear nevertheless

to have been aware of the uncertainties and dangers of de the departure from the East till "next year;" for in one o older settlements there was found an inscription over the g a Dutchman to the following effect:—

> " Here lies — Minheer, Who was to have gone home next year."

In Mr. Neison's Report on the Bengal Military Fun referring to the mortality of retired officers of the Indian ar given by Mr. Christie, it appears, that as compared to the m of England, that of retired officers is uniformly in excess, age; being for the ages of forty-eight and under sixty, about third more. After the age of sixty, the mortality of both officers and soldiers but little exceeds that which prevails civil life amongst persons of the same age, who have ne this country. But while the mortality amongst retired off proved to be as stated, that of the widows of deceased offic their return to Europe, is found to differ so little from the me of females of like ages in England and Wales, as to be appreciable. This fact can, as I conceive, only be attribu the circumstances, that-firstly, the European female is no jected to the hardships and direct solar and other expos which military officers are liable; and secondly, to the pe temperate habits of ladies in India, which save them from n those formidable diseases which there deteriorate the heal destroy the lives, of their husbands. Thus the widows officers come to England in the enjoyment of health, and th as well and as long as those who have never been absen their native country. \*

I would urge, in conclusion, one other important consider which should be held in recollection, before entering a medical treatment of the retired Indian—viz., that, though in years, or, least of all, in mental character or habit, become prematurely aged in his physical constitution. general principles of medical management must therefore be fully modified and specially adapted to this condition system.

Under no circumstances of disease can such a man be with the vigour that may be safe, or even proper, in perturbed the same age, but who have passed their lives in Er Indeed, no contrasts can be greater than such as are exhibit the moral and physical constitution of the European newly within the tropics, and in the same person on his return native climate after an absence of thirty years.

<sup>\*</sup> For these interesting and important statistical facts in corroboration of I am indebted to Colonel Tulloch and Mr. Neison. They were obtained long article was written.

In the first case we have a youth, buoyant with health and animal spirits, with all his functions exalted, through the stimulus of heat, and with the mind eager for employment and for distinction; while, in the last, we perceive, but too often, premature age, with obtunded sensibilities and depressed functions, coupled with profound mental depression, amounting, in some cases, to the most severe form of hypochondriasis—all resulting from the opposite states of the nervous, vascular, hepatic and other excretory functions, brought about, at first, by an unnaturally high and continued range of temperature, and now, by the sudden application and sedative influence of cold.

It is true that the Indian, coming home at the middle or more advanced stage of life, approaches the land of his fathers with hopes and expectations quite as exalted, though by no means possessing the same glow of enthusiasm and buoyancy, as when in mere youth he quitted the paternal roof:—

"And from the dregs of life hopes to receive What the first sprightly running could not give."

Lord Clive, though but forty-two years of age on his final retirement from India, writes to his friends whom he left in the East:-"I have suffered so much ever since my arrival in England, that I have not been able to interfere so much with public affairs as I could wish. Parliamentary concerns have embroiled me more than is good for my health, and I really begin to grow tired of them." He then exhorts his friends to be satisfied with a moderate competency, and return to England while they have youth to enjoy it, rather than by staying longer lose that youth, and sacrifice that constitution which no riches can possibly compensate for. Clive's disease is spoken of by his biographer, Sir John Malcolm, as "a severe bilious complaint, attended with spasms from gall-stones, loss of appetite, and indigestion; a continuation, or consequence of that derangement of the liver from which he has already suffered so much in Bengal." This great commander, in a letter to his friend George Granville, describes his personal sufferings as "very distressing to the mind and to the body;" and on hearing of Granville's death, he expresses "his indifference, not only to the world's politics, but to the world in general."

To give a more concentrated view of the physical circumstances above related, it may be said, that in hot climates, the air being expanded, less oxygen is taken in at each inspiration. The necessity for hydrocarbonaceous food is therefore lessened. Less of the "elements of respiration" ought to be taken in the food than would be taken in colder climates. In consequence of

the external increase of temperature, less internal heat

quired.

Exercise increases the heat of the body by increasing the of circulation and respiration. In a very hot climate, all in of heat is undesirable. Moreover, the excessive heat remuscular action impossible, because the circulation is directed (in consequence of the activity of the skin) to the skin order that fluid may be furnished for evaporation, to keep the heat of the body, and to prevent the parching of the skinch must otherwise ensue.

In consequence of the lessened muscular action, less albuminous constituents of the food are required to supp waste of the muscles. Hence, in hot climates, less of both of food should be taken, and Nature points this out in the a of appetite; to force an appetite, stimulants are taken, and the system is loaded with nourishment. The excessive pe tion requires an excess of liquids; but, instead of water sugar and spirit, the elements of respiration, are taken we water, in the form of beer, and the spirit, by its stimulating

perties, is doubly injurious.

The excessive flow of blood to the surface—the consequence the high temperature—no doubt prevents, for a time, the resulting from an excess of the two kinds of food. The changes and evaporation going on in the skin draw the circulation to the surface of the body, just as the flame of the lamp drawill up the wick. Whilst the high temperature lasts, this into action of a flow to the surface is kept up. It is probable that is thereby made feeble by the excessive of the skin, as we see it frequently is, temporarily, by the ration bath. As soon as the temperature falls the blood ceflow in excess through the skin. According to the degree of it is almost driven from the surface. It accumulates with congestions and inflammations are produced. Free action inner or outer surface of the body for a time relieves the of tions and enables the circulation to proceed.

After such alternations for years, the resident in the hot returns to a far colder home. There no heat leads the bithe surface; it accumulates in the vessels and capillaries internal viscera. The outward appearance is that of a whilst in reality an internal plethora frequently exists.

The more frequent and severe the temporary congestions viscera have been, whilst in the hot climate, the more per does the internal congestion become when the surface stantly exposed to cold. Though the face and extremit look void of blood, the capillaries of the liver and spleen with the full, and the engorgement of the venous trunks, which v

effect of previous congestion for years, becomes permanent when the coldness of the atmosphere leaves the blood to be circulated by the enfeebled heart alone.

If the nervous influence be the power which regulates the contraction and relaxation of the capillaries, by how much must this important part of the circulation be affected—first, by the long continued heat of the tropics—and secondly, by the cold and damp of the climate of Europe, so as, under both agencies, to disturb alternately, and conversely, this great function?

Tropical heat has had the effect of exalting for a time, and then diminishing the nervous energy, and thus of eventually relaxing and diminishing the circulating power of the capillaries; while cold, on coming to England, will have the contrary effect of constringing the capillaries, and thus of determining the great mass of the circulating fluid upon the internal organs:—such power, acting on the enfeebled European on his return from tropical

climates, must prove of extremely injurious effects.

originate disease."

Dr. Billing concludes his admirable "Principles of Medicine" by stating that they "have been deduced from physiology; and for the explanation of the pathological phenomena he has referred to the action of the capillaries and nerves—nerves and capillaries together, not artificially separated, but as they exist in nature—ramifying and supporting each other throughout; for by their combined action upon the blood sent to them by the heart, they produce the phenomena of health—in their deranged actions they

Dr. James Johnson justly observes that, "The powers of the constitution, however plastic, cannot immediately accommodate themselves to great and sudden changes of climate, even when the translation is from a bad to a good one; and the tropical invalid requires full as much caution and prudence in approaching the shores of England, as he did in landing at the former period on the banks of the Ganges." Speaking of the beneficial influences of the voyage home, round the Cape of Good Hope, the same authority states that, "When the European has become much debilitated by liver affection, dysentery, or fever, and its consequences, the main hope of recovery rests on change of climate, and, under such circumstances, the sea-voyage often effects a cure. Indeed, the instances are not few where more benefit is obtained by the voyage home than by the subsequent residence in England." But this voyage, with its uniformity and salubrity of the sea air, aided by the mental exhilaration of a homeward voyage, producing as it often does, "surprising effects on the animal economy," is not

The overland journey, as it is termed, subjects the invalid to numerous discomforts, both by sea and land—often to insufficient

now nearly so much resorted to as it ought to be.

accommodation, always to a more or less hurried meal—hurried and uncomfortable journey through Egypt; and if, happily, he arrives in England in the winter or spring, he ha encounter their rigours within six weeks, with very imperfec with no preparation, instead of the four months' voyage round Cape of Good Hope, with its rest, ease, and comforts, and with unequalled climate.

The clothing with which invalids are supplied in India is a gether insufficient to protect them against the cold of an east wind in the English Channel; and hence the frequency of c affections, diarrhoea, and abdominal congestions, even before

set foot on shore.

In a moral and physical point of view, and more through agency of moral and physical influences of an extraneous chara than through age, no two persons could present contrasts gre than may every day be recognised in the very individual unour consideration, as compared to the youth of five-and-twyears back.

The average age at which the Indian, whether civil or milit retires, may be taken at fifty to fifty-five years. Many of the who thus return home are, to all public purposes, worn-out mand but few of those even who are equal to the exertion, make effort to enter upon public duties in England—duties for witheir official aptitude, their administrative experience, and to

varied abilities, so eminently qualify many of them.

I would now refer to a question of great importance, and on which I have often had to interfere. Officers of all cla frequently leave India in such a state of ill health and exhaus as to feel hopeless of recovery within the time allowed by reg tion for ordinary leave of absence. In this state of moral physical depression, despairing of health, they often express t purpose to retire from public service and enter into private The advice I have almost invariably given in such cases has I to the following effect:- "You have at least two years' abso leave, at the expiration of which come to me and I will then you what you had better do. At present, borne down by me and bodily suffering, you are altogether unequal to the detern ing of any important question in life, and, for the rest, your is not hopeless." Often have I been warmly thanked in times for this counsel, and that not seldom, before the expira of the two years; many declaring with much feeling that "I saved them from ruin."

"We're not ourselves
When nature opprest commands the mind
To suffer with the body."

However desperate the case may appear, provided ther

system, so as seriously to complicate the case, and add to the difficulties of treatment. These considerations have induced me to select ANÆMIA as the first subject for attention; for the cachectic state now referred to must be very generally kept in view in the treatment of the European invalid, on his return from tropical climates. It has an important bearing on much that relates to the history, nature, and treatment of most of his disorders and diseases. The deprayed state of the blood, in diseases of the spleen, has been noticed in the first part of this work, and there the sub-

ject has received a separate and special consideration.

Anæmia is frequently produced by long residence in the malarious districts and stations of tropical climates, even in persons who have not been sufferers from other actual disease, acute or chronic, such as the fevers and fluxes of their more unhealthy localities. This is what may be termed the cachexia loci; -more properly paludal cachexia. European females of the better class though far less subject to the more formidable of the tropical diseases than males, are yet, through long continued disturbance of the nervous, vascular, and secreting functions, very liable to become anæmic. The excessive perspiration will of itself produce anæmia in weakly constitutions. The very treatment necessary to the cure of acute disease, especially bloodletting, mercury, and drastic purgatives, along with the most rigid abstinence, tends powerfully, even in young persons, to produce a deficiency of red particles; while the same course of treatment, in persons of more advanced age, along with their more protracted exposure to malarious influences, are sure to diminish the old blood-globules, and to prevent the formation of new. The "debility" spoken of by all writers who have noticed the condition of tropical invalids, as one of their main characteristics, will be found, in the great majority of cases, to be none other than the debility of the cachectic habitof anæmia, in fact.

Owing to this impoverished state of the circulating fluid, all the functions of the body are enfeebled—all the secretions defective, diluted, and deprayed; the patient is wanting in the power to repair any serious injury. I often find severe and protracted anæmia to result from the unguarded application of leeches. Three hundred leeches, and in one instance five hundred, were mentioned to me as having been applied to the abdomen in remittent fevers, attended with gastric or hepatic complications, and in

dysenteries

Mr. E—, then a youth of sixteen, fell into the hold of a ship, upon his loins. He writes:—"I was bled three times; hot fomentations were used for ten days, and I had seventeen dozen leeches on (204), and was bled once more."

Lieutenant S—— says:—"I had three different times sixty leeches applied, and I think twenty more afterwards, making in all

ANÆMIA. 475

two hundred in less than ten days." This officer was tall, slender, and of delicate frame, and only nineteen years of age. If, as generally believed, the Indian leech abstracts an ounce of blood—nay, if it be allowed to abstract but half the quantity, how large must be the aggregate amount of depletion in these cases?

Annesley ascertained that the Indian leech abstracts, on the average, one ounce and a quarter, "besides what flows from the bites." This simple fact should prove a warning not to apply leeches in the careless or indiscriminate manner witnessed by me in its results, a few more examples of which I will adduce.

Mr. —, a mercantile gentleman from one of the Indian Presidencies, states, January 3rd, 1855, that after a fall from his horse he had twenty-two dozen of leeches applied at once, by order of Dr. —, while Lieutenant — states, that in the course of two months he had thirty-six dozen of leeches applied to the abdomen,

while suffering from fever.

Lieutenant — was seized in August, 1849, with diarrhæa, followed by intermittent fever and enlargement of both liver and spleen. By order of his regimental surgeon "he was bled to twenty ounces, and about a hundred leeches were applied during the subsequent five days." This officer is described in the Medical Report of his surgeon as "of weak constitution." The patient went to another station, where another surgeon states that he subjected him to "depletion, keeping up a drain from the organ" (the liver), and occasionally a large number of leeches have been applied, as indicated by symptoms. Lieutenant — concludes:

—"I had altogether applied to my side, from December, 1849, to

January, 1851, fully 1,200, at the very least."

But all wonder must cease regarding the leechings of lay persons, when we come to the following case of a medical officer: - "When I went to India, my constitution was excellent, and I was able to withstand any amount of fatigue; but within three months of my arrival, I suffered from an attack of Bengal remittent fever, with great determination to the liver and colon, accompanied by spasms of the stomach and gall-duct; and the treatment of the early stages of the disease having been inactive and inert, it was the opinion of the late Mr. Twining that I would remain a martyr to the ill effects of the climate of Bengal for life. This opinion has been fully verified—disorders of the liver, spleen, mucous digestive surface and bronchial tubes being the result. Early in my illness, organized lymph tubes were voided from the bowels; and this state being regarded as purely inflammatory, leeches were applied in vast numbers. Occasionally fifty leeches were applied in the twenty-four hours; and for weeks in succession the number of leeches expended amounted to upwards of 150 per week. Great exhaustion followed, with a high degree of anasarca of the body and limbs. I came home on sick leave, and I was told by Dr. 97°. The patient was an officer of engineers, a man of high scientific and practical attainments.


How, and to what extent an anæmic condition of the brain, spinal cord, and nervous central ganglia, may affect the functions of the nervous system generally, the present state of our knowledge does not enable us to determine; but that such a condition of the blood must have an extensive influence, no one can doubt. I have seen an instance of what could only be termed nervous

apoplexy from anæmia of the brain.

Neuralgic pains of the abdomen are very commonly to be found on exploration of that cavity; and when they are present in the epigastric, or in either or both the the hypochondriac regions, there is a risk of their deceiving the inexperienced as to their cause: thus leading to very grave errors in treatment. The pains here spoken of, when of a neuralgic character, will always be found to disappear on the return of health; and I would seriously urge the necessity of care in explorations, lest the pains in question be mistaken for those produced by organic disease in either liver or spleen. These abdominal neuralgic pains are common to chlorotic females in all countries, and they seem to depend on a morbid sensibility of the nervous system induced by depravation of the blood, which in its normal condition is the natural support of the nervous system. Here, in Europe, the patient presents the complete reverse of the order of his original acute state of disease under the sun of the tropics-to the acute disturbance of all the functions of life, a torpor and feebleness of corresponding extent have succeeded.

There is now a general cachexia, with atony of all the vital functions. The motions of the heart, along with those of respiration, are feeble and perturbed on the least exertion, and a blowing murmur in the direction of the pulmonary artery, and frequently a loud venous murmur in the neck, can be discovered. Palpitations and throbbings in the head are common, accompanied by giddiness and tinnitus aurium. The voluntary muscles are attenuated, feeble, and relaxed, and there is a disinclination from all exertion.

The excretory functions are imperfectly performed; the bowels are torpid, the tongue being coated and exsanguine. It is evident that here, owing principally to an altered condition of the organic constituents of the blood, a corresponding alteration has taken place in the nutrition of the various tissues of the body. We find, in short, that the functions of the stomach, bowels, kidneys, liver, lungs, organic nerves, and brain, are seriously disordered, through deficiency of blood, or through blood of a depraved quality, or both. I have frequently seen a temporary fatuity to result from anæmia general and cerebral; but it occurs more generally in that form of the disease which follows upon



"The health of different parts of the country differs widely," says Dr. Farr, "and the difference is greatest in summer. In the summer of 1841—50, the mortality in 506 districts, comprising, when the census was taken, 10,126,886 people, was at the rate of 18·15 in a thousand annually; while in 117 districts, comprising the chief towns, and 7,795,882 people, the mortality was at the rate of 25 in a thousand annually. Thus at least 7 in every 25 deaths which occur in towns are the result of artificial causes."

Ever since my return from India, I have remarked that the most rapid, as well as effective restoratives to health, have resulted from a visit to the Highlands of Scotland. Doubtless the moorland exercises of walking and riding, with the cheering accompaniment of sport, and the general prevalence of temperance in diet, have, in this instance, contributed largely to the benefits derivable from inspiring pure air. But the peculiar local properties of the air have appeared to me to aid likewise in the beneficial results;—I mean the union of the ocean air to that of the mountains, such as we everywhere find it in the Highlands. A pure mountain air we appreciate in many countries; but the proximity of the Atlantic and German Oceans to the mountains of Scotland cannot fail to constitute a powerful and peculiar accessory.

Those who have to remark on the difficult and tedious convalescence of invalids in certain unfavourable localities, will require no arguments as to the importance of this subject: "but few have a clear conception," says Liebig, " of the influence of air and temperature on the health of the human frame." I seriously impress on the anæmic patient the necessity of avoiding all occasion for the use of mercury. If benefit does not accrue within a given time, I recommend travel, and it is surprising what an impetus this simple measure gives to the flagging functions, and to the restoration of the blood. With no class of invalids is the medicina mentis of more importance than here—the benefits to be derived from a well-regulated course of "dietetics of the mind" being second only to a proper adjustment of the ordinary medical appliances. "A journey," says Sir James Clark, "may indeed be regarded as a continuous change of climate, as well as of scene, and constitutes a remedy of unequalled power in some of those morbid states of the system in which the mind suffers as well as the body."

With the aids to be derived from a temperate climate, the medical management of these cases is generally easy: that is, when the patient is not advanced in life, and the anæmic condition is not complicated with organic disease. The various salts of iron may almost deserve to be termed specifics: this is so much the case that we can hardly go wrong in prescribing the citrate of iron, the citrate of iron and quinine, the compound iron mixture, the protoxide and sesquioxide of iron, the old wine of iron, with

ANÆMIA. 481

that, in the country, he became so much improved in health that in eight months he was enabled to discontinue all medicine. During eighteen months that I was in occasional communication with this officer, he continued to report favourably of his health and strength; and at the end of two years he returned to India.

Remarks.—This is an example of simple uncomplicated anæmia, resulting from remittent and intermittent fevers, and their necessary treatment by bloodletting, mercury, purgatives, and low diet. The patient had been under able and experienced medical management, but the malarious influences proved too powerful to be resisted even by his robust European constitution. By a careful attention to his habits of life, and to the simple rules of medical treatment mentioned, Capt. T—— left England in perfect health.

Case II.-Lieut. M-, aged twenty-six, of vigorous frame, served nine years in India, and latterly in Sindh, where he contracted severe intermittent fever. He was leeched largely and repeatedly, and took calomel, purgatives, and quinine. After ten months' residence in this unhealthy province he became so enfeebled by repeated accessions of fever, that a return to Europe was at length considered necessary to his recovery. He is now in the pale and bloated state characteristic of the diseased condition of the blood under consideration; being in what may be termed an acute state of anæmia. The pulse is hurried and feeble; the abdomen hard, full, and tense; but I cannot discover, on the most careful exploration, that there exists any enlargement of either liver or spleen. He had two attacks of ague on the way home; one in Egypt, and the other at Southampton. The complexion is very pale and bloodless; the mind feeble and desponding; the appetite is wanting, and the bowels constipated; the entire surface is harsh, dry, and heated. He is a cigar smoker, and I attribute much of the disturbance of the heart's action to this circumstance, having observed the same result in hundreds of instances in India. To regulate the bowels, he was directed to take, every morning, a solution of the sulphate of magnesia and quinine, with dilute sulphuric acid: and as a general tonic he took, twice a day, two ounces of a mixture containing the sulphates of quinine and iron, with dilute sulphuric acid. When, by these means, the tendency to fever had been subdued, the frequency of the circulation moderated, and the bowels brought into regularity, the more simple chalybeates were employed. He was then ordered a light diet, with a little wine or bitter ale, and no cigars. He used warm baths, with powerful friction to the entire surface of the body night and morning. In a few weeks he was able to proceed into the country.

Remarks.-This officer recovered but slowly, being subject to

ing there during twenty-four years, there occurred 63,810 cases of fever; and out of an aggregate British force of 82,342, serving in the Presidency of Madras generally, from 1842 to 1848, there occurred 22,923 cases of fever. Amongst officers and the better classes of Europeans, who, from their better habits of life, are not so subject to hepatitis and dysentery as soldiers, it will very generally be found that fever in some form has been the first disturbing cause of illness, and that to recurrences of fever are to be referred many of their subsequent more local diseases on their return to Europe, whether these last be seated in the liver or in the bowels.

The experience of the West Indies confirms the statistics of fevers in the Eastern hemisphere. Out of that portion of the inhabitants of George Town, Demerara, who are likely to resort to public institutions—the total population of the town amounting only to 20,000—there were admitted into the colonial hospital, from June, 1846, to June, 1847, 2938 cases of remittent and intermittent fever; and the annual consumption of quinine in Demerara and Essiquibo—countries somewhat resembling Lower Bengal in medical topography and climate—averages 3000 ounces!

If we look to the embouchures of the Mississippi, we shall find a state of public health, and a prevalence of fever especially, not less remarkable. The New Orleans Charity Hospital is one of the most extensive fever hospitals in the world. It appears from the records that, in a period of nine years, from the 1st of January, 1841, to the 1st of January, 1850, there were admitted into this hospital 73,216 patients; of which number were admitted, for all the different forms of fever, 33,381; and among these last, for

intermittent fevers, 17,217.

Great as are the advantages possessed, in India, by the officer over the common soldier, as to habits of life, accommodation, &c., it is still matter of every-day observation, that even amongst the better classes of Europeans there, both civil and military, the sequelæ to malarious fevers are both severe and frequent. Some of these I have already noticed; but here I shall confine my observations to the slighter forms of organic diseases, and to the functional disturbances and ill-health which follow on tropical fevers, remittent, intermittent, and continued: those conditions, in fact, which are common to the three types, and which are most frequently observed in England, as the results of tropical fevers. It should be held in recollection, also, in reference to the subject of this article, that most of our tropical remittent fevers are in a great degree, and very generally, of a gastrie character, one of the prominent conditions consisting in acute congestion of the mucous digestive surface; so much so, as to cause Robert Jackson to designate the bilious remittent fevers of the West Indian Islands by the term "gastric, or bilious remittent fever." This profound The obstructed and depraved state of the blood, and consequent depravation of all the secretions and excretions, and of assimilation also, must not be overlooked.\* We sometimes find the heart impaired in its functions, through long-continued and often-repeated engorgements of the great venous trunks during the cold or congestive stages of remittent and intermittent fevers, and from the violence of the subsequent bounding reactions. Through the operation of these causes, associated, as they often are, with an anæmic condition of the system, the heart is occasionally found to be flaccid in its texture, and dilated in its cavities, while its action is feeble and irregular. In every form of malarious fever, but more especially in "the tremendous remittents of hot climates," it is manifest that the sensibility and energy of the entire nervous system are blunted and enfeebled, and that the muscular system, voluntary and involuntary, is relaxed and enfeebled also, in consequence both of the lesion of the nervous system and of the altered condition of the blood, so common to tropical fevers, and as resulting from their treatment.

These disturbances, repeated during years, in a climate moreover which, of itself, and without the intervention of actual disease, disturbs greatly the functions of the nervous, vascular, and muscular systems, will go far, I think, to account for the *intermitting* state of the heart's action, so often noticed in these essays as characteristic of a large proportion of returned Indians. A long residence in the East, without the intervention of fever, may alone induce the irregular state of the pulse now under consideration; but a lengthened stay in India, together with repeated

fevers, is almost certain to cause it.

From many of the symptoms indicative of disturbed function of the heart's action, witnessed in malarious countries, and from the periodical nature of some of them, I have long entertained no doubt that they often have their origin in the efficient cause of fevers, remittent and intermittent. A cause which operates so seriously to the disturbance of the nervous and vascular functions generally, which induces paroxysmal fevers of the most deadly nature, visceral congestions, and neuralgic affections; a cause which produces most serious disturbances of the cerebral and organic nervous functions, may well be believed to aid at least, along with the general enervation and muscular relaxation, in occasioning the feeble and intermitting states of the heart's action, the epigastric pulsation, and the other nervous and vascular disturbances so frequently observable in the persons of returned Indians.

<sup>\*</sup> To such as would inquire more at large into the causes and consequences of tropical fevers, I would recommend the perusal of the admirable articles in Dr. Copland's "Dictionary of Medicine"—a great work that should be placed, by authority, in the hands of all naval and military surgeons.

functions of the spleen, while the general circulation, and the functions of the skin in particular, are oppressed and enfeebled.

All these conditions become aggravated when the sufferer from tropical fever arrives in England in the winter or spring seasons. In other instances we find congestive enlargement, with the commencement of interstitial deposits, more or less chronic, of the liver, spleen, and mesentery, accompanied, as in the former case of simple abdominal congestion, by torpor of the digestive powers, depravation of all the secretions, a feeble and oppressed circulation, and marked derangement of the cuticular functions. In both instances there is great sensitiveness to the impression of cold. The medical management of such cases, whether functionally or structurally affected, is always a matter requiring care, and the cure may occasionally, in persons greatly debilitated by fever, or by long residence in India, be protracted. We have to emulge with one hand, while we impart tone with the other. We have to maintain a gentle but persistent action in all the depurative organs, at the same time that by medicine, proper diet, free exposure to the air, and by exercise, we improve the quality and augment the quantity of the blood.

CASE I .- A gentleman of robust frame and short stature, aged thirty-eight, had resided sixteen years in India as an indigo planter, and during that time he had been continually, and in all seasons, much exposed to direct solar heats, to excessive rains, and to cold. His habits of life had been moderate, and for fourteen years his health had been excellent. Two years ago, while exposed to cold and damp weather, he was seized with severe intermittent fever, the stage of rigor lasting frequently for three hours. Medical aid was obtained from a distance, and but irregularly. Calomel and antimony, followed by brisk purgatives, and by quinine, overcame the immediate violence of the fever, but he has ever since been out of health. The least exposure to the sun, to damp or cold,-circumstances never considered when in health,-now excited him to fever, especially if such exposure occurred about the full and change of the moon. His fever by degrees, and under the influence of quinine, lost the severity of the rigor, until that symptom entirely ceased, and on the accession of the cold season it assumed the continued form. At length he became worn out by repeated irregular attacks of fever. His muscular system, which had been powerful, fell away, while the abdomen became large, full, and tense. In this state he was recommended to return to England, where he arrived in the month of

He now presents the true physiconia, or parabysma, of systematic writers, in the sense of "morbid congestion."

pure chalybeate, and the purgative was directed to be used every other day only. When the weather became warm in May, the convalescence was rapid; and in the following month he proceeded on a tour to the continent. He used the waters of Homburg; but in his own opinion the change of air and the continued exercise were of most effect. He returned to England in Septem-

ber, in perfect health.

Remarks.—This was a case of excessive abdominal congestion resulting from fever, accompanied by anæmia. This state of the blood was doubtless maintained by the stagnation of so large a portion of the circulating fluid in the abdominal venous trunks. The whole machinery of life was clogged. By the removal of intense abdominal congestion the general circulation was liberated, the quality of the blood improved, while the nervous system was thereby invigorated. Perspiration, digestion, and secretion were also set free towards the restoration of the general health. Youth, a robust constitution, and temperate habits of life, favoured the patient and aided his recovery. His treatment was simple but effective. In a case such as this, a persistent course of tonic purgatives is always necessary; and I believe that purgative medicines, when conjoined with tonics and chalybeates, as in the present instance, do not injure the functions of the colon, as they are too apt to do when given singly in large or repeated doses, or where given with or after calomel.

When this gentleman arrived in England, it was supposed by his medical attendant in the country, that effusion had taken place within the abdominal cavity, but this I was unable to verify, owing to the extreme fulness and tension of the entire region. Certainly the case wore a very unfavourable aspect, and dropsy, general or

local, might well have been apprehended.

Case II.—Major ———, of the Indian army, aged forty, had served twenty-two years in India. For the first sixteen years his health had been good; but during the last six years he was employed in field-surveying, the duties of which necessarily exposed him to great variations in temperature and humidity, as well as to the malarious influences. His constitution, originally delicate, appeared indeed to have been improved and invigorated during the first ten years of residence in India. After that his health remained stationary till six years ago, when he was seized, in the month of May, with the ardent remittent fever of the hot dry season. Violent determination to the head, with tumidness and pain of the right hypochondrium, took place immediately, accompanied by delirium. He was freely bled, generally and locally, and took calomel, sudorifics, and purgatives, in large doses. His recovery was rapid, and apparently complete. By the month of August he

gentle action of the bowels was promoted by the pilula aloes et ferri of the Edinburgh Pharmacopæia. Animal food was used but sparingly, while fruits and vegetables were allowed; also weak brandy-and-water to dinner. He took gentle exercise on horseback daily, or whenever the weather admitted; but the cold easterly winds of March he could not face. Such days were passed, huddled up in coats and blankets, by the fireside. Yet, with care and attention, his health amended, though very slowly, so that by the end of June he was able to ride ten miles on end, the appetite and power of digestion having greatly improved. By this time he had gained eleven pounds in weight, while his mind had become cheerful and hopeful. The medical treatment was continued till the end of August, by which time the Major acquired seventeen pounds in weight, along with considerable increase of vigour in all his functions. The bowels now acted without medicine, the biliary secretion being abundant. By the middle of September he proceeded to the Highlands of Scotland, where, with the aid of a pony, he was enabled to ascend the mountains, and enjoy grouse shooting; but he was incapable of any sustained muscular exertion. With little further aid from medicine, Major - continued to regain health and strength during two years that I was in occasional communication with him, and he returned to India at the end of three years " as well as ever he was."

Remarks.—A torpid discharge of all the functions of the body was the character of Major ——'s case on his arrival in England. To the previous excitation, within the tropics, of the various functions, and to the tumult and disturbance of tropical fever, had then succeeded a torpor and impaired tonicity of the circulation, respiration, and of the nervous functions, with interrupted and depraved secretion and excretion. The gastric and intestinal digestion appeared paralyzed, with characteristic dryness of the mucous surfaces, and atony of the muscular layer of the stomach and bowels. He did not appear to labour under any organic disease of the liver or spleen, nor could I discover through exploration, or through the symptoms, that any anatomical changes existed in the

other abdominal viscera.

Major — was so exhausted on his first arrival at home, that, viewing his recovery as hopeless within the time allowed by the regulations of the Indian army, he desired to resign his commission, and retire into private life. I objected to this resolution, urging that his mind was enfeebled, and that he was consequently not in a fitting frame for coming to the determination of that or of any other important step in life. He yielded to my reasoning, and thus a valuable appointment, useful to himself and his family, has been preserved. By similar advice, given in India and at

ates of quinine and iron, so much every morning as should procure two free evacuations from the bowels; while he used two doses in the day of the sulphates of iron and quinine, with dilute sulphuric acid, as a tonic and febrifuge. His diet to be carefully regulated, using animal food to dinner, but sparingly; no wine or fermented liquor allowed. Warm baths were directed to be used twice a week at bedtime.

Feb. 15th.—Great improvement has already taken place; the bowels and kidneys act freely, and he is not so much distressed by

the cold nor by gastric oppression; pulse 94.

26th.—Bowels violently moved in the night, when enormous quantities of frothy, fœtid matter were voided. This discharge has afforded great abdominal relief, and his waist is reduced four inches. Urine copious and less charged, the complexion clearing, while the shiverings have nearly disappeared.

Hitherto the bowels have acted but twice a day, under the influence of the tonic purgative, and the present discharge is viewed as the liberation of pent-up bile, rather than an ordinary

purgative effect. The medicines and diet as before.

28th.—Continues to improve in all respects, and is able to walk a little in the open air on dry days; pulse 78. The appetite is becoming keen, and requires restraint. Urine copious, and nearly natural; bowels freely moved twice a day. Treatment continued.

March 4th.—Has acquired nine pounds in weight, while the tumidness of the abdomen is greatly reduced. Pulse 76; urine copious and natural in appearance. The bowels continue to act

freely twice a day.

12th.—Getting well; complexion nearly natural and the waist of the usual dimensions. He is now able to wear his former clothes, buttoned, with ease. He takes long walks in the open air and increases daily in strength. The tonic aperient to be taken every other day, then every third and fourth day, so as to be gradually discontinued, while a pure chalybeate is substituted for the

quinine and iron mixture.

Remarks.—The case just related affords an example of the acute and immediate consequences of malarious fever. The rapid transition to Europe, by steam, brought the sufferer home with his tropical ills fresh upon him, with the addition that his fever had now assumed a low continued form. "Visceral disease," says Dr. Billing, "converts ague into continued pyrexia;" and in further illustration of this principle, the same authority adduces the hectic fever of pulmonary consumption, and the irritative fever of surgical cases, which latter he considers "a mixture of morbid sensibility and pyrexia from inflammation." It has appeared to me both in India and in England, that quinine, when given in large doses,

recovered rapidly; and from the time when the bowels were made to act freely, his disposition to fever daily subsided. In a month he returned to the country, greatly improved in health and strength, using only the tonic and febrifuge treatment.

Remarks.—In both the cases just described the tendency to fever disappeared in remarkable coincidence with the subsidence of abdominal congestion, through the use of gentle but persistent evacuants, combined with tonics. That a temperate climate and the absence of malaria, may have materially aided in these results, I entertain no doubt, although in the last recorded very obstinate instance, climate alone proved insufficient; and so it proved also in the case of a gentleman from the south of Spain, whom I lately treated, and who was much distressed by ague, until brought under the influence of the emulgent treatment here described. He had taken enormous quantities of quinine without effect.

Every one has seen intermittents, on the other hand, which resisted the influence of ordinary purgatives, and of every antiperiodic remedy, until mercurial purgatives were exhibited; in other words, until the necessary degree of freedom was given to the abdominal circulation, through powerfully emulgent means. Dr. Watson states that in Zealand, the biliary functions suffer so much during the intermittents of the country, that they are called

by the inhabitants gall-fevers.

That the partial stagnation of the venous and arterial circulation in the liver, spleen, and other abdominal viscera, tends, by the detention of the blood, to unfit it for the purposes of the general circulation, and to dispose also to attacks of intermittent fever, would appear highly probable. Dr. Prout goes so far as to declare that the blood in the portal system is deprived of a portion of its vitality. A condition of the blood exists in the cases here under consideration similar to that in varicose veins; and it has been observed by Mr. Bransby Cooper that the discharge of such blood by puncture of the diseased vessels "does not produce any constitutional effect, the blood contained within the varicose veins, being to a certain extent thrown out of the general mass of the circulation, and retained in a half-stagnant state within them."

ring erai, hee. and and 850, ther ring iver, DUIT-But -tilli-0, 15 APER. while Tri to nd m rrival as m m m ge of and nunid d the : the

lings

ct-

ting

dace

ittle

vere

ion.

very

eral

that

his

of a

OTTO-

tained oxalate of lime in greater or less quantity. The appetite is sometimes voracious, at others defective and capricious.

The secondary or chronic diarrhœas of Europeans who have resided long in hot climates are always difficult of cure; and when of long standing, they constitute a class of disease dangerous and

intractable beyond most others.

Whether the disease under consideration presents itself in the form of diarrhœa originally contracted as such in India, or as one of the sequelæ to acute tropical dysentery, as the result of hepatic disease, fevers, cholera, malaria, the abuse of mercury, and purgatives, or from errors in diet and habits of life, we seldom find the disease simple in its pathological nature—that is, confined to the mucous digestive surface alone. On the contrary, a careful exploration of the abdominal regions, coupled with an attentive consideration of all the antecedent and attendant circumstances, will generally show that diarrhea, in a large proportion of instances, is complicated with, if not mainly dependent on, chronic disorders and diseases of the liver. And here it is worthy of remark, that in India diarrhœa is a frequent and immediate evidence of dangerous congestion, or of inflammation of the parenchyma of the liver, and sometimes it results both in India and in Europe from abscess of that organ. The first approach of congestion or of inflammation of the liver suspends its secreting power, and diarrhoa is the result. An entire suspension, or its converse, an excessive flow of the biliary exerction, will generally produce diarrhea.\* That there are in India, on the other hand, diarrheas of a simple and uncomplicated nature, is a fact well known; but they are mostly acute and comparatively easy of cure, the subjects of them recovering on the spot, so that in effect none are sent home but the most severe, protracted, and complicated cases.

Although chronic diarrhea generally follows on previous tropical disease, I have seen it affect persons on their return to England, who, during their residence in India, had never suffered from any of the diseases of that country. The health of such patients, however, had been enfeebled, and they were thus rendered susceptible to the influence of cold or damp, or to what is always more injurious to the returned Indian—to both conjoined. The returning Indian is often seized with diarrhea from exposure to cold on board ship, especially if exposed to the spring easterly winds. Diarrheas, of course, vary in their nature with their causes, and these last con-

<sup>\*</sup> Dr. James Lind, describing hepatitis in Calcutta in 1762, says—"The cure of inflammation of the liver proved uncertain and tedious, as it was frequently followed by colliquative diarrhea, which speedily put an end to the patient's life;" so that, whether hepatic disease may be more immediately or more remotely antecedent, diarrhea is frequently found to be its result.

menced in India by enormous and long-continued "discharges of pure bile," which would appear to have exhausted the functions of the liver: then came the "white flux," with its complete suspension of the nutritive processes; the anamia of all the organs and tissues of the body; the anamic ulcerations (chronic aphthae) of the mucous digestive surface; and all the attendant difficulties and dangers, so as to call into trying exercise the toilsome and difficult requirements of the physician.

As an example of diarrhœa very formidable in its nature, and of

very frequent occurrence, I adduce the following :-

Colonel —, aged sixty-three, served forty-three years in India, where during many years he enjoyed an average share of good health. In 1840 he came home intending to resign the service, being possessed of a competent fortune; but the call made by the Indian government in 1842, on all military officers then on furlough to return to their duty in India in consequence of the disaster at Cabul, induced Col. ——, out of pure military zeal, to proceed to join his corps, despite of a certificate from myself, and from another medical friend, to the effect that he was not fit to serve in India. An accomplished and distinguished officer, he was soon placed in positions of varied and important trust, and he served in the campaigns of the Sutlej in 1845-6, and in that of the Punjab in 1849.

Colonel —— had previously been twice on sick leave to England, and by the recent exposures and exertions he began again to feel the exhausting influences of severe and unremitting duty, as well as those of climate, when, in January, 1850, he was unfortunately induced to accept a mission into a malarious district. Here he was seized with an acute attack of intermittent fever, from which he recovered but slowly, and he arrived in Calcutta from the N. W. provinces in April, 1850, weak and exhausted. The season and the opportunities for departure to England being considered unfavourable, Colonel —— was induced to remain in Bengal, making an occasional trip to sea; but the climate still proved inimical. In August, that being the height of the rainy season, he was seized with a wasting diarrhæa, which harassed him even after his embarkation for England, by the Cape route, in the end of January,

1851.

During a four months' voyage no improvement took place, notwithstanding the most prudent care in diet. He was, on the average, purged eight times every twenty-four hours after leaving Calcutta; and on reaching lat. 30° north, a great aggravation of purging, accompanied by vomiting, took place. The cold to which he had looked as a friendly and bracing agent, "penetrated into skin being now perspirable for the first time since he left India. and much desquamation of the cuticle having taken place under friction with the Indian hair glove, while the pulse, from being

hard and jerking, became soft and natural.

By the 18th of July, when Col. - left London for the country, the secretions had become quite natural in appearance; the appetite, digestive power, and strength having greatly improved. Within a month from his departure he wrote me: "I have not, after the manner of snuff-takers, rapped at the lid of a pill-box since I saw you, and I am almost persuaded I advance in convalescence daily, and that I gain strength, at least I feel that I walk more at ease, as the pilgrim said who boiled his peas." But convalescence did not continue to advance at this rate. On the 4th of November he wrote-" I am tolerably well, but I have not quite attained the point I could wish. There are but few things I can do with impunity. Over exercise prostrates me; a deviation in diet and beverages from strict regimen upsets me; and I cannot venture upon wine out of my own house, and then not daily, so that my customary strong drink is brandy-and-water. Pickles, preserves, and even honey disturb me. I have no pain nor uneasiness of any kind, but just a sense of a delicate state of the stomach and bowels. It is most sensible about the pit of the stomach—that is, the lower orifice of the stomach, on indulging in any kinds of acids. I take only two meals daily-a full though plain breakfast at half-past eight, and a spare dinner at five." Referring to the distressing irritability of temper so common in cases of this nature, Col. - says :- "Whilst I was nervous, wayward, and irritable, it was a great relief to me to sit down and pen a few lines to a friend or relative. Now, however, that I have become composed, and somewhat settled and at ease, it is an effort to me to place myself before a writing-desk, and forego the pleasure of whiling my time with books or wandering about."

In the beginning of December, Col. — came to London on private business, when I was glad to find him of quite his ordinary appearance of condition and of health. The bowels now acted but once in twenty-four hours, and the secretions were healthy. Being desirous to know the state of the urine, I had it again examined by my friend, Dr. ---, who was so kind as to send me the following very satisfactory report:- "I find nothing excepting the excess of urate of ammonia, and possibly the very slightest trace of albumen. But I am not sure of this, there is so little present." Col. returned immediately to the country, where he was directed to use the fluid extract of Bael with a view to give tone to the stomach

and bowels.

On the 26th January, 1852, the Colonel writes :- "Some time ago, owing to imprudent exposure and very cold feet, my bowels

kinds," but change of climate and medical treatment left him much altered for the worse, being greatly attenuated, with considerable ædema of the lower extremities—the discharges from the bowels being copious, frequent, watery, and of a light buff colour." Alteratives and astringents, with opium, having failed of doing any good, Mr. —— was sent to England in the end of 1842. Here he consulted the late Dr. James Johnson; but though receiving benefit from treatment, he has never during more than six years passed a day without astringent and opiate medicine. He was therefore

compelled to resign the service.

In April, 1849, I found Mr. - worn to the bones with diarrhæa, and with an immense enlargement of the liver. Dr. M'Lennan, from whose experienced and able direction the patient had derived much benefit while in India, being now in England, I felt desirous to have the advantage of his concurrence in the treatment by the nitro-muriatic acid bath, by immersion of the whole body. Dr. M'Lennan writes: "When I first saw him the whole appearance was so cachæmic as fully to bear out the description of one of his friends, that he was dropping into his grave; or, to use the very words of another friend, that he seemed like one who had been interred and exhumed again." He was emaciated to a great degree, with cheeks and eyes sunken, voice hollow, and debility so great, that I fully expected from that cause, to find the presence of anarsarca or ascites; -the former only was present to a slight degree. His complexion was of an ashy colour, deeply tinged with green-skin yellow, dry, and harsh-digestion greatly impairedappetite gone. Bowels harassed by incessant purging, which was only kept in check by vegetable and mineral astringents, aided by large doses of opium. The liver extended from the seventh to considerably below the margin of the false ribs, particularly in front.

In this state he commenced the use of the nitro-muriatic-acid bath to the whole body—i.e. to the trunk and extremities; and by a steady perseverance therein for several months, he very greatly improved. The enlargement of the liver steadily and progressively decreased—fulness and weight of the body increased—complexion became clear, and at length even pinkish—skin soft, of natural colour, and perspirable. Subcutaneous veins showed the presence of blood in sufficient quantity. He became cheerful and even active; and in this state of improvement embarked for Malta, whence I have just heard from him to the following effect:—"I have been examined thoroughly by my medical man, and he says all enlargement of the liver has disappeared, and that my secretions are in every respect natural and healthy. I am increased considerably in size, and have got a plump rosy face such as I have not had for these many years past."

sents no appreciable inflammation of any portion of the mucous digestive surfaces; indeed, when diarrhœa has existed for any length of time, we find an opposite state of the general system, and of the bowels, to that of inflammation—an atony of the system, and consequent atony of the digestive functions, and a specially atonic condition of the discharges from the whole of the secreting organs of the abdomen, including those of the mucous membranes. The brain and nervous centres are likewise in a state of complete anæmia. These circumstances have not been sufficiently considered, either in the pathology or in the treatment of the disease.

I have observed, both in India and in England, that a very severe form of chronic diarrhea frequently followed upon intermittent forms of malarious fever, and it was found in the general hospital of Calcutta, that amongst the European soldiers who had served in China, the diarrhea, which was then always a sequel to the agues of the country, proved very generally fatal. Out of one detachment of twenty-eight cases of chronic diarrhea admitted into the above-mentioned hospital from China, and of whose cases I have received reports, twenty-five men were noted as having previously suffered from intermittent fever, and eleven men were noted as suffering from pain in the epigastric region, and of the liver. These poor soldiers were in the last stages of chronic diarrhœa of the worst kind, and but few of them could have survived to reach their native country, fourteen out of the twenty-eight admitted into the Calcutta General Hospital having died there; in truth, their complicated state of disease left little or no chance of recovery.

The following post-mortem appearances are recorded in the hospital reports: - Excessive emaciation of the whole body; anasarca more or less general; serous effusion occasionally found in the cavities of the pleura and pericardium; a flaccid, flabby state of the heart; omentum "perfectly transparent, without a particle of fat;" in other cases, "omentum absorbed and shrivelled up;" pylorus and commencement of the duodenum abraded and granular; "pancreas hard and gritty;" the liver large, soft, and flabby, sometimes blue, together with occasional enlargement of the spleen, this organ being "softened like grumous blood," while in other cases it was "small and indurated." A pale, attenuated, "diaphanous" condition of the small intestines was generally observed; cæcum frequently ulcerated and thickened, exhibiting marks of former disease; colon sometimes attenuated, but more generally thickened, occasionally "cartilaginous," its mucous membrane being often found ulcerated in many places; mesenteric glands generally much enlarged, mesentery sometimes containing

calcareous deposits "as large as peas."

racter. It is, in fact, in many instances, but the last inevitable stage, or symptom, in the progress of complicated tropical disease.

Anæmia in its most aggravated form exists as the result, but certainly as the concomitant also of nearly all chronic diarrheas of tropical invalids; and this attenuated and depraved state of the blood, and of all the secretions derived from it, tends greatly to keep up diarrhœa: the serum of the blood appears to transude through the intestine. The diminution of the general mass of the blood, and its depravation in quality, must also produce an anæmic condition of the mucous membrane of the bowels, with occasional gelatinous softening of its texture, either or both of which conditions offer serious impediments to our curative endeavours,-impediments often increased by hepatic and other complications. In the instance of an officer of the Indian army, who had been brought to the verge of the grave by diarrhea of long standing, recovery appeared to be materially promoted by his living in the open air during summer and autumn. Could we in these cases impart oxygen to the blood, and thus impart to that fluid more life and consistency, we should, I think, assist the cure in a very material degree; and I make it an invariable rule with all such patients that they occupy a chair or couch, during temperate weather, in the open air, taking care that they are well-clothed at the same time. Where, as here, the blood-forming processes are suspended, no means should be neglected which can tend to vivify the circulating fluid.

In considering the best means of cure, it is absolutely necessary-first, to ascertain minutely the previous states of health and the habits of the patient; and, secondly, whether the diarrhoa be simple in its nature, or associated with hepatic or splenic disease. Along with these considerations, a careful examination of the matters voided is always requisite. Much will depend on the judgment and care given to these preliminaries, for much of the perplexity, differences of opinion and of practice, result from inattention to them. But of all the difficulties encountered in the treatment of chronic diarrhea, one of the greatest is that which relates to the due regulation of the diet, especially when the patient is of the male sex. In this most simple and all-important matter it is truly astonishing how unreasonable and unreasoning we often find the most able and excellent men. With females I have rarely known much difficulty, on the contrary; while with men I have often noticed a morbid craving amounting to fatal infatuation, requiring the active and vigilant guidance of others to counteract; for here the mind is greatly enfeebled, the power of self-control lost. In these unhappy instances men may in truth be said to "pepper themselves into appetite, that they may eat and drink and die." The climax of devotion to the luxuries of

cessarily called to the treatment when the diarrhea has existed long and made great progress; and even here how often do we see patients sliding into their graves through their own negligence, or that of relatives, whom we feel assured that we might save if they could but be placed within the systematic control of an

hospital?

In matters that relate to the conduct of the patient the circumstance next in importance to rigid care in diet is attention to the clothing; and here his sensations and wants will generally be found on the side of safety. The sufferer should be incased in flannel in all weathers, sometimes with the addition of a silk covering, and a double layer of flannel should invariably cover the abdomen, the apartments being maintained of a high and equable warmth, especially in the winter season. In every form and variety of chronic diarrhæa, warm baths, used occasionally at bed-time, will be found of much service; indeed, every means which determine to the surface of the body is most efficacious. In the instance of two officers under my care, who contracted diarrhæa in England, and who were forced to return to India when but half-cured, I predicted that the equable determination to the skin, resulting from external heat in India, would restore their

healths, and so it proved in both cases.

In determining the strictly medical treatment of chronic diarrhæa we ought to be guided by our knowledge of its nature and cause, for without this our measures are apt to become variable and uncertain in their aim and result. For practical purposes we shall do well to ascertain, by the most careful and minute examination, whether the disease be confined to the mucous membrane of the bowel, or whether, with or without anatomical changes in the intestine, there be disordered function, or actual disease, in the liver. This is indeed a matter of the first importance, as regards the selection of remedies, and the safety of the patient. Where the case is simple and uncomplicated in its nature—that is, confined to the mucous surfaces, we shall find it advantageous to begin our treatment by means calculated to allay general irritability and irritation of the mucous membranes. For these purposes the combination of mild opiates with alkalies will be found both grateful and useful. I sometimes found the liquor taraxaci. with soda or potash, and a sufficiency of opium to restrain overaction of the bowels, very beneficial; while, in other instances, the cretaceous preparations, combined with sudorifics, and with opium in larger proportions, and with aromatics, succeed better. In this, as in other morbid affections where acidity and rancidity of the stomach and bowels are so liable to exist together, it is remarkable how often we find the mineral acids fulfil all the indications ordinarily accomplished by alkalies. So much is this the

and diarrhœa. I have frequently seen it arrest ouress of these diseases in twenty-four hours, after all other at treatment had failed. On what the curative property of depends I know not; it is certainly not astringent to the or, at all events, very slightly so. I am inclined to believe much of its efficacy may reside in the thick mucilage which ands the seeds of the fruit. A singular property of the fruit that it does not merely restrain undue action of the bowels, diarrhæa and dysentery, but also in cases of obstinate habiconstipation acts as a mild and certain laxative. It may be in all cases to regulate the bowels." On the other hand, anbrother officer and friend, who went from Calcutta to Egypt, ring from chronic dysentery, writes me :- "The figs and pes of Egypt have done more for me than ever the Bael fruit Bengal;" but here, doubtless, the climate of Egypt proved a verful agent in aid of the fruits, which are everywhere conredly beneficial in such cases.

am indebted to Mr. Henry Pollock for the following account his examination of the preserved ripe fruit, as imported in its led and preserved forms from Calcutta, by several of our best bemists:—"The pulp and the dried shell of the fruit do not appear to me to differ chemically in any respect, except as to mantity. They both contain—I. tannic acid; 2. a concrete seential oil; 3. a bitter principle, which is not precipitated by tribasic acetate of lead, and a vegetable acid. The pulp, as I received it, also contained a considerable quantity of sugar, in which it was preserved. All three of the substances I have mentioned exist in the largest quantity in the dry rind. There is most

acid in the pulp."

Mr. Pound, of Oxford Street, states that in the fruit, as imported by him, the mucilage exists in about the proportion of

twenty per cent.

The preceding observations contain all that appears to the writer to be necessary towards establishing the principles of treatment in the chronic diarrhoa of tropical invalids. But as each individual case will present its special peculiarities, so it will require also special modifications of remedies in its treatment. I proceed now to another branch of this interesting subject.

The following cases are here presented with a view to illustrate

the pathology of chronic diarrhœa:-

Case I.—Mrs. ——, the subject of the following case, was seen by me about six weeks previously to her death. She was twenty-nine years of age, and had resided seven years in India. Very soon after her arrival there, she was seized with intermittent fever, which disordered her health, disturbing the functions of the

with ulcerations on their mucous coats. The gall-bladder was empty; the mesenteric glands free from tubercle or other disease."

Remarks.—The conduct of this lady, under great and protracted suffering, excited the anxious interest of every one who approached her. She expressed "a wish that her remains should be examined," it is believed from a desire that her husband in India might know the actual cause of her death. The case itself is remarkable, as illustrating the pathology of chronic diarrhea, in which, I believe, chronic abscess of the liver to be a not infrequent cause of death. Here, I venture to conclude, that the hepatic abscess was of about two years' existence; in other words, that it had been formed in India, and had existed during twenty months of residence in England. The anæmic condition was adverse to all progress in the abscess, and the "firm sac" was all that nature was capable of performing in its conservative endeavour to bring about absorption.

CASE II.—The subject of the following notice survived but nine days from his landing in England, and his case, like the preceding, is placed in the front rank, in order to exhibit the pathology of the disease under consideration. - , Esq., aged fortysix, resided twenty-eight years in India, eighteen of which were passed at sea, and the rest in Calcutta, in mercantile business. He states that in 1826 he suffered from a severe dysentery of Bengal, from which his convalescence was but slow, the bowels being irregular, sometimes constipated, and at other times loose, for a long time afterwards; but looseness ultimately became the dominant condition. Twelve years ago he had a severe attack of diarrhœa, and four years ago he suffered from a still more severe attack, after which he made a voyage to the Isle of France for the recovery of his health. For two years after his return to Bengal from this last-mentioned voyage, his health remained comparatively good; but in July, 1850, he was seized with the remittent fever of Bengal, since which his general health gradually declined, while the bowels became much relaxed, accompanied by extreme prostration of strength and anæmic giddiness. He struggled on for months, in the vain hope that a once powerful constitution would serve him; but tropical disease pays no respect to European vigour of body, and he was in the end constrained to come home.

He is now (August 23rd, 1851) in the last stage of emaciation and exhaustion from chronic diarrhea. The entire skin is of a dirty mahogany colour, dry, and cold; the tongue is pale and bloodless, with ulcerated edges; the pulse extremely feeble and slow. The erect posture occasions faintness, and for months he

panied him during a long sea voyage, and on her return to India she was, for the first time, seized with fever, which, however, lasted but a few days. After this she enjoyed very good health in all respects until July, 1848, when she contracted tertian intermittent fever, which, in the September following, was succeeded by what was considered ervsipelas of the throat, face, and scalp. During eight days in which this disease continued, she writes that "one thousand leeches were applied, a lady at the station having taken the trouble of counting them." It resulted from all this that Mrs. suffered from the usual consequences of large depletion, including violent palpitations of the heart, and she recovered but slowly, and with difficulty, a sufficiency of strength to enable her to undertake a journey to Simlah, in the Himalayah mountains; and here she arrived for the benefit of climate and change of air, in February, 1849. The anticipations of an improved state of health from the mountain air were not destined to be accomplished; for ague recurred severely, accompanied by diarrhoa, of which she had had a slight attack along with the original intermittent, and during a heavy fall of snow which took place soon after her arrival, the diarrhœa assumed a dysenteric character, and it has ever since become aggravated under the influence of cold weather.

During the homeward voyage her disease did not abate; for she says that, besides much suffering from continued sea-sickness, she was purged, on the average, from six to eight times in the twenty-four hours, in a passage of four months, the matter ejected being fluid, copious, frothy, and generally devoid of colour. Here, for the first time during her protracted illness, she ceased to menstruate.\*

<sup>\*</sup> The frequency with which women cease to menstruate during the voyage to India, and back to Europe, is a remarkable circumstance, and one that continually pre-

aphthæ of the mouth and throat, vomitings, occasional pain in the regions of the liver and of the cæcum, and excessive diarrhæa, occurred twice in every month, and latterly more frequently even.

Thus she became gradually wasted and exhausted.

On these distressing occasions, one circumstance always excited the attention of the patient; viz., that after many hours of retching and vomiting, if even the most minute streak of bile was perceptible on the side of the basin, an immediate remission of fever, vomiting, and diarrhœa took place, followed by a few days of comparative ease. But no real amendment resulted from any plan of treatment. Everything failed in succession, and at length she sank, emaciated and worn out. The following report of the postmortem appearances was recorded by my friend, Mr. Pollock, of

St. George's Hospital :-

"The body was extremely emaciated. Abdomen: The liver was not much enlarged; its colour was very different from the natural healthy condition, and instead of the dark red appearance, it was of a yellowish red colour; the peritonwal coat peeled off its surface most readily, and the structure was soft, fatty, and easily lacerated, as if rotten. The vessels were not unusually filled. The gall bladder was not distended, but was filled with bile; the peritonæal membrane covering it was much thickened, and of a pearly whiteness, and the sensation given to the finger when the bladder was pressed upon was that of a thick india-rubber ball, elastic, as if its contents were not capable of escaping. On cutting into it, some dark fluid bile escaped, also a large number of gall-stones about the size of the common pea, and granular on their surfaces. Under the blow-pipe they were entirely consumed. On examining the cystic duct, it was found contracted and thickened externally, and at the point of junction with the hepatic duct, where the two unite to form the common duct, one of these calculi were so firmly impacted, that it appeared as if very little bile would pass into the intestine. The stomach was contracted, contained very little fluid, and was apparently healthy. The small intestines were also rather empty; their muscular walls were attenuated. The lower portion of the ileum was laid open; the mucous membrane was thickened, but the glands were not affected. A darkish-red appearance presented itself in the inner surface, which was partly occasioned by the vascular condition of the lining membrane, and partly by its being covered with tenacious mucus tinged with blood. The excum and ascending portion of the colon were much distended with most offensive and acrid flatus, and also contained some dark fluid fæces; small isolated patches of congestion were observed on the mucous membrane, but no vestige of ulceration. The transverse and descending colon was contracted, and the small quantity of solid fæces contained in the bowel was of a light cured by the nitro-muriatic acid bath. Such are some of the exceeding difficulties in treating so complicated a disease as that under consideration.

Case IV.—The result of the following post-mortem examination, by Mr. Pollock, was exhibited in the case of the wife of a surgeon of the Indian army. As in the majority of the other instances, this lady's first illness was intermittent fever; after which, hepatic enlargement, distressing and irregular feverish attacks, general anæmia, aphthæ, and excessive wasting of the entire frame, at

length wore her out.

Mrs. -, aged thirty-two. Examination of the body forty-eight hours after death: The body was extremely emaciated, and decomposition rapidly advancing.-Thorax: The cavities of the pleuræ contained a small quantity of fluid. The apex of the left lung was adherent to the walls of the thorax, and in its substance were a few crude tubercles. The remainder of the lungs was healthy. The cavity of the pericardium contained a small quantity of fluid. The heart was pale and flabby, but healthy.-Abdomen: A small quantity of turgid fluid was contained in the cavity of the peritonæum, especially observed in the pelvis. The liver was flattened and spread over a large surface. It was much less red than natural, and mottled on its surface by numerous pale spots and marks. On cutting across these marks, they were found to consist of deposits of lymph and pus, some of them being situated in deeper portions, but for the most part confined to the surface immediately under the peritonæal covering. The gall-bladder was distended with dark thick bile. Some old and very firm adhesions existed between the surface of the liver and the walls of the abdomen on the right side. The stomach was contracted and healthy. The spleen was dark-coloured, and its surface and its substance studded with deposits of lymph and pus, in size varying from a pea to a pin's head. The surface of the lower portion of the small intestine showed increased vascularity of the peritonæal coat in patches, and also was marked by dark-blue spots every here and there, as if from some mischief to the interior of the bowel. On laying open the small intestine, these discoloured spots were found to correspond to ulcerations of the mucous membrane, some being the size of a sixpence, others much smaller: some healing, others quite cicatrized; while some had nearly perforated the peritonæum. The ulcers were all smooth on the surface, and the edges not raised, and confined to the lower portion of the small intestine. The large intestine was healthy.

Case V.— , a surgeon of eminence in the Indian army, aged fifty-eight, had served thirty-eight years in India, during

and the t illness. ch, being 4 he was soon folble him. sars. Im owed by blisters cury and vsenterac he lamer g at that here was neart and elaxarion n by Dr.

return to escribed: e medical 851. Fir egular == le resided England. rrbœa === us. There lower exnight and liver was ions being hough mos there were inine. time the with na everish wentyis time the oc d not and money the bo iscond ppeared ice of

Still no disposition to an improved nutrition manifested itself, and in the middle of October he contracted a severe bronchial cold, followed by copious viscid expectoration, some slight oppression, and gradual increase of debility. In the first week of November, there was a slight recurrence of diarrhœa, and on the 5th of that month he sank from utter exhaustion.

The following post-mortem appearances are recorded by Mr. Pollock:—"The body was extremely pale, exsanguine and emaciated.—Thorax: Each of the cavities of the pleuræ contained rather more than a pint of serum. The structure of the lungs was perfectly healthy, and crepitant, except that they were slightly emphysematous at their margins. The pericardium contained a small quantity of serum. The heart was rather large, and its surface somewhat covered with fat; the latter structure was infiltrated with serum to a very great extent. The muscular structure of the heart was pale-coloured (exsanguine) but firm. The right ventricle was somewhat dilated. The left ventricle was much hypertrophied; its lining membrane was somewhat thickened and whitish; the aortic valves were thickened, opaque, and contained spots of atheroma; their edges were slightly contracted. The root of the aorta was roughened by deposit of atheroma, as also other parts of that artery to its bifurcation. The blood was extremely thin, and perfeetly fluid, no trace of coagula or stain being perceived in the heart's cavities .- Abdomen: The liver was rather large, of a healthy colour, and smooth on its surface, its structure presenting the natural conditions. The gall-bladder was much distended; its coats were very thin; its contents were simply serum, with two or three small particles of inspissated bile. The spleen was healthy, The stomach was natural. The coats of the intestines were extremely thin, and their contents fluid, and like yeast in consistence and colour. There were no indications of recent ulceration of the large or small intestines; the mucous membrane of the cæcum was rather more congested than in health, and more so than the rest of the large intestine. The kidneys were very exsanguine, but not unhealthy in structure.

Remarks.—A long succession of severe tropical diseases—anæmia of several years' duration, chronic anæmic diarrhæa, defect of nutrition—inanition, in fact, destroyed this excellent officer. Could circumstances have admitted of his remaining at home in 1849, so as not to return any more to India, a valuable life might have been saved to his family; but two additional years of tropical service, in his then miserable state of health, left no hope on his last return to England, either from his native climate

or from any other remedial means.

General Remarks.—The post-mortem appearance above recorded, while they exhibit the irremediable lesions of structure the conditions most certainly conducive to the destruction of an organ; for if the mere retardation or detention of the blood in the larger venous trunks be sufficient to spoil the circulating fluid, and render it unsuitable for admission into the general mass, and a source of its contamination, what must be the consequences, both general and local, of a protracted state of congestion of so large and of such important organs as the liver and spleen?

Postscript.—After this essay was written, the following confirmation of many of the facts and views here contained was received

from an unexpected quarter :-

Dr. Stillman, of New York, writing of the diseases of California, says of the chronic diarrhea of that region:—"Many cases of Californian diarrhea were preceded by intermittents; and although many do not appear to have suffered in that way, still it must be admitted that all were exposed to the intermittent malaria. The discharges" (he says) "were copious, liquid, and voided without pain, or but momentary griping. The appetite was so insatiable that no opportunity of devouring whatever food could be obtained was neglected. By nothing was this affection more distinguished than by its fatality. Of twenty-five cases of chronic diarrhea fourteen died before the 1st of April (in three months), and most of those remaining are known to have died since. Before death, emaciation and anæmia became extreme, and the mind, which in most cases had been cheerful and hopeful, at last gave way to the most childish imbecility."

Dr. Schulhof, writing of the diseases of the Delta of the Danube, says justly:—" Where there is ague, dysentery is not far off, the former predisposing for the latter by the disorders which it produces in the abdominal organs. . . . That ague on the one hand, and dysentery on the other, will leave behind them tokens of their visits by liver affections, every one will readily believe." That ague predisposes to diarrhoa even more imme-

diately, would seem to be established.

## CHRONIC DYSENTERY.

CURTIS, writing on the Diseases of India in 1782-3, says:—
"The stomach and intestines, the liver, and mesentery, are here
the grand sources of almost every disease; or, what amounts
nearly to the same thing, in every standing complaint, if the
radical affection does not lie in these organs, they are sure to become the principal sufferers in the issue." The justice of this
observation is confirmed by modern statistics; for we find that
there occurred, according to Colonel Tulloch, out of an aggregate

nature of chronic dysentery, as contra-distinguished from chronic diarrhoa, which latter is proved to be the sequela of fevers more frequently than of any other disease. As seen in Europe, chronic dysentery has ceased to possess any inflammatory character, even of the most chronic nature.

It may sometimes be difficult to assign the reason, but chronic dysentery, whether complicated or otherwise, has always appeared to me to be a more manageable disease in this country than chronic diarrhœa. I find that the process of healing an ulcerated bowel in chronic dysentery is much more readily brought about than is the restoration of the hepatic function in cases of chronic diarrhœa. The history of this latter disease given in the last article will in some measure explain the different degrees of difficulty and danger to be apprehended in the treatment of the two diseases. I have certainly not seen chronic dysentery the sequela to the acute forms of the disease, so frequently associated with incurable diseases of the liver as is the chronic diarrhœa, notwithstanding the great frequency of the hepatic complication in the former disease. But though I state this fact as the result of my personal experience of the chronic dysentery as it appears amongst the returned Indians, I am very far from desiring to underrate its occasional dangers here in England.

In order to exhibit the relations of the dysentery of India to diseases of the liver, I need only quote the following important facts from the statistics of Dr. Macpherson, of the General Hospital, Calcutta:—

		ACUTE DYSENTERY OF BENGAL.	CHRONIC DYSEN- TERY.		
	_	Observation in the cases of 160 Europeans.	Observation in the cases of 55 Europeans.		
	The state of the s	Cases.	Cases.		
Liver found	to be altered in	84	31		
11 11	contain abscess in	21	6		
" "	be enlarged in	40	5		
" "	be gorged and turgid in	4			
" "	be small in	7	8		
" "	be pale in	26	11		
0 0	be granular in	22			
" "	be softened in	12	1		
" "	be indurated in		4		
" "	contain cicatrices in	5 3	1		
**	contain hydatids in		1		
-	be nutmeg in		6		
11 11	be cirrhosed in		1		

	· · · · · · · · · · · · · · · · · · ·						C	ases.	
The i	leum is noted as ov	er-vasc	ular	or co	ngest	ted in		21	
	t ulceration and abi								
Inas	state of sphacelus in	n .						1	
Muco	us coat of the st	omach	over	-vasc	ular	or s	of-		
ten	ed in							4	
Muco	us coat of the stom	ach ulc	erate	d in		-		1	
	intestine, chiefly						nd	-	
rec	tum ulcerated in all	the ca	ses.		-				
	m, transverse and			colo	n. fr	ee fr	om		
nle	eration in	debeen	9		.,		-	3 (	only
Large	intestine, but gene	rally th	e cm	eum.	perfe	rated	in	8	Jana
Ileo-c	ecal valve ulcerate	ed and	destr	oved	in	,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,	***	3	
	uration of appendix								
roc	orded in	vermi	ioi iii	is no	une	Omm	,	1 .	only
Thiel	ening and stricture	of into	etino						my
Maga	ntion . nteric glands enlarg	and on i	n flow	od in				17	
Splan	nerie giands emarg	ged of 1	ппап	led II				6	
Spice	n enlarged in . eys diseased in .			*				2	
Kidne	eys diseased in .	12		*				2	
In th	e 55 cases of chro	nie des	ente	re th	e fol	lowin	or fo	ota	ore
ecorde		me dy	SCHOO	1, 01	101	HOWIN	B 1a	010	arc
CCOLUC							C	ases.	
Large	intestines were ule	perated	in						
Color	contracted in .	cerateu	111						
Cmen	m nearly closed in		•					1	
Color	n nearly closed in								
Color	perforated in . ach noted as unhea	labor in							
Chass	ach noted as unhea nic inflammation an	d as fra			1.			30	
Unito	ne innammation an	d some	ming	III			-	2	
Dalas	ased vascularity in							2	
Pylor	us in a state of abra	asion in			*		*	3	
~ 2	, , , cano	er in	4		*			1	
Small	intestines noted as	unhea	Ithy :	in				12	
Ulcer	ation or abrasion of	the ile	um i	n				2	
Mese	nteric glands enlarg	ged in						16	
Splee	n enlarged .	16		*	151	10		4	

After this summary, we shall be prepared to learn, that out of 2,044 admissions from dysentery into the General Hospital of Calcutta, from 1830 to 1850, there occurred, according to Dr. Macpherson, 457 deaths, or 22.3 per cent.; and "the extremes of mortality have been 14.8 in 1833, and 34 in 1845.

Having shown so much of the antecedents of the cases of European sufferers from tropical dysentery, we have now to contemplate their condition as they present themselves in England.

val im ; and everal angereventh ttacks licine\_ of the reguright of the rbutie th are ning." ne Sah ed an ss the stool= vomithes to pium iment IS COMEbed as muco-

i Longe completion in the cin the cin

twenty-four hours. Thinks the pills cause nausea, and I therefore substituted two grains of gallic acid for the sulphate of copper. The complexion and general condition are improved, and he is allowed solid animal food.

May 19th.—He says he feels quite well, with the exception of "a sensation of cold in the bowels, and of a feeling that they would be moved by the use of cold drinks." Has but one consistent motion in the twenty-four hours. Ordered four grains of powdered nux vomica three times a day in pill, and allowed a tumbler of bitter ale at dinner.

June 5th.—He declares himself now "really well in all respects, and that he has acquired his ordinary condition and weight;" complexion much improved. Ordered a vinous solution of quinine and citrate of iron twice a day; and in the event of the bowels becoming constipated, he was directed to use an enema of cold water; when a more effective aperient became necessary, he was ordered to take a mild dose of magnesia and rhubarb with inecacuanha.

Remarks.—The case of this young officer had in India been of a dangerous character, for he voided exudation-tubes of the croupous character, and frequently also shreds or dark-coloured flakes of dead mucous membrane. The nervous temperament and the scorbutic complication also added greatly to the difficulties of early treatment. But, on arrival in England, the health had somewhat improved. There were then but slight traces or remains of ulceration of the mucous surface of the larger bowel, and there was no hepatic or scorbutic complication. The treatment was therefore simple and freed from difficulty. I believe that the formula originally prescribed would have conducted his case to a successful and safe conclusion; but such were his terrors of what he would designate "dysentery" that I felt constrained to use active mineral and vegetable astringents in order to appease his fears. The powdered nux vomica appeared to act very beneficially, and I generally find it so in the irritable state of the colon, which so frequently follows the ulcerative dysentery of tropical climates.

In the instance of this young officer, as in so large a proportion of all who come home sick from India, the appropriate winding-up to the treatment consisted of chalybeates, given with a view to remove the anæmic condition; but in this completion, as in the curative endeavour, the salutary effect was greatly retarded by the temptations of the London clubs.

Case II.—October 18th, 1849.—Assistant-Surgeon ——, of the Indian Army, aged thirty-eight, had resided eight years in India. His complexion was naturally sallow, temperament phlegmatic, habit abstemious. "From childhood," he says, "until shortly previous to the present dysenteric attack, he had been subject to

giment side. overed. seized paid\_ nt, he sufferok no udden three nentaused = pium. of pill rse totober\_ twelve lood. by the THATY. emen ention ectum ses of onselt me he in the flund ted to ended freeze of the e 30c3 aplete meral any serious organic change in the liver, or other viscus." Dr. Watson ordered nitro-muriatic acid conjoined to gallic acid, the use of which he continued until the 30th of August, when there was very copious purging of thin, watery, and bloody fluid, which

was "squirted out with violence."

Towards the end of December, not having derived benefit from any treatment, the patient went to Hastings, where the late Dr. Mackness directed him to take bismuth, logwood, opiates, &c.; but no improvement resulting, he returned to Croydon at the end of March, 1851, in a very feeble and exhausted condition, from the constant drain of large fluid bloody motions. By the 20th of April, he was actually sinking, and could not turn in bed without assistance. I now visited him at Croydon, and found him voiding enormous quantities of bloody serum. In consultation with Mr. Bottomley, a pill, composed of diacetate of lead and opium, with a solution of the same medicines in the form of enema after each motion, were ordered. On the 23rd of April Mr. B. reported to me that the patient was "living only from hour to hour," and certainly neither of us looked upon his recovery as at all probable; so far from it, indeed, that we spoke of the desirableness of a post-mortem examination. The patient had deliberately made up his mind to death, and he took what he thought to be a final leave of me. The means last used, however, were strictly persevered in against all disadvantages; and the result was remarkable. Recovery took place from this hour. On May 13th the patient states, in his notes of the case, that "the remedies last ordered seemed at once to check the purging, and the injection was only used twice or thrice, and retained for a short time. The bowels had now become so costive that tepid-water enemata were requisite in order to relieve them." On the 9th of July the patient visited me in London, when I ordered twenty minims of the pernitrate of iron to be taken in water three times a day, and which was afterwards changed for the sesquioxide.

Under this treatment, aided by a nightly anodyne, the anæmic condition of the system was gradually overcome; the bowels were moved by enemata, sleep was secured, and the anæmic state of the pulse disappeared. By the 24th of March, 1852, the patient had increased in weight from eight to twelve stone, while he and his family declared his then condition of health better than it had been for years. His necessities, as well as his inclinations, pointed to his return to India, "where," he said, "I flatter myself I know my duties as surgeon in the service; and I feel sure that I am at any rate much better qualified to discharge them than I am for private practice in this country, of which I have seen little and like less." He proceeded to India, and enjoyed good health for a

year or more, when he was cut off by epidemic cholera.

fluids and tissues, so as to prove a serious detriment and hindrance to cicatrization. Thus we perceive how both the complications here noticed must be attended with dangers and difficulties.

Whether the condition of the mucous membrane of the intestine be that of relaxation, thickening, or ulceration, we shall find our best means of cure in the mineral or vegetable astringents and

tonics, with or without opium.

Robert Jackson says, that where ulcerated surfaces are healed, or in a healing state, the foundations of congestion having been removed, and where purging continues from relaxation or irritability, kino, catechu, and simaruba have appeared to him "the most useful." But he found it necessary "to vary the form of tonic," as the effect, "often considerable for a few days at the first trial, is soon lost." . . . "It is to this condition of disease that the arsenical solution particularly applies; and it is here a remedy of the greatest value." In the great majority of instances, I find it necessary, in order to secure the full effects of both astringents and tonics, to conjoin them with opiates, in more or less minute proportions, according to the urgency of the symptoms.

Whether we employ the mineral astringents—such as the diacetate of lead, sulphate of copper, nitrate of silver, or quinine, arsenic, or mercury, the mineral acids, the vegetable tonics or astringents, the rule of shielding them with one or other of the

preparations of opium, is equally applicable.

In every stage of dysentery, whether acute or chronic, we must endeavour, by every means, to quiet irritation and movement of the bowels, as by such quieting means we most certainly guide the patient towards the healing of abrasions or ulceration of the intestinal mucous membrane. I repeat that astringents and tonics, mischievous in the inflammatory stage, are here much to be relied upon. A bland and spare diet, proper medicaments, including warm clothing and warm baths, mental and bodily rest—these are the rational means found on experience to secure the end in view. But though this principle and these means ought never to be lost sight of, we shall have occasionally, in the management of chronic dysentery, to use gentle aperients, with a view to remove accumulations of matters in the cæcum and colon, organs often thickened, narrowed, and weakened by previous disease.

These last means will be more specially necessary as we approach the period of cure; and I have found castor-oil, in small doses, very useful. Occasionally, however, the oil produces griping and tenesmus, and thus proves very mischievous; so that the question of the agreement or disagreement should always be ascertained beforehand. The aperient which I have found most generally to agree, is a combination of rhubarb powder and magnesia, with those of columba and ipecacuanha, and bicarbonate of soda; a combination of magnesia, sulphur, and ipecacuanha, with some

the tenesmus of months, or even years, no wonder that the sphincter and the entire perinæum should prove inordinately flaccid. This state of perinæal relaxation causes very distressing uterine descent likewise, even in very young females. The pain and irritation from hæmorrhoidal tumours keep up a disposition to void the contents of the bowels, and to ineffectual straining, while the loss of blood so frequently attendant on these affections greatly aggravates the concomitant anæmia. Along with relaxation and anæmia there exist here immediate mechanical sources of injury: the descending mucous lining, or the tumours, as may be, from within or without the bowel, and the hypertrophied integument external to the sphincter, keep that muscle in a permanently relaxed or halfclosed state, impairing its tone; and the hæmorrhoids, whether blind or bleeding, proceed from bad to worse. In general, it will be found proper to wait for some improvement in the general health before any kind of operation is had recourse to; but where the bleeding from the hæmorrhoid is depressing, the application of nitric acid to the bleeding point or points is advisable. In other cases, and where the habit is in a better condition, I recommend the operation of Hey and of Abernethy-that preferred also by Liston and Mr. Fergusson. This operation, always safe, and generally successful, consists first, in replacing the obtruded internal pile, or relaxed mucous membrane, with the oiled finger; and secondly, in drawing out the loose hypertrophied external integument with a common vulsellum, and in clipping off portions of it with strong scissors, the directions of the excisions corresponding with that of the bowel-the blades of the seissors, according to Mr. Syme of Edinburgh, being directed from the circumference towards the centre of the anus. This operation at once removes the mechanical external impediment to the natural contractile action of the sphincter, while the cicatrices resulting from the wounds afford permanent buttresses to support both the sphincter and perinæum, or parts immediately external to and around the anus.

In cases where the relaxed, thickened, or otherwise altered mucous membrane, with or without condylomata, protrudes, the application of nitric acid, as first recommended by Dr. Houston of Dublin, or else the galvanic cautery, should be made. Like every operation for the cure of hæmorrhoids, the galvanic cautery and the nitric acid require care in the application;—and so does the ligature, which some operators still prefer. The acid should be of the strongest, so as utterly and at once to destroy the vitality of the parts to which it is applied; and it is better to repeat the application to small points at a time than to cover too large a surface at once with the acid. The same rule applies to the use of the actual cautery, likewise a safe remedy when used in the manner recommended by Mr. Henry Lee, of King's College

Hospital. Mr. Lloyd, at St. Bartholomew's Hospital, habeen in the habit of smearing the whole surface of membrane, and likewise the congested hæmorrhoidal with solid lunar caustic, it is said with uniform success rarely finds it necessary to repeat the application above four times.

But whatever operation may be preferred for the r hæmorrhoids, condylomata, or altered and prolapsed much brane, the clipping of the hypertrophied loose folds of it around and external to the anus is a measure of absolute to the cure. It should constitute the first and earliest too, whatever else may be ultimately required in respect conditions within the sphincter. The clipping external by restoring the free action of the sphincter, is very ofte of what are termed internal piles, and this circumstance itself secure for it a priority and a preference; but whe pens otherwise, the early removal of loose and thickene integument, impeding the contractile action of the spl necessary to the eventual success of any and every mean directed to within the anus.

I state these circumstances upon observations of an nature, both in India and at home. But I would obser of these cases, and, indeed, of all cases of returner equiring surgical operations, that to treat them wirequires a careful regard to their peculiar habit—care by and care in the after-treatment being necessary. The system of the returned Indian is excitable, and, if but from the East, he is often an amic—conditions of the which obviously require much attention. I saw a lady, but arrived in England, in whom the opening of a small tumour nearly proved fatal; and, in the case of a gentle same dangerous symptoms resulted from the removal of internal hamorrhoidal tumour, by ligature.

The benefits derivable from horse exercise, in the pand cure of hæmorrhoidal affections, are not generally un yet they are very great. Riding on horseback acts bene accelerating the course of circulation along the veins brings the sphincter and adjoining muscles into alternathus contributing, the one action with the other, to preve engorgement, and to impart tone to the relaxed muscles.

I propose now to conclude with the histories of some interest, with a view further to illustrate the treatment of dysentery.

Case III.—April, 1854.—Captain ——, aged thirty, h in India six years, enjoying excellent health. In April, corps was ordered to Rangoon, where he underwent me

sure and fatigue, being indifferently fed at times. In September, 1853, while serving at Prome, he was seized with acute dysentery. He was actively treated, with mitigation of suffering; but purging continuing, he was ordered home, and arrived in England in February, 1854. Purging still continued to distress him, and he obtained leave to reside with his relatives. Here he took calomel to salivation, with apparent benefit at first; but after three weeks all his distresses returned, and he came to London on the 18th of

April, 1854.

He was then pale and anæmic, and much emaciated; skin cool and damp; pulse feeble and slow; tongue clean. He was purged from eight to ten times in the twenty-four hours, with much straining, voiding quantities of mucus streaked with blood. There was no hepatic complication, but he had tenderness on exploration along the cæcum and colon, with feeling of soreness in the course of the rectum. Ordered the following mixture, of which two tablespoonfuls were used three times a day, and five grains of Dover's powder, with two of gallic acid, at bed-time:-Infusion of simaruba, seven ounces and a half; extract of logwood, and tineture of krameria, of each, two drachms; dilute nitric acid, and dilute hydrochloric acid, of each, one drachm; tineture of opium, one drachm and a half. The patient was at the same time enjoined to avoid all exercise, to observe great caution in diet, especially as to quantity, and the use of warm clothing and the warm bath at bedtime.

April 28th.—Had a visit from the patient, stating that though he had walked out daily during cold weather, he felt then "quite well," and that after the fourth day he had discontinued the night pills, contenting himself with the mixture. He added that he had but one formed motion in the twenty-four hours, and that unattended by either blood or mucus in the discharges, and that straining and all uneasiness had disappeared. Ordered the continuance of the mixture twice a day, with a view to obviate any tendency to relapse.

July 18th.—Says he has been living in the country, and that by exposure to cold while fishing, and by irregularities in diet, he has brought back "looseness of the bowels on several occasions, but no dysenteric feelings." He then had recourse to his mixture, and always with complete success. He was again ordered to take five grains of Dover's powder, with two of gallic acid, every night, and urged to use temperance in diet. This course of medicine he continued for a month, with how much temperance, however, I know not; but he became well.

Remarks.—It were unsafe to conclude that, in a case such as this, because it yielded so readily to treatment, it was therefore free from ulceration, and from the more grave dangers of secondary purulent deposit in the liver. The appearance of the intestinal tion did of conily was specific

went to e years ttack of since , which alth in

d with d him alaya, 1844, is then is since, and on tion

nce of in the

febrile where

conuring

s; the h pain feeble, ed by frothy

interionally about losing all effect, each in succession. Most of the vegetable and mineral astringents were used in conjunction with Dover's powders, morphia, simple opium, and bitter tonics, including the nitromuriatic acid with opium and quinine,-all apparently with no other result than arresting the progress of the disease. On two occasions a grain of blue pill with morphia, given at bed-time, appeared to promote the function of the liver, to improve the secretions, and to correct the irritable condition of the mucous membrane; but at length extreme rectal distress supervened, with straining and inability to retain the contents of the bowels for an instant, when once the disposition to void arose; the matters discharged being extremely offensive, and containing mucus and pus. A drachm of finely-powdered charcoal in barley-water was now exhibited night and morning in the form of enema, with a remarkably soothing effect, while gallic acid and morphia in pill were continued three times a day. The effect of the charcoal was the most marked of all, the relief being very decisively evident; and by the middle of August the patient was sent into the country for change of air.

In February, 1855, Major —— called on me to ask my opinion as to his fitness to accept a command in Turkey. I then found that for some time previously his bowel complaint had been changed for constipation, and that he had gained considerably in flesh and strength, although not to the extent to permit of his

entering on the active duties of his profession.

Remarks.—The case of Major — is remarkable for the severity of succeeding tropical diseases during two-and-twenty years, and for the severity of the eventual dysentery. Originally of a powerful constitution, he went through an enormous amount of illnesses, one after another, without sinking. On arrival in England, and for months subsequently, his case appeared hopeless; but by a persevering course of treatment, frequently changed to meet the altering circumstances of the case, by home nursing, and by the utmost attention to diet, clothing, &c., this excellent officer eventually recovered.

In 1827, he came to England, ailing, and returned to India in 1830 restored to health. He served in the Affghan war of 1842-43, and returned to India in the enjoyment of good health. On being removed to Calcutta, however, in 1844, he was speedily seized with dysentery. Two short voyages to sea "did him some

I saw the patient, in all, five or six times only, in the early summer of his arrival. The bowels were at this time irregular—sometimes relaxed, and again constipated, blood and mucus being occasionally observed mixed with the contents of the bowel; but there was neither pain nor straining in the act of defectation. The skin and pulse were natural; the tongue clean; the appetite good; the intestinal and renal excretions were natural. There was no tenderness on pressure in any of the abdominal regions—no enlargement of the spleen or liver; and the disorder, slight as

it appeared, was confined to the rectum.

the

s so

the

HIE

ore

arly

nes

our

DII-

fu-

nd

Great care in diet and clothing was enjoined, avoiding all exercise, other than a gentle drive in a carriage; and in the way of medicine, he was directed to take compound powder of kino in infusion of simaruba, confection of senna with sulphur and bitartrate of potash being used when the bowels required aid. An enema, to be retained in the bowel, was directed to be used at bed-time, composed of alum in decoction of oak-bark.

It was six months from this time that I next saw General —, by accident, when he told me that he had been residing in the country, and been there "treated for the liver, by means of calo-

mel, which had done him much good."

He now consulted an eminent physician in London, under whose care "he rallied much during the winter of 1852-53." In the spring of the latter year the patient was attended in the country by another practitioner, who, on careful examination, found the spleen dark, indurated, and enlarged; the liver containing a large abscess. The physician who saw most of this case in London gives it as his opinion, that "the suppurative action in the liver was, in a great measure, if not entirely, secondary to the ulceration of the lower bowel;" while the other physician, who was consulted early, gives the following summary:—

"1st.-Injury to the intestinal surface, from tropical influences

of long standing.

"2nd .- Injured surface ulcerating in the last attack.

"3rd.—Abscess of the liver derived from absorption of pus from

the ulcerated intestine."

This gentleman adds, that when he was consulted, he could, "on a careful examination, detect no sign whatever of liver abscess;" nor was there indeed at the time of my first seeing him, any indication of hepatic disturbance, functional or structural.

Remarks.—The two cases here last recorded present a remarkable contrast—the first being immeasurably the severer of the two. The patient was in the last stage of exhaustion—bedridden; and when he stood erect, there was an instant call to void by stool.

In the last case, on the contrary, the symptoms, on arrival in England, were so trifling as to constitute to all appearance but a slight indisposition, the patient having been in London a fortnight before he thought of consulting me respecting his state of health; yet his disease proved eventually fatal. Here we perceive, in one out of many cases, the dangers attaching to chronic dysentery,

even when it appears to be of the mildest character.

Suppurative phlebitis, and consequent purulent deposit,—to use the ordinary but unsatisfactory expressions,—may here be inferred, but what the actual morbid process consists in, does not, as yet, appear to be demonstrated. Purulent secondary deposits are mentioned by many surgeons as resulting from the application of ligatures to hæmorrhoidal tumours; and this is but another instance of the so-termed purulent deposit from ulceration of the mucous membrane of the large intestine. Nor are such abscesses in the liver limited to cases in which ulceration exists in the mucous digestive surface; for they are observed to result from other lesions as well. On the 15th of March, 1854, a stableman, aged forty-four, died in St. George's Hospital, of "phagedena of the great toe; secondary deposits in the liver, and pleurisy."

On questions relating to the medical treatment of the case of General ——— I shall offer no opinion, as the patient came under my personal observation in the very earlier stage of his disease only, and at a time when his condition promised a speedy

recovery.

nature. te comder and atment will end v see im ring the ngestion r proper mie disresummer the prodomina s patient nstances that time ne other quality e, as in lifficul upidly to instan IS TOTAL tive cli

allow,
age
residence
ier is
ce of
s appliates,
surface
fluids
eces
118

nto acuar

tion, through the agency of cold, is also much increased where the individual has suffered from malarious fevers, and from consequent disturbance of the balance of circulation. The feeling of the patient is one of epigastric fulness, uneasiness, and oppression, rather than of pain of the region of the liver. The organ is oppressed by an accumulation and stagnation of blood, while its function is paralysed. With all this we have frequently a weakened condition of the heart and arteries, and a diminished power of reaction, consequently. It is only, however, when the hepatic congestion is associated with a similar condition of the cerebral organs, or of the lungs, that danger to life need, under careful management, be apprehended. In the cerebral, and in the thoracic complication, there is indeed a pressing and immediate danger, even under the best remedial means known to us.

I find that in the first part of this work, when speaking of the congestive fever of the cold season in Bengal, the following observations occur; and I here quote them, to show how prevailing must be the liability to hepatic congestion in European invalids coming at once from hot climates into the winter and spring seasons of England :- "It is believed, that in animals whose pulmonary system is less perfect, there is a greatly increased quantity of blood transmitted through the liver. In hot climates, then, where respiration is less perfectly carried on than in cold ones, owing, according to Tiedemann and Gmelin, to the greater rarefaction of the air in warm regions, a vicarious decarbonization of the blood is established by an increased flow of bile; and hence it is that the function of the liver, weakened and torpid, in proportion to the excitement of the hot and rainy seasons, becomes disposed to congestion, or inflammation of its parenchyma, during the cold season; and thus are produced the dangerous states of disease noticed. An irritable or inflamed state of the mucous digestive surface is also a frequent complication; and these two together constitute the great dangers of our congestive fevers of the cold season, as well as of those of the autumnal fevers of the more unhealthy countries of Europe."

There is no more efficient cause of congestion of the liver than antecedent torpor of its function—especially where this last disorder has been of long duration. To the tumult of the nervous, vascular, and secreting functions, within the tropics, has now succeeded an exhausted condition of all three. The system at large, and the organ now principally at fault, have lost their power of resisting the cold and damp atmosphere of Europe. To be more precise, the circulation through the skin, and also its function, which had been raised to the greatest degree by the high temperature of the tropics, is reduced to the opposite extreme by the cold and damp atmosphere of our northern climate. The blood

purgative draught of compound decoction of aloes, infusion of senna, and jalap powder, with aromatics. After the third exhibition, these medicines produced copious biliary discharges, so as rapidly to diminish the volume of the liver. This plan of treatment was accompanied from the beginning by sinapisms to the hepatic region, warm baths every other night, and friction to the entire surface of the body. On the ninth day congestion had so far subsided, through copious excretion, that the treatment was concluded by the nitro-muriatic acid, in combination with fluid extract of taraxacum, given three times a day, and purgative pills were continued at bed-time, composed of compound extract of colocynth, compound galbanum pill, each half a drachm, with two minims of croton oil, made into twelve pills. Of these, two were given at first, and then one, until they were gradually discontinued. Warm baths and friction were used at longer intervals, till every symptom had vanished, when exercise in the open air, and mild chalybeates, concluded the treatment. The diet at first consisted of farinaceous articles, with thin soup, vegetables, and fruit; but, as health improved, animal food with diluted wine was allowed.

CASE II.—Lieutenant ——, of the Indian cavalry, states that soon after parting with his wife and child, who were obliged to leave India on account of ill health, he became depressed in spirits, and lost his appetite and flesh rapidly. Pain and tumour in the right side soon indicated inflammation of the liver, for which he was very largely bled, followed by a course of calomel purgatives, and blisters. Abscess, however, resulted, and the matter was discharged into the colon. He arrived in England in the month of April, in a very enfeebled and emaciated state, having slight cough, with little appetite, the digestion being much impaired, and the bowels constipated. There was no fever, the pulse being languid, the skin cool, dry, and harsh to the touch. The general condition was anæmic.

I ordered the decoction of taraxacum with dilute nitro-muriatic acid, mild aperient pills devoid of mercury, warm baths, and powerful friction to the skin; the diet to be spare, light, and nutritious, with a little bitter ale to dinner. Strict injunctions were

given to avoid exposure to cold.

On this plan he rapidly recovered flesh, and by September his

appearance indicated returning health.

I was requested to visit Lieutenant — in the January following, in consultation with Mr. Squibb, of Montague Place. I then learned that, three weeks previously, the patient had taken a long walk, after which, while still heated, he went to hear the afternoon service in a neighbouring church. Here he felt chilled,

having all along been but little restricted. For the last month he has been residing near London, suffering greatly from pain of the right side and shoulder. His days have been passed sitting by the fire side, supporting his head on his hands, while his elbows rested on his knees. His nights have passed in a burning fever,

but he has had neither rigors nor perspirations.

He drove to my house on the 16th of September, and walked with the body bent forward on the pelvis, sitting in the same attitude, and breathing rapidly. His complexion was of a leaden yellow hue; the skin dry, and of a pungent heat; the pulse feeble, and at 108; the abdomen was tense, round, and hard as a barrel, with an enormous bulging of the right hypochondrium, the liver extending four inches downwards from the margin of the false ribs, and pushing up into the right cavity of the chest. Percussion and exploration of every kind gave no pain but only a slight uneasiness. He complained of headache, the eyes being suffused, yellow, and muddy; the tongue deeply coated, and red at the edges, the urine was of a madder-colour; the bowels constipated. The nature of his case was explained to Major ---, and he was urged to remain in bed, to which he seemed averse. He was ordered to take two pills composed of extract of colocynth, calomel, and ipecacuanha, with a quarter of a grain of morphia at bed-time, followed by a purgative draught, using a warm bath every other night; the diet to be purely farinaceous.

On the 19th September, the patient again visited me, expressing himself as being much relieved. For the first time during months past he had had refreshing sleep; the skin was somewhat relaxed; the pulse 94; pain of the liver, head, and shoulder, somewhat diminished. The medicines had acted powerfully on the bowels, the urine remaining as before. Major — was now informed that his treatment could not be safely or effectively conducted abroad; and that, in such cold weather, it was dangerous to leave his house, or even his bed. The former medicines were

directed to be continued.

Sept. 21st.—Called to see Major—at his residence, where I requested a consultation with two medical officers who had lately been in attendance on him. It was here agreed that mild mercurials, with sudorifics, and small doses of morphia, should be exhibited over night, followed by active purgatives in the morning, while warm baths were used at night, with frequent hot fomentations to the side during the day. The rapid and feeble state of the circulation, and the general exhaustion, precluded bloodletting in any form, either general or local.

27th.—On this plan of treatment great relief of symptoms ensued; the complexion improved; the skin became soft; the pulse was reduced to 80; while the tumid and tense state of the abdomen

tation of the sea voyage, aided, perhaps, by nausea and sea-sickness. But in the second congestive attack no such favourable influences existed. I have confined my narrative of this interesting and instructive case to a bare recital of facts, such as they were reported by the patient, and to a detail of the symptoms, which I personally witnessed, avoiding all comment on antecedent circumstances; but I feel bound to state, that I received the most candid co-operation and able assistance from the two medical men with whom I consulted in the treatment of this distinguished but unfortunate officer. Up to the time when the mild influence of mercury was evidenced by the action on the gums, and by the mercurial fætor, so much improvement had taken place as to warrant the balance of opinion to lean towards a favourable result; and so, also, with respect to the action of the nitro-muriatic-acid bath, up to the fourteenth day before death, the unfavourable circumstances being, that by none of the means used was a healthy secretion obtained either from the liver or kidneys. This was the result anxiously looked for, and never obtained. The important practical question here was, When did the suppurative process commence? My own impression was, first, that the formation of abscess had taken place within the fortnight before death; and secondly, that the abscess was of but small extent; for in India, in the instances of large hepatic abscesses, I have never seen one case in which rigors, colliquative sweat, and a permanently furred tongue, were all absent, as in this instance. The pain of the right shoulder, amounting, during several weeks of July and August, to agony, indicated the seat of the active inflammation to have been centred in the upper convex surface of the liver, involving its peritonwal covering, for there is seldom much pain, and very often none at all, when the parenchyma alone is the seat of the inflammation. In general, these circumstances point to the actual seat of the subsequent abscess.

Major-General ——, aged seventy-one, served thirty-six years in Bengal, and has resided fifteen years in England. Eight or ten years ago he is reported to have suffered from repeated attacks of intermittent fever, and for which quinine was administered largely. These attacks were finally overcome by the use of calomel and purgatives, quinine having proved of no avail. His habit is spare but muscular, and in diet he has been through life abstemious to the degree of extreme, having avoided wine and malt liquors altogether for years, and used animal food during the last two years but sparingly, once in twenty-four hours. He has been used to active walking exercise all his life; but during the last year official duties have interfered much with this salutary habit, while they have agitated and over-excited his mind. Within

the last month he has been often exposed to severe cold at night when returning from dinner parties. He relates that his habit has always been to use cold ablution in the morning; but owing to the severity of the last winter and spring, and to the cold water proving ungrateful to his feelings, he has for some months substituted ablution with warm water.

On getting out of bed this morning, April 4th, he felt giddy t staggering, with nausea, and some degree of faintness; and, a the weather was warmer than of late, he bathed in cold water took his usual breakfast, and drove to his office, where, during th last two days, his mind was unusually excited and agitated. T add to this distress he states, that for more than an hour of th forenoon he was persecuted by solicitations with which he could not comply. At noon he was seized with a violent rigor, accom panied by extreme pallor and shrinking of the surface of the body, which was soon followed by vomiting. In this state h drove to my house. Violent shivering, with sense of extreme cold, lasted for two hours, but by warm drinks, and warmth applied externally, imperfect reaction was established, accompanied by moisture of the surface. The tongue is loaded with a thick whit fur: urine for some time past scanty and surcharged; bowel habitually regular, but of late the excretions have been scanty and clay-coloured.

On examination of the abdomen, the region of the liver wa found tumid, bulging, and very painful on pressure, so much so that he started when it was touched, the abdomen throughout being puffed and inelastic. He felt no pain any where else. A ful dose of calomel, with James's powder, was given immediately At 7, p.m., reaction still continued moderate, with a warm gentle perspiration. Ordered two pills, composed of compound extract of colocynth and calomel, to be taken at bed-time, and a purgative draught for the early morning.

At midnight I was summoned to the patient, and found that, or getting out of bed, he had fallen on the floor without the power to recover himself. The face and neck were of a reddish livid hus with deep somnolency, and difficulty of being roused; skin ho and dry; pulse full, but easily compressed. I found that anothe medical gentleman had been summoned at the same time with me and we agreed at once that the patient should be cupped from the neck, while a blister was applied to the right hypochondrium. The cupping abstracted twelve ounces, with marked relief from cerebral oppression; but it was not till copious and repeated dis

patient was so far restored as to reply to questions.

April 5th, 10, a.m.—General —— is now perfectly restored to consciousness, and expresses himself as much relieved, especially

charges of vitiated bilious evacuations from the bowels, that th

from the sense of pain and weight in the side. No headache; skin moist and warm; pulse 90; secretion of urine copious, but of a deep porter colour. Ordered a pill, composed of two grains each of calomel and colorynth, to be given every two hours. Diet farinaceous.

9, p.m.—General —— was seized with a violent rigor, accompanied by stupor, but both speedily gave way to diffusible stimuli, ammonia, and the use of warm drinks. The following reaction was but trifling, and it now became apparent that ultimate recovery might be regarded as more than ever doubtful. In consultation with Drs. Chambers and Watson, simple calomel was ordered to be given frequently during the night, while a blister was applied to the nape of the neck.

April 6th.—After midnight, relaxation of the surface with flagging circulation rendered the use of wine and soup necessary. The powers of life are now (10, a.m.) sinking, and he is dependent on the frequent use of stimulants; consciousness but partial.

Died at 6, p.m.

Remarks.-Here we perceive hepatic congestion of the most severe character, followed by the same condition of the brain, and accompanied by diminished power in the heart and arteries, all taking place fifteen years after the return to the native climate of the patient. The organ primarily affected was the liver-that which, during youth and manhood, had been most excited in its function—and that, too, in a person of unusually abstemious and active habits of life. It would seem that with some persons no amount of care will prove a sufficient protection against hepatic disease. All the circumstances of constitution, and of habit of life, mental and bodily, were in favour of this gentleman's living in health to an advanced age; yet were they all frustrated by the antecedent circumstances of service, and of long exposure to the influence of a tropical sun. In these respects, and in respect of the influence of mental excitement on a nervous system, very impressible in its nature, the case is most interesting. From the time that General — experienced discomfort in the use of the cold bath, it may be inferred that hepatic congestion and functional disturbance were present; and there can be no doubt that the great and repeated mental excitements of the past year laid the foundation of the cerebral complication, which proved fatal, and but for which the hepatic disease would easily have been overcome. It is remarkable how often we find, towards the decline of life, and when the vigour and the balance of the functions have been impaired, that the liver-the organ most excited in hot climates-assumes disease, and that in persons who had not previously suffered from actual disease of that organ, either while residing in hot climates, or during a protracted residence at home. Numerous such instances have come under my observation, and

Anæmia, and consequent torpor of the liver, often follow on long-

continued hyperæmia of that organ.

In all cases of torpor of the hepatic function we find the process of digestion and assimilation much impaired. Owing to defective quantity and quality of the gastric, biliary, and pancreatic fluids, the nutritive functions are manifestly impaired. The arrest of the cuticular discharge, so abundant within the tropics—so powerful, too, in carrying off acids and other redundancies from the blood—occasions in many persons, on their return to Europe, an excessive acidity of the stomach, not to speak of the other more import-

ant alterations of functions already referred to.

It may be said, truly, that most of the influences of climatethat most of the diseases resulting from these influences—and that many of the habits of life of Europeans in India, and other hot climates, tend powerfully to disturb the functions of the liver, and thus lead ultimately to serious disorders, and to actual disease of that organ. The summary offered by way of introduction to these essays will have furnished the reader with examples of numerous climatorial causes of hepatic disorders and diseases; but there is one other efficient cause, much overlooked, but which, on the score of extensive influence, must not here be omitted-viz., mercury, internally exhibited. It has been observed in Europe that persons who have used mercury, either in the liberal or rapid manner, or as a protracted alterative course, have frequently been afflicted subsequently with severe forms of hepatic disease. There can be no doubt that, through the repeated use of this mineral, even when necessarily exhibited, the hepatic function is inordinately stimulated, to be proportionately enfeebled thereafter. The action of mercury on the liver may be viewed as having a close analogy to the primary excitement, and subsequent exhaustion, of the functions of the stomach, produced by dram-drinking.

In referring to the treatment of the troublesome and obscure affection now under consideration, we must recollect, not only that we have to deal with torpor of the secreting and excreting power, but with viscidity and adhesiveness of the secreted fluid, and also with diminished power in the gall-bladder and ducts to eliminate their contents, with the addition, perhaps, of thickening of the parietes of the vesicle and its canals—thus affording additional and mechanical obstruction to the discharge of the bile. In some persons this torpor of function is so great, and the case proves so little amenable to treatment, that we are forced to suspect some alteration, more or less extensive, or some permanent injury to the

elementary secreting structures of the liver.

The slightest reflection on the physiological and pathological changes, here but hinted at, will show the great difficulties that meet us in our endeavours at effective treatment. What little bile

This addition to the dandelion proves in many cases powerful in effect as regards the function of the liver, and generally tonic to the digestive organs. The addition of the acids proves specially useful in cases where tropical dysentery, or diarrhæa, has previously existed, or where an irritable state of the mucous membrane is present. In this latter instance, however, its use is not borne long, and, in many persons, not at all, without the addition of the tincture of opium. Such means, though slow in their effects, are infinitely more appropriate and safe than mercurials, in these chronic forms of disease. They slowly but safely promote increase of secretion, while they impart tone to the functions of digestion.

In treating torpor of the liver, with its complications, it must be borne in mind that our object is to solicit rather than to urge the restoration of the impaired functions, and that such can only be effected with safety by gentle and persistent means, always holding in view the improvement in the nutritive qualities of the blood. We must here be content to look to the actions resulting from the means in use, however gradually brought about, rather than to specific operations. We must even be careful not to push reme-

dies too far.

Whether exhibited as an aperient, or in purgative doses, I have found aloes in its several forms, superior to other purgatives. We are familiar with its irritant effects on the mucous surface of the rectum, and it is my opinion that such a drug cannot be other than active in the duodenum. In effect it is so, and it will be found, when given with ox-gall, taraxacum, the alkalies, the salts of iron, and ipecacuanha with colchicum, that its exhibition is at all times useful in stimulating and evacuating the duodenum, and that it thereby excites the secretion, and facilitates the discharge of bile.

Anæmia is uniformly productive of torpidity of the liver, the biliary secretion being depraved in common with its source, and having, in some cases, more of the character of serum than of bile. It ought here to be evident, that the endeavour to elicit a more abundant and healthy secretion, by mercurials and drastic purgatives, while the nutritive functions are impaired, and the blood re-

mains impoverished, must be alike vain and injurious.

The most cursory inspection of such a patient, the most superficial examination into the history of his case, ought at once to satisfy the medical observer that with mercury especially, he should, in this instance, have nothing to do. The tropical invalid will generally be found to have had quite enough of that mineral during the acute stage of his disease, and when it was necessary to his cure. The true remedies in the anæmic complication are,—the alteratives already mentioned, bitter tonics, chalybeates, mild

2. The patient should remain in the bath from fifteen to twenty minutes, a can of hot water being kept at hand and added in small

quantities at a time to maintain the temperature.

3. While the patient is in the bath, hot and dry towels should be provided, in order that immediately he leaves it, the body may be quickly and thoroughly dried; after which the patient should at once retire to a well-aired and warmed bed. To prepare the second and the following baths, remove on each occasion about one-third of the liquid, then add one quart bottle of the dilute acid, and sufficient hot and cold water to raise the temperature to the proper degree. Should the bath excite too much irritation of the skin, less than one bottle of acid may be employed on each succeeding occasion.

(II.)

### THE FOOT AND SPONGING BATH.

1. Two gallons of water are generally sufficient for an ordinary foot and sponging bath, which should be kept by frequent addition of hot water at a temperature of from 98° to 100°. To the two gallons of water, six ounces by measure of the prepared dilute acid are to be added and thoroughly mixed. While the feet are immersed, a warm sheet or some other suitable covering to protect the patient from draughts should be thrown over the shoulders. By means of a large soft sponge, the insides of the thighs, the right side under the ribs (the region of the liver), and the arm-pits, should be constantly bathed; at the same time, several folds of flannel may be immersed in the hot acid bath, and wrapped round the body. These baths should be used from fifteen to twenty minutes, night and morning, and on each occasion it is necessary to attend to the usual precautions of thoroughly drying the body and legs with hot towels before dressing or retiring to bed.

2. It must be remarked that earthenware or wooden baths should be employed, as all other materials destroy the efficacy of the prepared acid. The sponges and towels should after each bath, be thoroughly washed in cold water to prevent their being

destroyed by the acid.

I have been so much in the practice of prescribing this remedy, that, in order to save trouble to the patient and to myself, I have had the above directions printed; and I invariably place a copy in the hands of such as are to be treated. A course of two months of this bath is found generally to restore the healthy action of the liver; nor does it interfere with the use of bitter tonics, or even with mild chalybeates, but, on the contrary, it rather gives effect to their operations. In the severer instances, hereafter to be considered, I have continually to use this means for a much longer period. The use of the bath should be discontinued whenever

abundant, but it is sometimes high-coloured and surcharged. There is no organic disease discoverable; the pulse is feeble and slow.

He was ordered to take a pill every night, composed of watery extract of aloes, with a grain each of the sulphates of quinine and iron. The gentle action of the bowels being thus obtained, he took, during nearly two months, the fluid extract of taraxacum, with bicarbonate of soda, using a light but nutritious diet, and taking exercise on horseback daily. The warm bath, at 96°, was used twice a week, at the bed-side, and just before retiring to rest, while active friction with the Indian hair-glove was applied to the surface of the body,\* great care being observed at the same time to avoid exposure to the cold and damp of our winter and spring seasons. All these directions were rigorously attended to by the patient, but the progress towards recovery was slow. On the return of warmer weather, however, in April, the digestive power improved, along with the secretions, the complexion having assumed more of the European hue. The weight of the body, though much under the average of health, was gradually augmenting, while the power of increasing his exercises, with the enjoyment in them, steadily returned. Mild chalybeates were now ordered, and these, along with an improved diet and increased exercise, brought a more rapid increase of health. The hepatic and digestive functions acquired tone perceptibly under this course of treatment, and in the month of August Major - went to the moors. In October he returned to London, on his way to travel on the continent, and if his weight was still some pounds under his Indian standard of health, he nevertheless admitted that he was more muscular and energetic than when he last enjoyed tigerhunting.

Remarks.—In this case, as in many others of the like nature, the anemic condition was the pervading injury, and the liver partook largely of it; indeed, I believe that the proper pathology of torpor of the liver, in the instance of tropical invalids in general, and where actual disease is not present, may more justly be expressed by anemia of the organ than any other phrase; and the term becomes important if, as I conceive, it represents a true condition. Certain it is, that in the case of Major ———, the most marked and decided improvement took place under the appropriate operation of chalybeates, and so I continually find

it to be in such cases.

<sup>\*</sup> This glove, made from goat-hair cloth (very superior to the common horse-hair glove, which frets the skin severely), is manufactured by Mr. Savory, of New Bond Street, at my suggestion, after the pattern of those in universal and immemorial use in Hindustan, the country of all others where the purification of the skin is most and best attended to.

surface night and morning, were ordered, and gentle horse-exercise was used on fair days, while a moderate allowance of animal food and stale bread composed his dinner. In little more than a month under this treatment, the discharges from the bowels became so consistent as to amount to constipation, when a pill, containing a small quantity of aloes, with the sulphates of iron and quinine, was taken every night. Captain — then proceeded to the country, whence he wrote me, in January, that his digestive power and the alvine secretions had much improved. He was now directed to use a weak solution of quinine and sulphate of iron, with dilute sulphuric acid, while the diet and exercise were increased.

During the spring and summer months, pure chalybeates, gradually increased in quantity, were exhibited, and by the middle of autumn he was so much improved in health and strength as to proceed on his travels. It required three years' residence in England to restore this officer to health, when he returned to India full of high hope and expectation; and after four years he writes me that his health continues excellent.

Remarks.—Looking to the previous fevers, and the diarrhœa that followed them, and also to the existing febrile condition (which latter state I considered as referable to intestinal irritation), I directed the use of the nitro-muriatic acid treatment, as being febrifuge, emulgent, tonic, and refrigerant. In such cases it also proves, along with the taraxacum, an excellent preparative for more active tonics. Captain — 's condition appeared at first to promise an unfavourable result.

Torpor of the liver, like congestion of that organ, is frequently associated with very severe cerebral disturbance. Of this fact the

two following cases will afford illustrations:-

Case III.—Lieut.-Col. —— served twenty-eight years in India, and has been throughout of careful and regular habits. In 1831 he came to England at my recommendation, suffering dangerously from remittent fever, and consequent enlargement of the liver. He remained two years at home, and regained health despite an attack of bronchitis. Colonel —— returned a second time to England on sick leave, in 1849. The official statement of his case, from the surgeon of his regiment, sets forth that, for some years after his first sick furlough to Europe, "he had enjoyed good health," but that latterly he became 'subject to frequent attacks of pain in the head, the slightest irregularity in diet producing it." On the 6th of January, 1849, the weather being cold, and his corps encamped before Moultan, Colonel —— became jaundiced, his urine being of a blood-red appearance, and the head heavy and oppressed. While under the influence of active purgative treat-

ment he was, on the 8th of January, seized with a fit. "He suddenly fell back in his chair in a state of insensibility," and was found by his surgeon lying on the floor of his tent, "his face tinged of a leaden hue, pupils dilated, breathing oppressed and laboured, skin cold and clammy, and his pulse slow and full. He was immediately bled in the right arm, but the blood not flowing very freely the vein of the other arm was opened, and a large quantity of blood was abstracted, when his pulse became small soft, and more frequent. During the bleeding he had two very severe attacks of convulsive movements of the right side, with throbbings of the right corner of the mouth and winking of the eye of that side, the pupils becoming very much contracted. These convulsive fits lasted three minutes."

For three hours he remained insensible, after which sinapisms fomentations, large doses of calomel, followed by purgatives, were used, and he recovered. Though but forty-seven years of age he now appeared (September, 1849), prematurely old, and much emaciated. Hepatic torpor, and torpor of all the functions of the body, accompanied by severe headaches, were now the symptoms complained of. He was completely anæmic, and no wonder, for the blood-letting he declared to have been enormous. In the first winter and spring seasons after his arrival at home he suffered severely after meals from sleepiness, uneasiness, and occasional pain in the head, accompanied by sense of weight and oppression of the epigastric region, and fits of coughing. These symptoms after four or five hours disappeared; but it was long before he regained proper digestive power, healthy secretions, or muscular condition, sufficient to enable him to take exercise without fatigue

Remarks.—This officer was seen by many physicians in various quarters, and various remedies were prescribed. I directed the use of the mineral acids, chalybeates, and latterly of the cod liver oil. All three were followed by some improvement; but two years and a half of furlough proved insufficient to restore the vigour of the digestive functions, and he returned to India in a condition which did not promise that he could render effective service.

Case IV.—Major ——, ætatis fifty, served thirty years in India, and had enjoyed an average degree of good health until within the last six or seven years, when, according to the statement of his surgeon, he suffered "at periods, varying from a week to a month, from severe headaches of a most intense character. The affection commences primarily with a weight and heaviness in the head, which gradually increasing, ceases not until it assumes a degree almost insupportable. During the attack, and until the operation of medicine, he is incapable of moving, and often of lying down, the only relief experienced being in the sitting posture

ment he was, on the 8th of January, seized with a fit. "He suddenly fell back in his chair in a state of insensibility," and wa found by his surgeon lying on the floor of his tent, "his factinged of a leaden hue, pupils dilated, breathing oppressed an laboured, skin cold and clammy, and his pulse slow and full. H was immediately bled in the right arm, but the blood not flowin very freely the vein of the other arm was opened, and a larg quantity of blood was abstracted, when his pulse became smal soft, and more frequent. During the bleeding he had two ver severe attacks of convulsive movements of the right side, wit throbbings of the right corner of the mouth and winking of the ey of that side, the pupils becoming very much contracted. Thes convulsive fits lasted three minutes."

For three hours he remained insensible, after which sinapism fomentations, large doses of calomel, followed by purgatives, wer used, and he recovered. Though but forty-seven years of age h now appeared (September, 1849), prematurely old, and muc emaciated. Hepatic torpor, and torpor of all the functions of th body, accompanied by severe headaches, were now the symptom complained of. He was completely anæmic, and no wonder, for the blood-letting he declared to have been enormous. In the first winter and spring seasons after his arrival at home he suffere severely after meals from sleepiness, uneasiness, and occasions pain in the head, accompanied by sense of weight and oppression of the epigastric region, and fits of coughing. These symptom after four or five hours disappeared; but it was long before he regained proper digestive power, healthy secretions, or muscular condition, sufficient to enable him to take exercise without fatigues.

Remarks.—This officer was seen by many physicians in variou quarters, and various remedies were prescribed. I directed the use of the mineral acids, chalybeates, and latterly of the cod live oil. All three were followed by some improvement; but two years and a half of furlough proved insufficient to restore the vigor of the digestive functions, and he returned to India in a condition which did not promise that he could render effective service.

Case IV.—Major ——, ætatis fifty, served thirty years in India, and had enjoyed an average degree of good health untwithin the last six or seven years, when, according to the statement of his surgeon, he suffered "at periods, varying from a week to month, from severe headaches of a most intense character. The affection commences primarily with a weight and heaviness in the head, which gradually increasing, ceases not until it assumes a degree almost insupportable. During the attack, and until the operation of medicine, he is incapable of moving, and often of lying down, the only relief experienced being in the sitting posture.

From the previous history, present state, and observing the action of medicine on the case, I came to the conclusion that the disordered functions of the brain are entirely caused by torpidity of the liver, and derangement of the digestive organs generally, the retention, in fact, of some of the component parts of the secretion of bile, which are not eliminated from the blood under deficient action of the hepatic apparatus.

In the intervals between the attacks, irregularity of the bowels and extreme torpidity of the liver have always been present, evinced by great dryness of the skin, sallow countenance with loaded conjunctive, occasional pains in the right shoulder, and

heaviness in the hepatic region, &c.

The disposition to torpor, and consequent disturbance in the sensorium, has always been here increased, and brought on, by the slightest change in the seasons or weather, by change of habits from cantonments to camp, and vice versa, while the slightest irregularity of diet, or even a change of the dinner hour, has

always produced its punishment."

Such is the statement of the able surgeon of Major --- 's corps. "The treatment," he says, "was in the first instance, and between the attacks, to restore due actions of the hepatic system, by mercurial alteratives and purgatives, particular attention being paid to the habits and mode of living; but the treatment during the attack has never been altered. On the warning of his headache approaching, pills, containing a large dose of calomel and colocyuth, were immediately taken, followed, some hours after, by an aperient draught of salts and senna, and after copious dark evacuations the headache has generally subsided, leaving him however pallid and depressed in spirits for some days. The action of mercurial purgatives in this case was remarkable, the earlier their operation on the biliary system was affected, the less severe the attack; and if by any delay the medicine was not directly taken the more severe the affection." Seeing that length of residence in India produced "increased severity and frequency" in the disease, and that a return to Europe was deemed "the only means now in his power for the due restoration of the hepatic function," Major - obtained sick leave, and arrived in England in the spring season.

Remarks.—Major —— came under my observation only at intervals. He visited Brighton, Bath, Cheltenham, and other places, and sought various opinions. For the first year he continued exactly as described in India, headache attacking him on an average once a month. The fluid extract of taraxacum with alkalies appeared to me to be of much service to him, and had he persevered in their use his recovery, I think, would have been

more rapid and progressive.

In the treatment of all the disorders and diseases affecting persons who have resided in hot climates, I endeavour to impress or the patient the vast importance of attention to the habits of life—to the diet, to the clothing, especially in the winter and spring seasons, to exercise in a pure air, to the functions of the skin, &c. and that without such care no medical treatment whatever can avail them; in short, that all those means which are preventive of disease are also curative.

Dr. Budd, in his admirable treatise on Diseases of the Liver speaking of the injurious effects of indulgence at table, and of the neglect of exercise, especially in persons of middle age, in whom

respiration has therefore become less active, says :-

"We may often see inverse evidence of those relations in the effect of pure air and active exercise in relieving various disorders that result from repletion and from the retention of principles, which, if burnt out in respiration, should pass off by the liver as bile. Every sportsman must have remarked the effect of a single day's hunting in clearing the complexion. It has no doubt much the same effect on the liver as on the skin."

If the truth and force of these important observations were more generally appreciated in practice, we should have fewer hepatic disorders, and fewer still of those hepatic diseases, which, taken together, render the days of many amongst us, presently as miserable, as they are sure ultimately to shorten them. Exercise in the open air must also prove powerfully beneficial in another manner—viz., by accelerating the actions of respiration, it thus increases the power of the circulation, and thereby promotes a more active return of blood from the liver, spleen, and other important organs, so often engorged and oppressed in the instances of invalids from hot climates.

"Exercise on horseback," says Dr. Billing, "has innumerable advantages over every other kind, as it gives motion to the viscera without fatiguing the limbs; very little motion is given to the

viscera in walking or in carriages."

If it be true that "the portal circulation is carried on in a great measure by the pressure of the abdominal muscles," how great must be the benefits resulting from well regulated exercises in the open air in such cases. The quantity of carbonic acid exhaled from the lungs is proved to be greatly increased under active exercise, and thus we possess another channel of powerful depuration, the active promotion of which should never be neglected. The same result under exercise is found by Lehmann in the depuration by the kidneys; for, while the proportion of uric acid to urea in a state of rest was 1 to 38, after active exercise it fell to 1 in 77. The truth is that the amount of good that may be effected, in the way of prevention and cure of disease, by pure air and

water, proper diet and active exercise, is a matter which up to this day has not received all the attention that is due to it.

In no class of disease is the most rigid adherence to strict rule of diet more necessary than in such as affect the function and structure of the liver; and this truth should be strenuously impressed on the attention of the returned Indian: he should be made to understand that no cure can be effected without it.

Dr. Budd observes justly that by increasing the activity and effect of the respiratory process, that in the vast power we have in modifying them by regulations having reference to the great conditions of air, exercise, temperature, and food, we have means much more effectual than any other, in dealing with biliary disorders.

Patients, in their ignorance of their present condition, and referring only to our heroic measures of cure, under directly opposite circumstances, in India, will sometimes look anxiously to "active treatment," and to speedy and substantive results. They must be made to understand that there are many morbid conditions for which we cannot pretend to find any direct and immediate remedies; and that even in withdrawing things that tend to injury, or to retard the curative operations of Nature, the physician does much;—much more than can be hoped for by random attempts at rapid or "decided" measures of cure, by powerful medicinal agents applied to an exhausted constitution. They must, in short, be induced to look hopefully to the future, and to judge of the power and value of their medical management by the result—by the gradual restoration of health. It is seldom, indeed, that we shall find reasoning such as this disregarded.

### CHRONIC ENLARGEMENT OF THE LIVER.

Next to the liability to periodical fever and bowel complaints, such as dysentery and diarrhea, engendered by residence in malarious countries, stands the liability to disorder and disease of the liver, engendered by antecedent circumstances of climate and of habit of life. It is well known to such as are acquainted with the health history of the British army, that the soldiers employed on the expedition to Walcheren never afterwards proved serviceable in the Spanish campaigns, or when sent to the intertropical colonies.

The least exposure to malaria, to cold and damp, night duties, a shower of rain, or wet feet, was sure to induce fever of an intermittent character. The same latency of disposition, and subsequent liability to fever on the application of slight causes, has all times been remarked, in the East Indies, of Europeans when have once been affected with remittent (jungle) fever.

The chronic enlargement of the liver will generally be found in persons who have suffered from previous acute inflammation, of from repeated congestions of that organ, from fevers, both remitter and intermittent, and from dysentery and diarrhæa. This form of hepatic disease is often associated also with dysentery, acute an chronic, and with diarrhæa; so that, in fact, it may be said to precede, to accompany, and to follow those bowel diseases.

In the East Indies we find that acute hepatic disease stands the Hospital Returns as third in order of frequency, fevers an dysenteries being in the first and second relations. But though this order will be found generally correct, it must be remembere in addition that, in the course of fevers, dysenteries, and diarrhœa hepatic diseases are very apt to arise, and to follow upon then so as, in fact, to render the total amount of cases of hepati disease which occur in the East very great in the aggregate—fa greater than can be exhibited in any Hospital Returns. Out an aggregate European force of 25,431 men of her Majesty's arm serving during ten, eight, and ten years respectively, between 182 and 1826, at the stations of Calcutta, Chinsurah, and Berhampore all in Bengal Proper, there occurred, according to Colonel Tulloci 1,354 cases of original and uncomplicated hepatitis; but if w could to this enumeration add the amount of consecutive hepati disease, the grand total would be very great. Whether th Returns made from the Madras Presidency be more accurate tha those of Bengal, I know not; but we find that out of an aggregat British force of 82,342 men, serving in the former Presidence from 1842 to 1848, there occurred 5,911 cases of liver disease "In India," says Sir James M'Grigor, "the liver seems to be the seat of disease, in nearly the same proportion that the lungs ar in England."

I have, on the other hand, treated numerous cases of hepatienlargement at home, in persons who had never suffered from an tropical disease, nor yet from any functional disorder, or from structural disease of the organ in question, so far as could be ascertained, but who had nevertheless been exposed in India to the general influences explained in the opening chapter of these essays.

The history of these latter instances is very generally as follows. The patient has been enfeebled by many years of residence in he climates:—he comes to England in an impaired state of generally, with torpor of the liver:—he is exposed to cold; an along with this, perhaps, are associated excesses in food and drink

persistence, or to curative effect, it never could have attained to

any repute in the practice of medicine.

N.B. Should the bowels not act with sufficient freedom of themselves, or should the bath not excite them to sufficient action, I am in the habit of ordering a mild aperient draught of Epsom salts in some bitter infusion, every other morning, so long as the use of the bath may be in progress.

CASE I. --, Esq., had served thirty-eight years in India, and there enjoyed excellent health, terminating his official career as member of one of the medical boards. His constitution had been robust; but he was now seventy years of age. He had never suffered from fever, dysentery, or hepatitis, though his habits had always been those of one who is said to "live well." He had retired from the service, and resided in England twelve years, and until eight months previously to his consulting me his health had continued good. At the time now stated he began to fall off in condition, was observed to look sallow, and complained of dyspeptic ailments with fulness in the epigastrium; but there was no pain. Latterly tumidness of the whole abdomen attracted his notice, and that of his family. I found the liver enormously enlarged, and extending down to the umbilicus, accompanied by considerable serous effusion into the abdominal cavity, and by edema of the lower extremities. The bowels were constipated. the excretions being clay-coloured, the urine turbid and scanty. There was no disturbance of the heart's action.

In consultation with Dr. Chambers it was determined to use the nitro-muriatic-acid bath by sponging the limbs and abdomen. using a mild and aperient draught every second morning. This was done by the patient's wife and her female attendant, but after ten days they were obliged to desist, both of them being seized with bilious diarrhea. The person next employed in sponging used gloves of oiled cloth, and the remedy was continued for more than three months, with a powerful effect on the secretions of the liver, bowels, and kidneys. The liver was now not to be felt under the margin of the ribs, and the patient rapidly acquired appetite, condition, and strength. Mr. -- now proceeded to his country residence, returning to London for the winter season, and enjoying excellent health, which he retained for upwards of two years. At the expiration of this time he complained of nausea after meals, accompanied by great languor and distress at the stomach. Vomitings after meals soon succeeded, the matters rejected being latterly of a dark coffee colour, and he died greatly emaciated, three months from the accession of his latter illness, in his seventy-third year.

The contents of the abdomen were examined by Mr. Pollock,

ther ccathe wer, e so mall sure

of a was d of deen

years
liver,
ither
long
ease;
ears—
ln renitroa state
emove
ropsy,
lth reoint of
t have

afferment
estion
layan
her of
to the
in the
sness
happy
I had
g straff
to be

but little advice. A medical man I saw at -, said there was not much the matter with me, and I wanted a change to sea. On reaching Calcutta, Mr. Nicolson said my liver was much enlarged. I was now almost unable to leave my room—almost my bed. Mr. Nicolson gave me, I believe, calomel and opening medicines, and something to operate upon my kidneys. I did not get worse in Calcutta, and was put on board ship. Here I took lots of calomel three times a day, and digitalis. On reaching the Cape of Good Hope I was much better, and a month afterwards very much improved indeed-in fact, I considered myself well. On board ship, Dr. — latterly gave me some sort of acid to rub into my side, but he had not much in store, and I used it sparingly. On reaching England, I consulted -, who attended me for three months, and during that time I grew as bad, or worse than I had been on leaving Calcutta. I had calomel and digitalis. After that, I was treated by Mr. J. R. Martin with acid baths, and occasional medicine at night, and in three months grew very much better, with the constant use of the bath night and morning, and in about six months became a sound man, I hope. My internal organization has worked soundly for about two years, and I embark to-morrow, July 30th, 1846, again for India, without apprehension of a return of my disease. I think I took eight hundred grains of calomel between November, 1842, and August, 1843!"

At the time here last referred to, this gentleman's condition was most unpromising. The abdomen was full of serum, and there were signs of commencing effusion into the cavities of the pleura and pericardium, the heart being slightly enlarged. The liver was of enormous size, and anasarca had extended to the face and eye-

lids.

It was under these circumstances that the nitro-muriatic acid bath was commenced, using daily various diuretics, and a draught composed of compound powder of jalap in aromatic water, every other morning.

The result of this plan of treatment is stated above by the patient. He landed in Calcutta in November 1846, as he wrote to me, "in excellent health." During the voyage by the Cape of Good Hope from England, he added that he had been "remarkably well, and never troubled the doctor or his chest."

Unhappily, Mr. —— remained for months in lower Bengal, where he was seized with remittent fever. The result was ruinous to his general health. The abdominal functions became torpid and depraved; the heart became perceptibly enlarged, with a return of dropsy of the abdominal cavity, and general anasarca. He died in August, 1847—nine months from the date of his return to Bengal.

On examination after death, the heart was found by Dr. —— of Calcutta, to be enlarged, but all the valvular structures were sound. The abdomen "contained an enormous quantity of fluid." The

Professor Bennett, states, that this morbid change in the liver consists of hypertrophy of the fibrous element between the lobules of the organ, and its subsequent contraction whereby its volume is diminished, and the secreting cells compressed and atrophied. As a further result, the large venous trunks are also compressed, and their commencing ramifications so congested that effusion into, or dropsy of the peritonæal cavity is induced.

Dr. Handfield Jones, on the other hand, regards the thickening and condensation of the fibrous tissue of the liver, not so much the effect of inflammatory action, as of a low degenerative process. He states that there are two forms of cirrhosis; "the one depending on thickening of the Glissonian Capsule, and the other, in which the capsules themselves are involved in fibroid degeneration, the abolition of tissue, and of function being much more complete in the latter!"

More than sixty years ago, Dr. Dick, an intelligent physician of Calcutta, noted that, "In chronic cases, where there is no fever, but only an obtuse pain in the side and shoulder, with a fulness in the side and about the pit of the stomach, keeping up a constant uneasiness, mercury seems to me to have but little good effect. When used freely it removes the symptoms for the time, but they generally return as soon as the mercury is left off." Dr. Dick states, also, that "such liver attacks very often succeeded long courses of mercury." Mercury, in the case here last related, as in other similar cases, did no good, but, on the contrary, much harm, though tried in various ways, and for longer and shorter continuance.

Dr. Girdlestone made the same observation in Madras in 1781—83. He ascribes hepatic disease to the abuse of calomel, even when used "as a prophylactic," but not so in any degree to mercurial frictions, which, however long-continued, did not, he thought, produce such results. He also declares that "after the hepatitis is removed, the constant but cautious use of acids, in hot climates, proves the best means of preventing a relapse."

Dr. — had never been in a hot climate, and he stated that his case had originally been one of syphilitic ulceration, treated without mercury. Secondary and tertiary forms of syphilis followed, and for these three protracted courses of mercury were had recourse to, with no benefit. When I saw him his constitution was entirely destroyed, and his liver was of enormous size, having become so after the last mercurial course.

For the following interesting case I am indebted to Mr. Pollock, of St. George's Hospital:—

"27. Grosvenor Street, January 5, 1850.

CASE III.—" I send you the particulars of the case I mentioned to you of scrofulous disease of the hip-joint and enlarged

liver, the recovery from which I attribute solely to nitro-muriatic-acid bath, used as recommended by

In July, 1848, I was requested to see a boy (fifteen years of age, with extensive scrofulous dis The region of the joint was externally marked by cicatrices, and several sinuses still discharged the interior. The thigh was contracted on the body much distorted. His aspect was very unhealthy delicately pallid. It was impossible to ascertain tion of the joint, or position of the head of the consolidation of the surrounding soft parts, and al fering any movement of the parts produced in the was no doubt that considerable mischief had occur and had been going on for some time.

The abdomen was also found to be very larg small quantity of fluid in the cavity of the perit enlargement was chiefly dependent on the great which was very prominent, reached down below ileum, and filled up the greater part of the abdomable to sit up, and moved with difficulty. Pulse v appetite good; tongue clean; but every structure

sanguine condition.

When six years of age, and nine years before he had a fall down stairs. Some months subseque observed him walking lame, and, as they describe tended by a doctor for the disease of the hip." H by degrees, and for the symptoms referred to the issue made in the back part of the thigh, which w some eighteen months. He appeared to improve ment, but soon after an abscess formed in the sic which was opened, allowing the escape of much time until I saw him many abscesses opened an constant discharge continued from the hip, and ocof carious bone came through the openings. months of my seeing him his parents observed th of the abdomen. About a week after I saw him, using the nitro-muriatic-acid foot-bath, and spu with the nitro-muriatic acid, as recommended b first week of its application, the size of the abdor diminished one inch and a half, and during the weeks, one inch every week. After this the pa keep the measure accurately; but during the wh from the commencement of the use of the bath, proved in every respect; he gained strength, colour, and all the sinuses healed up rapidly; and afterwards able to move about upon crutches. Is

April, 1849, so much improved that I did not know him again. Every sinus was closed round the joint, and quite sound; he could place his foot to the ground, and even bear some weight upon it. He had gained flesh, and could walk about with the help of one crutch only, and wished much to have a high-heeled shoe, which, however, I recommended not to be tried too soon. The liver could not be detected larger than natural. The bath was used for some three or four months, being omitted for the interval of a week or a fortnight occasionally; and in the interval he took internally small doses of the nitro-muriatic acid. The entire treatment consisted in this; and to the use of the bath I cannot but attribute the favourable result of the case.

I enclose you a letter from the parents, of which you can make any use you may think proper.

GEORGE D. POLLOCK.

To J. R. Martin, Esq."

Case IV.-Mr. -, a surgeon of eminence, states that, when fourteen years of age, he had an attack of jaundice. In January, 1840, "after being much worried and fatigued," he had a severe bilious attack, with incessant vomiting of bile, followed by jaundice. During a month, his health was not "improved by any sort of treatment;" but on his "mind being powerfully acted on by agreeable intelligence, and stimulated by hope, the jaundice, from that day, began to give way, and in a week it disappeared."

With exception to occasional bilious attacks, one of which was caused by anxiety, he enjoyed good health from this time till March, 1849, when he was "worried about professional matters,

and underwent great fatigue."

On the 29th of April of this year he was seized with rigors, and sense of cold in the back, headache, and tenderness of the eyes, followed by much fever; and while still suffering from anxiety and depression of mind, he had profuse and offensive perspirations From this time till the 18th of October, 1849, when I first saw the patient, the "wave chills," as he calls them, increased, followed by fever, pain of the right hypochondrium and epigastrium. The bowels were constipated, the excretions depraved. For this he took purgatives, and large doses of quinine, which removed the rigors; but fever, followed by profuse and offensive night-sweats, harassed him. The abdomen had for some time been much enlarged, and "a hard illdefined swelling at the epigastrium" was perceived by his friends (16th of September). By this time, great emaciation had taken place; the pulse ranged from 84 to 90; the complexion was "of a dusky yellowish white; the lips were pale and dusky; the tongue covered with white pasty fur; urine, like porter dregs."

softened and flattened, but still nearly of the same extent as before. The tumour of the abdomen had much subsided; the complexion had improved; the urine had become less turbid; and the frequency of the pulse was reduced. The circumference of the body had undergone "a diminution of four and a half inches in less than a month, and this in spite of a considerable increase of fat, for his ribs were now decently covered, and his arm measured an inch more in circumference. The excretions were now nearly natural, with exception of an abundant deposit of lithates in the urine, which was passed in very large quantities. The sleep was good, and the appetite excellent."

Dec. 4th.—Had slight bilious diarrhea. The spleen scarcely

perceptible, and the liver greatly diminished.

December 7th.-Took a seidlitz powder, which was followed

by "a copious discharge of pure, honest, laudable bile."

28th.—"During the last few days I have felt some irritability and depression of spirits; my gums are red and turgid, and bleed on the least suction. Mr. Martin made a careful examination, and finding that the spleen is no longer perceptible, and that the liver has also recovered its natural size, and being of opinion that malaise, and the condition of the gums, indicated that the system was fully under the influence of the nitro-muriatic acid, permitted me to leave off bathing, which has been steadily pursued day after day, with one or two exceptions only, since the 23rd October.

January 5th, 1850.—I now feel in every respect in excellent health; stronger in body and more cheerful than I have been for many months; I may almost say for twelve months. I have gained considerably in bulk; the functions of the alimentary canal are healthily performed; the pulse is firm and natural; the skin soft and healthy; the muscles firm; and I am equal to a great amount of exertion.

I cannot conclude without expressing my conviction that, under Providence, I owe my life to the treatment directed by Mr. Martin. On reviewing my case, it will be seen that there was a steady progress from bad to worse. There were amendments from time to time, but each amendment was succeeded by symptoms more severe and more intractable than had before appeared. It will be further seen that the ordinary treatment, under the direction of most able physicians (whose friendly attention I can never forget), although producing temporary lull, quite failed in removing the cause of the mischief. The state of debility to which I was reduced by the fearful night sweats forbad depletion. The enlarged and tumid spleen put mercury out of the question. Quinine had been tried most fully, and had disappointed us. The mineral acid and taraxacum, though affording some relief at first, soon ceased to produce a beneficial effect, and iodine absolutely disagreed. Thus the line of treatment still open was so circumscribed that the most skilful

mination, you gave as your opinion that the case congestion of the liver; that no suppuration has place, though disorganization was threatened, un be found to emulge this great viscus. Mercury saying you believed it would produce the very were anxious to avert. Captain W --- was ac nitro-muriatic-acid sponging bath twice a day; an the only medicine he was directed to take was so aperient, to maintain a free state of the bowels. rest of the case need require but few words. No used from first to last, but the bath was most ho in, and a few short weeks sufficed to render the st The pale yellow gave place to a more healthy co as the food once more became assimilated, and elaborated. In less than three weeks all pain h on firm pressure. His appetite increased; hi elastic; his limbs were evidently more muscular, diminution in the size and prominence of the t abdomen took place. In six weeks he was w only that additional vigour that change of air gi lescent.

Since the period I have referred to, Captain Voyages to China, and returned in perfect health the last three years, and is now, in China, and only a few days back, from Hong Kong, stabeen uniformly well during this his last reclimates.

In this history I have not dwelt upon any but nent symptoms, nor have I thought it necessary by day progress. The case is mainly valuable exact adaptation of the treatment to the exigen for like as a ship once within the influence of need to trim her sails for weeks perhaps together, ation of treatment was required from the time Co out upon his voyage of recovery until he arrived health."

(Signed)

60, Grosvenor street, Jan 22, 1850."

Case VI.—For the following interesting case Mr. Donald Dalrymple, of Norwich:—

"The Rev. T. C—, aged fifty, the subject tall, bony man, nearly six feet high. His father vigorous, at an advanced age; his mother suffers of gall-stones. He has lost some of his fami He was very active in his habits, a teetotaler by

Mr. — also used the acid bath under grave disadvantages. He commenced this course of treatment in the end of October, and continued it through winter cold of the two following months; and with all this he was in continual and active professional exercise, exposed to weather and other injurious influences; yet, despite such serious disadvantages, the result was complete in curative effect; and he is now, November, 1855, in perfect health.

Case V. - Through the kindness of the late Mr. Dalrymple

the following case has been brought to my recollection:-

"I have great pleasure in giving you an account of the case of hepatic disease about which I consulted you some few years ago; although the relation wants the precision of accurate notes taken at the time.

Captain W., then about thirty-six years of age, had already made several voyages to India and China, and had usually enjoyed excellent health. He was a well-built, muscular man, exceedingly temperate, and of very active, energetic habits. During his last voyage to the East, he had suffered from bilious fever and dysentery, which had been checked, though not cured, by some American physician whom he consulted in his voyage back round Cape Horn, in the Brazils.

On his arrival in London, I was called to see him, and I found him greatly emaciated in his limbs, but protuberant in the abdomen. His complexion was that of a pale lemon, no red blood even in his lips. His bowels were obstinately costive; the evacuations clay-coloured, without a trace of bile. He suffered grievous pain all over the upper part of the abdomen; his strength was gone; and his spirits, in spite of his natural firmness and energy, de-

pressed to the last degree.

On examination of his abdomen, I found the hypochondria expanded, the epigastric region bulging, the liver extending upwards, as determined by percussion, and invading the right thoracic space. The pain produced by even moderate pressure over the hepatic region was exquisite. Under these circumstances, I believed he had acute inflammation of the liver, and that mercury energetically given was indispensable to his recovery. It was fortunate, however, for my patient, that I determined to fortify myself by the advice of another, and before beginning the treatment I asked the opinion of the late Dr. James Johnson. On his first examination of the patient's abdomen, and from observing pressure and percussion produced pain amounting to agony; he told me he feared that abscess of the liver had already formed; but he declined to prescribe, and advised us to consult you, giving me an introduction to yourself for that purpose.

After you had heard our history, and had made a careful exa-

In conclusion, I would add, that during the course of this long and interesting case, no remedy or plan of treatment that could be devised was left untried, though it would be impossible, in the limits of a case, to detail it at length."

The patient reports, on the 30th of Jan., 1850, that his eyes and skin are clear of all yellow tinge, and that his urine is of a healthy appearance, the intestinal secretions having the same character. "The abdominal enlargement, as far as visible to the eye, is quite gone."

"I attribute the improvement in our patient's health," says Mr. Dalrymple, "to the action of the acid bath, and to the emulgent

effect of the croton oil."

Mr. C—, on my recommendation, used the bath again during part of June and July, and he now writes:—"I feel as if I never was in better bodily health," while the weight of the body has increased from nine stone four pounds to about twelve stone.

This gentleman is now, November, 1855, reported to be "stout

and well."

TOF

no

ad

rst

tor

se.

rth

He

int

3

of

ir.

ras

om

d:

ed.

the

an

ler

th

st,

în

Case VII.—April 6th, 1853.—I was requested by Mr. Fergusson, of King's College Hospital, to visit Colonel ———, of the Indian army, an officer of about 65 years of age, having gone to India in 1805. During his course of service, which exceeded thirty-five years of actual residence in India, he had been a keen sportsman, and had consequently been continually exposed to direct solar influence, while his habits were what is called free. He had had three severe attacts of remittent, jungle fever, which were followed by often repeated attacks of intermittent fever. He also suffered from severe rheumatism, for the cure of which he came to England in 1812, returning to his duty in good health, after fourteen months' residence at home. By constitution he was gouty, his mother having died of that disease. He came home again, on sick leave, in 1842, suffering from emaciation and debility, and returned to his duty in India in November, 1846.

While at his presidency, he was nearly destroyed by a large quantity of arsenic, administered by one of his servants, with a view to murder and robbery. From this poisoning he recovered rapidly, and returned to England finally, in March, 1847. From this time he describes his health as always indifferent; but it has been more especially so, during the last three years. In India, from 1835, he had occasional slight attacks of gout, but in England they have become more severe, and more frequent of occurrence, especially since exposure to wet and cold, while shooting in the

Highlands of Scotland two years ago.

In January, 1853, Colonel ———, became seriously indisposed, and confined altogether to his house. His skin became of a lemon

l, and by a eased. arrowretion quarynding ssure, ongue n, the t and anght a was ibiteci neans l; but iolent h was owels from ion of come from ithous erient ses of wentybdned. on on re flow of moanied

n bile, food.

and

L By

appe-

d with tinued s confarred Ordered a repetition of the bath every night, and a magnesian draught in the morning.

June 7th.—Two dark green motions followed by much relief from pain. Urine examined carefully, and found nothing unnatural. June 8th.—Much better. Had three dark green motions, followed by much relief from pain. Urine abundant and straw-

coloured. To continue the bath at bed-time only.

Towards the end of June, Colonel —— had so far recovered in condition and strength as to take regular and daily active exercise, and he then went into the country. I saw no more of him till the 10th September. He stated that for some three weeks previously, the epigastric pain had returned, with frequent vomiting, and some tendency to diarrhæa. All food, even the most bland, caused excessive pain. He had again fallen off in condition; the pulse was frequent, and the countenance had become extremely pallid.

Sept. 12th.—Visited Colonel —— in company with Mr. Pollock. The patient was now much exhausted from constant vomiting and pain, the matter vomited being streaked with blood, as were the contents of the bowels. Having left London at this time, I saw no

more of the patient, who died on the 11th October, 1853.

For the following report of the post-mortem examination by Dr. Quain, I am indebted to Mr. Fergusson:—

"Stomach.—There was no disease of this organ, save some thickening of the coats over the pylorus. The mucous membrane was

everywhere sound.

Liver.—The whole of this organ was throughout paler than natural—the tissue was tolerably firm—towards the anterior margin there was some contraction, rounding and puckering, giving it to a limited extent, the appearance of cirrhosis.

Spleen .- This organ was small, firm, and traversed by an

unusual number of fibres, or trabeculæ.

Kidneys.—These organs appeared healthy.

The small intestines were healthy.

Large Intestine.—This bowel appeared elongated—much convoluted—distended at some points—contracted at others. On laying the bowel open, it was found filled with a chocolate coloured viscid substance, about the consistence of treacle. This substance was bounded above by the ileo-cœcal valve, and occupied the bowel to the anus. On washing the surface of the intestine, the mucous membrane was seen to be of a deep chocolate colour—the follicular structure of the cœcum was particularly well marked—the membrane in some parts was extremely thin—in others presenting marked vascularity. There were seen also the perfect cicatrices of some old ulcers."

vellow, hard and dry to the touch. The pulse became rapid he had no natural sleep; the appetite failed, accompanied most violent "burning pain" in the epigastric region, incre even on swallowing the most bland food, as milk, sago, or a root. The urine was scanty and blood-red, the alvine exce being clay-coloured. He had greatly emaciated since Jan The liver was, on April 6th, found greatly enlarged, exter down to near the umbilious, hard, and very painful on pres especially in the epigastric region:-pulse one hundred; to much coated. Ordered, in consultation with Mr. Fergusson nitro-muriatic acid bath for the whole body, to be used night morning, followed every other morning by a mild aperient dra To secure sleep, a mild anodyne of solution of morphia used every night, while the dilute nitro-muriatic acid was exhi internally three times a day, in bitter infusion. By this n the fever, pain, and restlessness were in some degree relieved on the fifth day, the treatment was discontinued, owing to a vi gouty inflammation in the knee-joint: on April 25th, the bath resumed, and about May 12th, its action on the liver and be became perceptible, in two or three large bilious motions the bowels in the twenty-four hours, and in a copious excretistraw-coloured urine. By this time also the pulse had be natural, the tongue clean, the skin perspirable, and free jaundice; and food of every kind was borne, not only wi pain, but it was taken with appetite. The anodyne and ape draughts were now discontinued, as were the internal dos dilute acid; and the bath was used only once in the two four hours, viz. at bed-time. The fever had now been subo and there was but slight uneasiness in the hepatic regio pressure, none in the epigastrium. After a few more days the of green and yellow bile became more copious, the number of tions averaging from five to six in the twenty-four hours, accompa by slight malaise, with redness and tenderness of the gums tongue, so that on May 23rd, all treatment was discontinued. this date, Colonel — had used thirty-seven baths.

May 26th.—Owing to the frequent passage of acrid green he felt griped and uneasy, and had but little inclination for Ordered, bicarbonate of soda in bitter infusion, and a magne aperient in the morning.

28th.—Draught acted twice, and he now feels much better, a

tite returning; has improved in condition.

June 6th.—Has continued improving in all ways until towhen owing to overloading of the stomach, he was seized epigastric pains of a severe character: the bowels have contiopen twice or thrice a day, without medicine, the discharges taining a greenish-yellow bile; urine high coloured, tongue fu during two years by occasional spasms in the pit of the stomach. In October, 1854, I came down to Calcutta for medical advice, when a great enlargement of the liver was discovered, the jaundice having by this time become intense, the spasms recurring monthly, and the complexion being a deep green and yellow. Treatment having failed of doing me good, I embarked, by the advice of my medical attendant, for England in January, 1855. During the voyage round the Cape I had partly the assistance of the surgeon of the vessel, but without any avail, and in April of the same year I arrived in England, and on the 26th of that month consulted Mr. J. R. Martin of Grosvenor Street, and formerly of Calcutta. At this time I was emaciated and enfeebled to the last degree; my stomach was large and tumid; my urine was like porter, thick and dark, and the motions from my bowels were white. I was instantly put under a course of acid baths which I continued without intermission for four months. I am now in a state of feeling of comfort and health, such as I have not experienced for a couple of years; I eat with an appetite and sleep well; my complexion is very clear, and the urine apparently healthy looking. The liver, my medical adviser in the country says, is not now to be felt, and my strength is rapidly returning, so much so that I can walk six and seven miles with ease. Formerly, my friends and medical attendant, who had not seen me for some years, did not recognize me, so emaciated and altered had I become in every way; my sister, whom I had not met for eight months, received me as a stranger; and even my wife declared she would have passed me in the street without recognition. When I first put myself into Mr. Martin's hands, which was at the end of April last as above stated, I weighed only nine stone (my original weight in health having been thirteen stone ten pounds), and in August last, when I again weighed myself, I weighed ten stone. I now weigh about eleven stone."

November 15th, 1855.

## CONCLUSION.

It will be seen that in the management of convalescence from acute diseases, and for the treatment of those of a chronic kind, I have seldom had recourse to mercury as a therapeutic agent; believing that the powerful actions of this mineral are most appropriately applied to acute and dangerous maladies. To use mercurials perseveringly for disorders of the health, or for the cure of chronic affections, is to confound that which is appropriate to

ed would es of iml anguard e was more favour.
The man anguard the en commend on th

se worthy

inder the

lusion:

-L

is aspect diced. I adergone is, nitrolost four ne clear iver, the e tongue ction, the ith bile."

1855."

India, in d. The

became

a constitution of the cons

Chirurgical issued, Dr. ercurials in f that merof recently a remedies, g in cases of ment by the preventing leposited, its that of meri- and endo-

nd I quote it forth in page we shall be enabled effectually to cure acute tropical and other diseases, without the assistance of mercury; but it has not as yet arrived. In the mean time, and in the absence of a better knowledge, what we should try to obviate is—not the use of mercury, but its abuse.

Of the sufferers who have passed through the storms of tropical disease described in this work, it will be seen that many recover immediately on the spot, a certain proportion sinks, while others come to their native country for the restoration and confirmation of health. In the cases of these last, it is evident that a careful consideration of all their antecedent circumstances is absolutely necessary to their just management, and that we must understand the nature of their acute diseases in India, and their treatment there, to appreciate aright their chronic ailments on their return to Europe. Such a review will, I think, generally satisfy the medical observer at home, that with mercury in any form his emaciated, relaxed, enfeebled, and generally anæmic tropical patient need now have but little to do.

I have seen more mercury administered in this country, for what may be termed the disorders of health in adults, and more "gray powder" given persistently to children, by female relatives, governesses, and nurses, than is ever seen in India; and in many instances the results have been more lasting and lamentable than anything I witnessed in the East. It ought to be made a rule in families that no preparation of mercury should be used, excepting under competent and careful medical advice; and I feel assured that the carrying out so salutary a domestic reform would confer a greater benefit on young women especially, than all those lighter accomplishments with the acquisition of which their earlier years

are teazed.

These circumstances I think it right to mention; and certainly I do so in no arrogant controversial spirit, or with a view to finding fault.

Angry criticisms and unseemly wranglings, on the subject of the efficacy, or otherwise, of certain remedies, and on the merits of certain kinds of practices, I have ever seen to be prejudicial at once to the character of the profession of medicine, and to in-

dividual credit and reputation for usefulness.


If we cannot carry our point by justice, truth, and reason, we must wait till they come to our aid, as eventually they are sure to do. Cabanis said, more than half a century ago:—"Let us unite our efforts, and endeavour to introduce into the study and practice of our art that superior reason and philosophy, without which so far is it from affording useful aid, that it becomes a real public scourge."

acute, with what is applicable to protracted forms of organic lesion. In this our climate a sparing application of the means in question, for a short time, may be justly made where congestion threatens greatly to disturb the function, or to affect the structure of the liver; but I know nothing so greatly to be reprehended as the indiscriminate or inconsiderate use of so powerful a class of remedies.

In India, whether we have to treat an European suffering from remittent fever, dysentery, hepatitis, or cholera, the symptoms and progress of disease are so acutely violent as to leave but a very few days between recovery or death: the means of cure, therefore. to be effective must be energetic, that is, if we would save life, and prevent the occurrence of dangerous organic diseases. In fever, dysentery, and hepatitis, the patient, as a first measure, is bled to an extent proportioned to the severity of the disease, reference being had to age, constitution, habits of life, and length of residence in India; and blood-letting is repeated, generally, or locally, or in both forms, until a sufficient abatement is obtained in the force and frequency of the actions of the heart and arteries. The disease still proceeding with dangerous force and rapidity, we have next to employ calomel in aid of blood-letting, followed by purgatives, sudorifics, and diuretics, so as to excite all the depurative functions, and thus bring about a solution of acute and dangerous disease. These means we gradually abate as symptoms yield, and generally no injury is done to the patient's constitution, convalescence being rapid; but they are heroic remedies nevertheless. which demand much knowledge, discernment, and experience, for their just application, and even for their withdrawal. In the present state of our knowledge we possess no other means of cure than those here briefly described, aided by antiperiodics and tonics.

Under the sanction of the Royal Medical and Chirurgical Society of London, in its Thirty-eighth Volume, just issued, Dr. Garrod, in contrasting the actions of alkalies and mercurials in the treatment of acute rheumatism, expresses his belief that mercurials "have more power in causing absorption of recently deposited lymph, than alkalies or any other known remedies. This point he has recently had opportunities of testing in cases of pleuritis, in which it appeared that the alkaline treatment by the bicarbonate of potash, possessed great influence in preventing deposition, but that when the lymph was already deposited, its value in causing its absorption was far inferior to that of mercurials; he believes that the same holds good in peri- and endocarditis of a rheumatic character."

I believe this statement to be essentially true; and I quote it here as being altogether confirmatory of what is set forth in page 267 of the present work. The day may come when we shall be



ressord to d to mile

EW:

ICAL

PRCSE

TE.

d to all

and Scientific insure four SE a

THE

Iedical Warks iteral Sciences nd. Post 3 va

DICINE

ed by Forman eries.

IAL.

e Empire.

RECTORY

## A CLASSIFIED INDEX

## MR. CHURCHILL'S CATALOGUE.

DENTISTRY.

ANATOMY.	
	CE
Anatomical Remembrancer	3
Denie on Liver	5
Hassall's Micros. Anatomy	15
Holden's Human Osteology	15
Jones' and Sieveking's Patho-	
logical Anatomy	17
Maclise's Surgical Anatomy Paget's Catalogue	19
Paget's Catalogue	21
Sibson's Medical Anatomy	25
Toynbee's Catalogue	28
Wheeler's Handbook	30
Wilson's Anatomy	31
********	
CHEMISTRY.	
Abel & Bloxam's Handbook	4
Bowman's Practical Chemistry	7
Do. Medical do	7
City of the second Williams City on Taken	8
Fownes' Manual of Chemistry	12
Do. Actonian Prize	12
	12
The Chamberland Making	12
Fresenius' Chemical Analysis	
Calleman's Chemical Analysis	12
Galloway's First Step	12
Do. Analysis	12
	12
Grimths Four Seasons	13
Horsley's Chem. Philosophy Jones.—Mulder on Wine	16
Jones.—Mulder on Wine	17
Odling's Practical Chemistry	21
Plattner on Blowpipe	22
Speer's Pathol Chemistry	26
*********	
CHOLERA.	
Acland on Cholera at Oxford	9
Role and Call Percets	4
Baly and Gull—Reports Snow on Cholera	115
Show on Cholera	25
CLIMATE.	
Francis on Change of Climate	12
Hall on Torquay	
Hall on Torquay	14
Lee on Climate	18
Martin on the Undereliff	19
Martin (J. R.) on Tropical	20
marin for my on andhom	-

DEFORMITIES, &c.

Bigg on Deformities
Bishop on Deformities
Do. Articulate Sounds
Brodhurst on Spine
Do. on Clubfoot
Hare on Spine
Hugmaa on Hip Joint
Inman on Spine
Tamplin on Spine

St Killer - OL

SECTION - 08

Odontological Soc. Transactions 21					
DISEASES of the URINARY					
and GENERATIVE ORGANS,					
and SYPHILIS.					
Acton on Reproductive Organs 3					
Do. Urinary and Genera-					
tive Organs 3					
Coote on Syphilis 9					
Coulson on Bladder 10					
Do. on Lithotomy 10					
Egan on Syphilis 11					
Judd on Syphilis 17					
Milton on Gonorrhœa 20					
Parker on Syphilis 22					
Todd on Urinary Organs 28					
Wilson on Syphilis 31					
DISEASES OF WOMEN					
AND CHILDREN.					
Bennet on Uterus 5					
Do. on Uterine Pathology 5					
Bird on Children 6					
Brown on Women					
Eyre's Practical Remarks 11					

DISEASES OF WOMEN	
AND CHILDREN.	
Bennet on Uterus	5.
Do. on Uterine Pathology	
Bird on Children	6 7 7
Brown on Women	7
Do. on Scarlatina	7
Eyre's Practical Remarks	11
lood on Crowing	16
ee's Ovarian and Uterine Dis-	31
eases	18
Lee on Diseases of Uterus	
	18
Roberton on Women	
Rowe on Females	
smith on Leucorrhea	
Tilt on Diseases of Women	
Do. on Change of Life	
Inderwood on Children	
West on Women	
Vhitehead on Abortion	30
NAMES AND ADDRESS OF THE PARTY	
HYGIENE.	
Seale's Laws of Health	4
Do. Health and Diseases	4
Slundell's Medicina Mechanica	6
arter on Training	4 6 8
Cornaro on Long Life	9
lartwig on Sea Bathing	
Do. Physical Education	14

	Tilt on Diseases of Women	27
	Do. on Change of Life	27
		26
		25
	Whitehead on Abortion	
ı	MANAGE STATE OF THE STATE OF TH	
ı	HYGIENE.	
١	Beale's Laws of Health	
١	Do, Health and Diseases	
ı	Blundell's Medicina Mechanica	к
	Carter on Training	
ı	Cornaro on Long Life	в
١	Hartwig on Sea Bathing	1
	Do. Physical Education	1
	Hufeland's Art	i
	Lee's Watering Places of England	
	Do. do. Germany,	
	France, and Switzerland	1
	Lee's Rhenish Watering Places	ï
	Pickford on Hygiene	2
		2
		2
	Rumsey's State Medicine	20
	Van Oven's Decline of Life	2
	Wilson on Healthy Skin	3
	Do. on Mineral Waters	3

### MATERIA MEDICA and PHARMACY.

DENTISTRY.	MATERIA MEDICA SIIG
Blundell's Painless Extraction . 7	PHARMACY.
	PAGE
Clark's Odontalgist, 9	Bateman's Magnacopia 4
Gray on the Teeth 13	Beasley's Formulary 5
Odontological Soc. Transactions 21	Do. Receipt Book 5
WALKARY.	
	To Transfer of transferons
DISEASES of the URINARY	Lane's Materia Medica 17
and GENERATIVE ORGANS,	Pereira's Selecta e Præscriptis 22
	Pharmacopœia Londinensis 22
and SYPHILIS.	Prescriber's Pharmacopœia 23
Acton on Reproductive Organs 3	
	Royle's Materia Medica 24
Do. Urinary and Genera-	Spurgin's Materia Medica 26
tive Organs 3	Squire's Pharmacopœia 26
Coote on Syphilis 9	Steggall's Materia Medica 26
Coulson on Bladder 10	Do. First Lines for Chemists 26
Do. on Lithotomy 10	Stowe's Toxicological Chart 27
Egan on Syphilis 11	Taylor on Poisons 27
Judd on Syphilis 17	Wittstein's Pharmacy 31

MEDICINE.	
Adams on Rheumatic Gout	4
Addison on Cells	1
Alexander on Rheumatism	1
Arnott on a Local Anæsthenic	3
Barclay on Diagnosis	1
Barlow's Practice of Medicine	4
Billing's First Principles	1
Bird on Charcoal	4
Brinton on Ulcer	1
Budd on the Liver	1
Do. on Stomach	3
Chambers on Digestion	4
Davey's Ganglionic	11
Eyre on Stomach	11
Fuller on Rheumatism	15
Gairdner on Gout	E
Garrett on E. and N. E. Winds	13
Granville on Sudden Death	1
Gully's Simple Treatment	1
Habershon on Stomach	1
Hall on Apnœa	14
Hall's Observations	1
Harrison on Lead in Water	1
Headland on Medicines	1
Hooper's Medical Dictionary	1
Hooper's Physician's Vade-	
Mecum	1
Lugol on Scrofula	11
Peacock on Influenza	2
Do. on Heart	25
Pym on Yellow Fever	2
Robertson on Gout	2
Savory's Compendium	2
Semple on Cough	2
Shaw's Remembrancer	24
Steggall's Medical Manual	2
Do. Gregory's Conspectus	2
Do. Celsus	2

Do. Celsus.
Thomas' Practice of Physic
Wegg's Observations.
Wells on Gout
What to Observe
Whitehead on Transmission
Williams' Principles
Wright on Headaches

DR. ACLAND.

MEMOIR ON THE CHOLERA AT OXFORD IN THE YEAR 1854; with Considerations suggested by the Epidemic. 4to. cloth, with Maps, 12s.

## MR. ACTON, M.R.C.S.

THE FUNCTIONS AND DISORDERS OF THE REPRODUC-TIVE ORGANS IN YOUTH, IN ADULT AGE, AND IN ADVANCED LIFE. Considered in their Physiological, Social, and Psychological Relations. Second Edition. 8vo. cloth, 7s.

IL.

PROSTITUTION: Considered in its Moral, Social, and Sanitary Bearings, with a View to its Amelioration and Regulation. 8vo. cloth, 10s. 6d.

III.

A PRACTICAL TREATISE ON DISEASES OF THE URINARY AND GENERATIVE ORGANS OF BOTH SEXES, INCLUDING SYPHILIS. Second Edition. 8vo. cloth, 20s.

DR. WILLIAM ADDISON, F.R.S.

CELL THERAPEUTICS. 8vo. cloth, 4s.

II.

ON HEALTHY AND DISEASED STRUCTURE, AND THE TRUE
PRINCIPLES OF TREATMENT FOR THE CURE OF DISEASE, ESPECIALLY CONSUMPTION
AND SCROPULA, founded on MICROSCOPICAL ANALYSIS. 8vo. cloth, 12s.

DR. ALEXANDER, F.R.C.S. BY EXAM.

RHEUMATISM: its Nature, Causes, and Cure. GOUT: its Nature, Causes, Cure, and Prevention.
Post 8vo. cloth. 7s. 6d.

MR. ANDERSON, F.R.C.S.

THE CAUSES, SYMPTOMS, & TREATMENT OF ECCENTRIC NERVOUS AFFECTIONS. 8vo. cloth, 5s.

THE ANATOMICAL REMEMBRANCER; OR, COMPLETE POCKET ANATOMIST. Fifth Edition, carefully Revised. 32mo. cloth, 3s. 6d.


## DR. JAMES ARNOTT.

ON THE REMEDIAL AGENCY OF A LOCAL ANÆSTHENIC OR BENUMBING TEMPERATURE, in various painful and inflammatory Diseases. 8vo. cloth, 4s. 6d.

ON INDIGESTION; its Pathology and its Treatment, by the Local Application of Uniform and Continuous Heat and Moisture. With an Account of an improved Mode of applying Heat and Moisture in Irritative and Inflammatory Diseases. With a Plate. 8vo. 5s.

PRACTICAL ILLUSTRATIONS OF THE TREATMENT OF OBSTRUCTIONS IN THE URETHRA, AND OTHER CANALS, BY THE DILATATION OF FLUID PRESSURE. 8vo. boards, 3s.

a 2



(18c-01

MR. HENRY HEATHER BIGG.

THE MECHANICAL APPLIANCES NECESSARY FOR THE TREATMENT OF DEFORMITIES. Post 8vo. cloth, 4s.

ARTIFICIAL LIMBS; THEIR CONSTRUCTION AND APPLI-CATION. With Engravings on Wood. 8vo. cloth, 3s.

PRACTICAL TREATISE ON THE DISEASES OF CHILDREN AND INFANTS AT THE BREAST. Translated from the French of M. BOUCHUT, with Notes and Additions. 8vo. cloth. 20s.

# DR. GOLDING BIRD, F.R.S.

URINARY DEPOSITS; THEIR DIAGNOSIS, PATHOLOGY, AND THERAPEUTICAL INDICATIONS. With Engravings on Wood. Fifth Edition. Post 8vo. cloth, 10s. 6d.

ELEMENTS OF NATURAL PHILOSOPHY; being an Experimental Introduction to the Study of the Physical Sciences. Illustrated with numerous Engravings on Wood. Fourth Edition. By Golding Bird, M.D., F.R.S., and Charles Brooke, M.B. Cantab., F.R.S. Feap. 8vo. cloth, 12s. 6d.

#### MR. JAMES BIRD.

VEGETABLE CHARCOAL: its MEDICINAL and ECONOMIC PRO-PERTIES; with Practical Remarks on its Use in Chronic Affections of the Stomach and Bowels. Second Edition, 8vo. cloth, 3s. 6d.

### MR. BISHOP, F.R.S.

ON DEFORMITIES OF THE HUMAN BODY, their Pathology and Treatment. With Engravings on Wood. 8vo. cloth, 10s.

ON ARTICULATE SOUNDS, AND ON THE CAUSES AND CURE OF IMPEDIMENTS OF SPEECH. 8vo. cloth, 4s.

LETTSOMIAN LECTURES ON THE PHYSICAL CONSTI-TUTION, DISEASES AND FRACTURES OF BONES. Post 8vo., 2s. 6d.

PRACTICAL OBSERVATIONS ON CERTAIN DISEASES OF THE CHEST; and on the Principles of Auscultation. 8vo. cloth, 12s.


## DR. JOHN W. F. BLUNDELL.

MEDICINA MECHANICA; or, the Theory and Practice of Active and Passive Exercises and Manipulations in the Cure of Chronic Disease. Post 8vo. cloth, 6s.

4

Care - oi

10 -- DE



## MR. CHURCHILL'S PUBLICATIONS.

MR. WALTER BLUNDELL.

## PAINLESS TOOTH-EXTRACTION WITHOUT CHLOROFORM;

with Observations on Local Anæsthesia by Congelation in General Surgery. Second Edition, 2s. 6d, cloth. Illustrated on Wood and Stone.

#### MR. JOHN E. BOWMAN.

PRACTICAL CHEMISTRY, including Analysis. With numerous Illus-

trations on Wood. Third Edition. Foolscap 8vo. cloth, 6s. 6d.

MEDICAL CHEMISTRY; with Illustrations on Wood. Third Edition. Fcap. 8vo. cloth, 6s. 6d.

DR. BRINTON.

THE SYMPTOMS, PATHOLOGY, AND TREATMENT OF ULCER OF THE STOMACH. Post 8vo. cloth, 5s.

DR. JAMES BRIGHT.

ON DISEASES OF THE CHEST AND AIR PASSAGES; with a Review of the several Climates recommended in these Affections. Second Edition. Post 8vo. cloth, 7s. 6d.

MR. ISAAC BAKER BROWN, F.R.C.S.

ON SOME DISEASES OF WOMEN ADMITTING OF SUR-GICAL TREATMENT. With Plates, 8vo. cloth, 10s. 6d.

ON SCARLATINA: its Nature and Treatment. Second Edition. Fcap. 8vo. cloth, 3s.

## MR. BERNARD E. BRODHURST.

ON LATERAL CURVATURE OF THE SPINE: its Pathology and Treatment. Post 8vo. cloth, with Plates, 3s.

ON THE NATURE AND TREATMENT OF CLUBFOOT AND ANALOGOUS DISTORTIONS involving the TIBIO-TARSAL ARTICULATION. With Engravings on Wood. 8vo. cloth, 4s. 6d.

DR. JOHN CHARLES BUCKNILL, & DR. DANIEL H. TUKE.

A MANUAL OF PSYCHOLOGICAL MEDICINE: containing the History, Nosology, Description, Statistics, Diagnosis, Pathology, and Treatment of Insanity. 8vo. cloth, 15s.

### DR. BUDD, F.R.S.

## ON DISEASES OF THE LIVER.

Illustrated with Coloured Plates and Engravings on Wood. Third Edition. 8vo. cloth, 16s.

ON THE ORGANIC DISEASES AND FUNCTIONAL ORDERS OF THE STOMACH. 8vo. cloth, 9s.

DR. BURGESS.

THE MEDICAL AND LEGAL RELATIONS OF MADNESS; showing a Cellular Theory of Mind, and of Nerve Force, and also of Vegetative Vital Force. 8vo, cloth, 7s.

## DR. BURNETT.

THE PHILOSOPHY OF SPIRITS IN RELATION TO MATTER,
8vo. cloth, 9s.
INSANITY TESTED BY SCIENCE. 8vo. cloth, 5s.

DR BRYCE.

ENGLAND AND FRANCE BEFORE SEBASTOPOL, looked at from a Medical Point of View. 8vo. cloth, 6s.

#### MR. ROBERT B. CARTER, M.R.C.S.

B. CARTER, W.R.O.S.

ON THE INFLUENCE OF EDUCATION AND TRAINING IN PREVENTING DISEASES OF THE NERVOUS SYSTEM. Feap. 8vo., 6s.

THE PATHOLOGY AND TREATMENT OF HYSTERIA. Post 8vo, cloth, 4s. 6d.

#### DR. CARPENTER, F.R.S.

PRINCIPLES OF HUMAN PHYSIOLOGY. With numerous Illustrations on Steel and Wood. Fifth Edition. 8vo. cloth, 26s.

PRINCIPLES OF COMPARATIVE PHYSIOLOGY. Illustrated with 300 Engravings on Wood. Fourth Edition. 8vo. cloth, 24s.

A MANUAL OF PHYSIOLOGY. With numerous Illustrations on Steel and Wood. Third Edition. Fcap. 8vo. cloth, 12s. 6d.

THE MICROSCOPE AND ITS REVELATIONS. With numerous Engravings on Wood. Second Edition. Fcap. 8vo. cloth, 12s. 6d.

#### MR. CHARLES CHALMERS.

ELECTRO-CHEMISTRY, with Positive Results; and Notes for Inquiry on the Sciences of Geology and Astronomy: with a Tract of Miscellanies. 8vo. cloth, 3s. 6d.

#### DR. CHAMBERS.

DIGESTION AND ITS DERANGEMENTS. Post 8vo. cloth, 10s. 6d.

DR. G. C. CHILD.

ON INDIGESTION, AND CERTAIN BILIOUS DISORDERS OFTEN CONJOINED WITH IT. Second Edition. 8vo. cloth, 6s.

(1900 - DI-

10-->4

th, 5s.

Illustrations

otes for Inqui llanies 8vo

o, cloth, 1 Os.

6000-+01

## MR. CHURCHILL'S PUBLICATIONS.

MR. H. T. CHAPMAN, F.R.C.S.

THE TREATMENT OF OBSTINATE ULCERS AND CUTA-NEOUS ERUPTIONS OF THE LEG WITHOUT CONFINEMENT. Second Edition. Post 8vo. cloth, 3s. 6d.

VARICOSE VEINS: their Nature, Consequences, and Treatment, Palliative and Curative. Post 8vo. cloth, 3s. 6d.

### MR. J. PATERSON CLARK, M.A.

THE ODONTALGIST: OR, HOW TO PRESERVE THE TEETH, CURE TOOTHACHE, AND REGULATE DENTITION FROM INFANCY TO AGE. With plates. Post 8vo. cloth, 5s.

#### DR. CONOLLY.

THE CONSTRUCTION AND GOVERNMENT OF LUNATIC ASYLUMS AND HOSPITALS FOR THE INSANE. With Plans. Post 8vo.

### LEWIS CORNARO.

SURE METHODS OF ATTAINING A LONG AND HEALTH-FUL LIFE. Thirty-eighth Edition. 18mo., 1s.

#### MR. COOLEY.

COMPREHENSIVE SUPPLEMENT TO THE PHARMACOPCEIAS.

THE CYCLOPÆDIA OF PRACTICAL RECEIPTS, AND COLLATERAL INFORMATION IN THE ARTS, PROFESSIONS, MANUFACTURES, AND TRADES, INCLUDING MEDICINE, PHARMACY, AND DOMESTIC ECONOMY; designed as a Compendious Book of Reference for the Manufacturer, Tradesman, Amateur, and Heads of Families. Third and greatly enlarged Edition, 8vo. cloth, 26s.

MR. BRANSBY B. COOPER, F.R.S.

LECTURES ON THE PRINCIPLES AND PRACTICE OF SUR-GERY. 8vo. cloth, 21s.

MR. W. WHITE COOPER.

ON NEAR SIGHT, AGED SIGHT, IMPAIRED VISION, AND THE MEANS OF ASSISTING SIGHT. With 31 Illustrations on Wood. Second Edition. Fcap. 8vo. cloth, 7s. 6d.

## MR. COOPER.

A DICTIONARY OF PRACTICAL SURGERY; comprehending all the most interesting Improvements, from the Earliest Times down to the Present Period. Seventh Edition. One very thick volume, 8vo., 1l. 10s.

MR. HOLMES COOTE, F.R.C.S.

A REPORT ON SOME IMPORTANT POINTS TREATMENT OF SYPHILIS. 8vo. cloth, 5s.

SE 6800-01

SIR ASTLEY COOPER, BART., F.R.S.

ON THE STRUCTURE AND DISEASES OF THE TESTIS.

Illustrated with 24 highly finished Coloured Plates. Second Edition. Royal 4to.

Reduced from £3. 3s. to £1. 10s.

### DR. COTTLE.

A MANUAL OF HUMAN PHYSIOLOGY FOR STUDENTS; being a Condensation of the Subject, a Conservation of the Matter, and a Record of Facts and Principles up to the present Day. Fcap. 8vo., 5s.

### DR. COTTON.

ON CONSUMPTION: Its Nature, Symptoms, and Treatment. To which Essay was awarded the Fothergillian Gold Medal of the Medical Society of London. Second Edition. 8vo. cloth, 8s.

PHTHISIS AND THE STETHOSCOPE: a concise Practical Guide to the Physical Diagnosis of Consumption. Foolscap 8vo. cloth, 3s. 6d.

# MR. COULSON.

ON DISEASES OF THE BLADDER AND PROSTATE GLAND.
The Fifth Edition, revised and enlarged. 8vo. cloth, 10s. 6d.

ON LITHOTRITY AND LITHOTOMY; with Engravings on Wood.

ON DISEASES OF THE JOINTS. 8vo. In the Press.

### MR. CURLING, F.R.S.

OBSERVATIONS ON DISEASES OF THE RECTUM. Second

A PRACTICAL TREATISE ON DISEASES OF THE TESTIS, SPERMATIC CORD, AND SCROTUM. Second Edition, with Additions. 8vo. cloth, 14s.

MR. JOHN DALRYMPLE, F.R.S., F.R.C.S.

PATHOLOGY OF THE HUMAN EYE. Complete in Nine Fasciculi: imperial 4to., 20s. each; half-bound morocco, gilt tops, 9l. 15s.

### DR. DAVEY.

THE GANGLIONIC NERVOUS SYSTEM: its Structure, Functions, and Diseases. 8vo. cloth, 9s.

ON THE NATURE AND PROXIMATE CAUSE OF IN-

600 ---

· --- ----

11

E TESTIS.

TUDENTS;

eatment. To

ractical Guide

TE GLAND

ings on Wood

Press.

UM. Second

IE TESTIS

Nine Fasci

are, Functions

E OF

DR. HERBERT DAVIES.

ON THE PHYSICAL DIAGNOSIS OF DISEASES OF THE LUNGS AND HEART. Second Edition. Post 8vo. cloth, 8s.

MR. DIXON.

A GUIDE TO THE PRACTICAL STUDY OF DISEASES OF THE EYE. Post 8vo. cloth, 8s. 6d.

DR. TOOGOOD DOWNING.

NEURALGIA: its various Forms, Pathology, and Treatment. THE JACKSONIAN PRIZE ESSAY FOR 1850. 8vo, cloth, 10s. 6d.

DR. DRUITT, F.R.O.S.

THE SURGEON'S VADE-MECUM; with numerous Engravings on Wood. Seventh Edition. Foolscap 8vo. cloth 12s. 6d.

MR. DUNN, F.R.C.S.

AN ESSAY ON PHYSIOLOGICAL PSYCHOLOGY. 8vo. cloth, 4s.

DR. JOHN C. EGAN.

SYPHILITIC DISEASES: THEIR PATHOLOGY, DIAGNOSIS,
AND TREATMENT: including Experimental Researches on Inoculation, as a Differential Agent in Testing the Character of these Affections. 8vo. cloth, 9s.

SIR JAMES EYRE, M.D.

THE STOMACH AND ITS DIFFICULTIES. Fourth Edition. Fcap. 8vo. cloth, 2s. 6d.

PRACTICAL REMARKS ON SOME EXHAUSTING DIS-EASES. Second Edition. Post 8vo. cloth, 4s. 6d.

DR. FELL.

A TREATISE ON CANCER AND ITS TREATMENT. 8vo.

This work embraces an account of the Remedies, external and internal, employed by the Author; the mode of their Preparation and Application.

DR. FENWICK.

ON SCROFULA AND CONSUMPTION. Clergyman's Sore Throat, Catarrh, Croup, Bronchitis, Asthma. Fcap. 8vo., 2s. 6d.

MR. FERGUSSON, F.R.S.

A SYSTEM OF PRACTICAL SURGERY; with numero trations on Wood. Fourth Edition. Fcap. 8vo. cloth, 12s. 6d.

6000-01

-14

## MR. CHURCHILL'S PUBLICATIONS.

SIR JOHN FORBES, M.D., D.C.L. (OXON.), F.R.S.

NATURE AND ART IN THE CURE OF DISEASE. Second Edition. Post 8vo. cloth, 6s.

### DR. D. J. T. FRANCIS.

CHANGE OF CLIMATE; considered as a Remedy in Dyspeptic, Pulmonary, and other Chronic Affections; with an Account of the most Eligible Places of Residence for Invalids in Spain, Portugal, Algeria, &c., at different Seasons of the Year; and an Appendix on the Mineral Springs of the Pyrences, Vichy, and Aix les Bains. Post 8vo. cloth, 8s. 6d.

C. REMIGIUS FRESENIUS.

ELEMENTARY INSTRUCTION IN CHEMICAL ANALYSIS. AS PRACTISED IN THE LABORATORY OF GIESSEN. Edited by LLOVE BULLOCK, late Student at Giessen.

> QUALITATIVE; Fourth Edition. 8vo. cloth, 9s. QUANTITATIVE. Second Edition. 8vo. cloth, 15s.

### MR. FOWNES, PH.D., F.R.S.

A MANUAL OF CHEMISTRY; with numerous Illustrations on Wood. Sixth Edition. Fcap. 8vo. cloth, 12s. 6d. Edited by H. Bence Jones, M.D., F.R.S., and A. W. HOFMANN, Ph.D., F.R.S.

CHEMISTRY, AS EXEMPLIFYING THE WISDOM AND BENEFICENCE OF GOD. Second Edition. Fcap. 8vo. cloth, 4s. 6d.

INTRODUCTION TO QUALITATIVE ANALYSIS. Post 8vo. cloth, 2s. IV.

CHEMICAL TABLES. Folio, price 2s. 6d.

### DR. FULLER.

ON RHEUMATISM, RHEUMATIC GOUT, AND SCIATICA: their Pathology, Symptoms, and Treatment. Second Edition. 8vo. cloth, 12s. 6d.

DR. GAIRDNER.

ON GOUT; its History, its Causes, and its Cure. Third Edition. Post 8vo. cloth, 8s. 6d.

#### MR. GALLOWAY.

THE FIRST STEP IN CHEMISTRY. Second Edition. Fcap. 8vo. cloth, 5s.

A MANUAL OF QUALITATIVE ANALYSIS. Second Edition. Post 8vo. cloth, 4s. 6d.

CHEMICAL DIAGRAMS. On Four large Sheets, for School and Lecture Rooms, 5s, 6d, the Set.





13

DR. GARRETT.

ON EAST AND NORTH-EAST WINDS; the Nature, Treatment, and Prevention of their Suffocating Effects. Fcap. 8vo. cloth, 4s. 6d.

MR. ROBERT GARNER, F.L.S.

EUTHERAPEIA; or, AN EXAMINATION OF THE PRINCIPLES OF MEDICAL SCIENCE, including Researches on the Nervous System. Illustrated with 9 Engravings on Copper, and Engravings on Wood. 8vo. cloth, 8s.

MR. GAY, F.R.C.S.E.

FEMORAL RUPTURE: ITS ANATOMY, PATHOLOGY, AND SURGERY. With a New Mode of Operating. 4to., Plates, 10s. 6d.

A MEMOIR ON INDOLENT ULCERS. Post 8vo. cloth, 3s. 6d.

DR. GRANVILLE, F.R.S.

ON SUDDEN DEATH. Post 8vo., 2s. 6d.

MR. GRAY, M.R.C.S.

PRESERVATION OF THE TEETH indispensable to Comfort and Appearance, Health, and Longevity. 18mo. cloth, 3s.

MR. GRIFFITHS.

CHEMISTRY OF THE FOUR SEASONS—Spring, Summer,
Autumn, Winter,
8vo. cloth, 7s, 6d,
Illustrated with Engravings on Wood. Second Edition. Foolseap

DR. GULLY.

THE WATER CURE IN CHRONIC DISEASE: an Exposition of the Causes, Progress, and Terminations of various Chronic Diseases of the Viscera, Nervous System, and Limbs, and of their Treatment by Water and other Hygienic Means. Fifth Edition. Foolscap 8vo. sewed, 2s. 6d.

THE SIMPLE TREATMENT OF DISEASE; deduced from the

DR. GUY.

HOOPER'S PHYSICIAN'S VADE-MECUM; OR, MANUAL OF THE PRINCIPLES AND PRACTICE OF PHYSIC. New Edition, considerably enlarged, and rewritten. Foolscap 8vo. cloth, 12s. 6d.

GUY'S HOSPITAL REPORTS. Third Series. Vols. I. to III., 8vo., 7s. 6d. each.

VALYSIS,

peptic Pel-

ible Planes of

ix les Bairs

Second

on Wood

F.R.S.

M AND

ro.cloth

HATICA 12s. Gal

ition\_

Fcap Sm

cond Edition

School a

( CO-01-

DR. HABERSHON.

OBSERVATIONS ON DISEASES OF THE ALIMENTARY CANAL, ŒSOPHAGUS, STOMACH, CÆCUM, and INTESTINES. 8vo. cloth, 10s, 6d.

### DR. MARSHALL HALL, F.R.S.

1.

PRONE AND POSTURAL RESPIRATION IN DROWNING AND OTHER FORMS OF APNŒA OR SUSPENDED RESPIRATION. Post 8vo, cloth. 5s.

11,

PRACTICAL OBSERVATIONS AND SUGGESTIONS IN MEDI-CINE. Post 8vo. cloth, 8s. 6d.

DITTO. Second Series. Post 8vo. cloth, 8s. 6d.

DR. O. RADGLYFFE HALL.

TORQUAY IN ITS MEDICAL ASPECT AS A RESORT FOR PULMONARY INVALIDS. Post 8vo. cloth, 5s.

MR. HARDWICH.

A MANUAL OF PHOTOGRAPHIC CHEMISTRY. Fourth Edition. Foolscap 8vo. cloth, 6s. 6d.

MR. HARE, M.R.C.S.

PRACTICAL OBSERVATIONS ON THE PREVENTION, CAUSES, AND TREATMENT OF CURVATURES OF THE SPINE; with Engravings. Third Edition. 8vo. cloth, 6s.

MR. HARRISON, F.R.C.S.

THE PATHOLOGY AND TREATMENT OF STRICTURE OF THE URETHRA. 8vo. cloth, 5s.

MR. JAMES B. HARRISON, F.R.C.S.

ON THE CONTAMINATION OF WATER BY THE POISON OF LEAD, and its Effects on the Human Body. Foolscap 8vo. cloth, 3s. 6d.

### DR. HARTWIG.

L

ON SEA BATHING AND SEA AIR. Fcap. 8vo., 2s. 6d.

ON THE PHYSICAL EDUCATION OF CHILDREN. Fcap.

MR. ALFRED HAVILAND, M.R.C.S.

CLIMATE, WEATHER, AND DISEASE; being a Sketch of the Opinions of the most celebrated Ancient and Modern Writers with regard to the Influence of Climate and Weather in producing Disease. With Four coloured Engravings. 8vo. cloth, 7s.

Kar-ot

10-->13

LIMENT IE

REGIET -OF

DROWNING

S IN MEDI

RESORT FOR

TRY. Form

EVENTION E SPINE:

ICTURE OF

HE POISON

6d.

REN. Fear

Sketch of the i to the Inflormant ingravings.

DR. A. H. HASSALL.

THE MICROSCOPIC ANATOMY OF THE HUMAN BODY, IN HEALTH AND DISEASE. Illustrated with Several Hundred Drawings in Colour. Two vols. 8vo. cloth, £1. 10s.

DR. HEADLAND.

ON THE ACTION OF MEDICINES IN THE SYSTEM.

Being the Prize Essay to which the Medical Society of London awarded the Fothergillian Gold Medal for 1852. Second Edition. 8vo. cloth, 10s.

MR. HIGGINBOTTOM, F.R.S., F.R.C.S.

AN ESSAY ON THE USE OF THE NITRATE OF SILVER IN THE CURE OF INFLAMMATION, WOUNDS, AND ULCERS. Second Edition. Price 5s.

ADDITIONAL OBSERVATIONS ON THE NITRATE OF SIL-VER; with full Directions for its Use as a Therapeutic Agent. 8vo., 2s. 6d.

MR. JOHN HILTON, F.R.S.

ON THE DEVELOPMENT AND DESIGN OF CERTAIN POR-TIONS OF THE CRANIUM. Illustrated with Plates in Lithography. 8vo. cloth, 6s.

DR. HINDS.

THE HARMONIES OF PHYSICAL SCIENCE IN RELATION
TO THE HIGHER SENTIMENTS; with Observations on Medical Studies, and on
the Moral and Scientific Relations of Medical Life. Post 8vo., cloth, 5s.

DR. DECIMUS HODGSON.

THE PROSTATE GLAND, AND ITS ENLARGEMENT IN OLD AGE. With 12 Plates. Royal 8vo., cloth, 6s.

MR. LUTHER HOLDEN, FR.C.S.

HUMAN OSTEOLOGY: with Plates, showing the Attachments of the Muscles. Second Edition. 8vo. cloth, 16s.

DR. G. CALVERT HOLLAND.

THE CONSTITUTION OF THE ANIMAL CREATION, expressed in Structural Appendages, as Hair, Horns, Tusks, and Fat. 8vo. cloth, 10s. 6d.

MR. C. HOLTHOUSE.

ON SQUINTING, PARALYTIC AFFECTIONS OF THE EYE, and CERTAIN FORMS OF IMPAIRED VISION. Fcap. 8vo. cloth, 4s. 6d.

LECTURES ON STRABISMUS, delivered at the Westminster Hospital.

10-04

DR. W. CHARLES HOOD.

SUGGESTIONS FOR THE FUTURE PROVISION OF CRIMI-NAL LUNATICS. 8vo. cloth, 5s. 6d.

MR. P. HOOD.

THE SUCCESSFUL TREATMENT OF SCARLET FEVER:
also, OBSERVATIONS ON THE PATHOLOGY AND TREATMENT OF
CROWING INSPIRATIONS OF INFANTS. Post 8vo. cloth, 5s.

DR. HOOPER.

THE MEDICAL DICTIONARY; containing an Explanation of the Terms used in Medicine and the Collateral Sciences. Eighth Edition. Edited by Klein Grant, M.D. 8vo. cloth, 30s.

MR. JOHN HORSLEY.

A CATECHISM OF CHEMICAL PHILOSOPHY; being a Familiar Exposition of the Principles of Chemistry and Physics. With Engravings on Wood. Designed for the Use of Schools and Private Teachers. Post 8vo. cloth, 6s. 6d.

DR. HUFELAND.

THE ART OF PROLONGING LIFE. A New Edition. Edited by Erasmus Wilson, F.R.S. Foolscap 8vo., 2s. 6d.

MR. W. CURTIS HUGMAN, F.R.C.S.

ON HIP-JOINT DISEASE; with reference especially to Treatment by Mechanical Means for the Relief of Contraction and Deformity of the Affected Limb. 3vo. cloth, 3s. 6d.

DR. HENRY HUNT.

ON HEARTBURN AND INDIGESTION. 8vo. cloth, 5s.

MR. T. HUNT.

DISEASES OF THE SKIN: a Guide for their Treatment and Prevention. Second Edition. Fcap. 8vo., 2s. 6d.

DR. INMAN.

THE PHENOMENA OF SPINAL IRRITATION AND OTHER FUNCTIONAL DISEASES OF THE NERVOUS SYSTEM EXPLAINED, and a Rational Plan of Treatment deduced. With Plates. 8vo. cloth, 6s.

DR. ARTHUR JACOB, F.R.C.S.

A TREATISE ON THE INFLAMMATIONS OF THE EYE-BALL.

DR. JAMES JAGO, A.B., CANTAB.; M.B., OXON.

OCULAR SPECTRES AND STRUCTURES AS MUTUAL EXPONENTS. Illustrated with Engravings on Wood. 8vo. cloth, 5s.

4600casi-

10-->0)

DR. HANDFIELD JONES, F.R.S., & DR. EDWARD H. SIEVEKING. A MANUAL OF PATHOLOGICAL ANATOMY. Illustrated with numerous Engravings on Wood. Foolscap 8vo, cloth, 12s. 6d.

### MR. WHARTON JONES, F.R.S.

A MANUAL OF THE PRINCIPLES AND PRACTICE OF OPHTHALMIC MEDICINE AND SURGERY; illustrated with Engravings, plain and coloured. Second Edition. Foolscap 8vo. cloth, 12s. 6d.

THE WISDOM AND BENEFICENCE OF THE ALMIGHTY. AS DISPLAYED IN THE SENSE OF VISION; being the Actonian Prize Essay for 1851. With Illustrations on Steel and Wood. Foolscap 8vo. cloth, 4s. 6d.

DEFECTS OF SIGHT: their Nature, Causes, Prevention, and General Management. Fcap. 8vo. 2s. 6d.

A CATECHISM OF THE MEDICINE AND SURGERY OF THE EYE AND EAR. For the Clinical Use of Hospital Students. Fcap. 8vo. 2s. 6d-

A CATECHISM OF THE PHYSIOLOGY AND PHILOSOPHY OF BODY, SENSE, AND MIND. For Use in Schools and Colleges, Fcap, 8vo., 2s. 6d.

#### DR. BENCE JONES, F.R.S.

MULDER ON WINE. Foolscap 8vo. cloth, 6s.

ON ANIMAL CHEMISTRY, in its relation to STOMACH and RENAL DISEASES. 8vo. cloth, 6s.

### MR. JUDD.

A PRACTICAL TREATISE ON URETHRITIS AND SYPHI-LIS: including Observations on the Power of the Menstruous Fluid, and of the Discharge from Leucorrhoa and Sores to produce Urethritis: with a variety of Examples, Experiments, Remedies, and Cures. 8vo. cloth, £1.5s.

### MR. KNAGGS.


UNSOUNDNESS OF MIND CONSIDERED IN RELATION TO THE QUESTION OF RESPONSIBILITY IN CRIMINAL CASES. 8vo. cloth, 4s. 6d.

### DR. LAENNEC.

A MANUAL OF AUSCULTATION AND PERCUSSION. Translated and Edited by J. B. SHARPE, M.R.C.S. 3s.

DR. HUNTER LANE, F.L.S.

A COMPENDIUM OF MATERIA MEDICA AND PHARMACY; adapted to the London Pharmacopæia, 1851, embodying all the new French, American, and Indian Medicines, and also comprising a Summary of Practical Toxicology. Second Edition. 24mo. cloth, 5s. 6d.



nows.

STEM OF

45.52

103

RCS.

19

MR. H. W. LOBB, L.S.A., M.R.C.S.E.

ON SOME OF THE MORE OBSCURE FORMS OF NERVOUS
AFFECTIONS, THEIR PATHOLOGY AND TREATMENT. With an
Introduction on the Physiology of Digestion and Assimilation, and the Generation and
Distribution of Nerve Force. Based upon Original Microscopical Observations. With
Engravings. 8vo. cloth, 10s. 6d.

LONDON MEDICAL SOCIETY OF OBSERVATION.

WHAT TO OBSERVE AT THE BED-SIDE, AND AFTER DEATH. Published by Authority. Second Edition. Foolscap 8vo. cloth, 4s. 6d.

M. LUGOL.

ON SCROFULOUS DISEASES. Translated from the French, with Additions by W. H. RANKING, M.D., Physician to the Suffolk General Hospital. 8vo. cloth, 10s. 6d.

MR. JOSEPH MACLISE, F.R.C.S.

SURGICAL ANATOMY. A Series of Dissections, illustrating the Principal Regions of the Human Body.

The Second Edition, complete in XIII. Fasciculi. Imperial folio, 5s. each; bound in cloth, £3. 12s.; or bound in morocco, £4. 4s.

II.

ON DISLOCATIONS AND FRACTURES. This Work will be Uniform with the Author's "Surgical Anatomy;" each Fasciculus will contain Four beautifully executed Lithographic Drawings, and be completed in Nine Numbers. Fasciculus I., imperial folio, 5s.

### DR. MAYNE.

AN EXPOSITORY LEXICON OF THE TERMS, ANCIENT AND MODERN, IN MEDICAL AND GENERAL SCIENCE, including a complete MEDICAL AND MEDICO-LEGAL VOCABULARY, and presenting the correct Pronunciation, Derivation, Definition, and Explanation of the Names, Analogues, Synonymes, and Phrases (in English, Latin, Greek, French, and German,) employed in Science and connected with Medicine. Parts I. to VI., price 5s. each.

DR. WM. H. MADDEN.

THOUGHTS ON PULMONARY CONSUMPTION; with an Appendix on the Climate of Torquay. Post 8vo. cloth, 5s.

DR. MARCET.

ON THE COMPOSITION OF FOOD, AND HOW IT IS ADULTERATED; with Practical Directions for its Analysis, 8vo. cloth, 6s, 6d.

DR. MARTIN.

THE UNDERCLIFF, ISLE OF WIGHT: its Climate, History, and Natural Productions. Post 3vo. cloth, 10s. 6d.

26000-01

GY, DIAG

Post Se

for the Tose

WAY

of Inveter

Second Edit

WITH

cal Obser

28-686-+OI

t, 8vo. clock 6

8vo. clock

Summer Course. Post 8vo. cloth, 4s. 6d.

TRANSACTIONS OF THE ODONTOLOGICAL SOCIETY OF LONDON. With Plates. Vol. I., 8vo. cloth, 8s. 6d.

#### MR. PAGET.

A DESCRIPTIVE CATALOGUE OF THE ANATOMICAL MUSEUM OF ST. BARTHOLOMEW'S HOSPITAL. Vol. I. Morbid Anatomy. 8vo. cloth, 5s.

DITTO. Vol. II. Natural and Congenitally Malformed Structures, and Lists of the Models, Casts, Drawings, and Diagrams. 5s.

THE PRESCRIBER'S PHARMACOPEIA; containing all the Medicines in the London Pharmacopeia, arranged in Classes according to their Action, with their Composition and Doses. By a Practising Physician. Fourth Edition. 32mo. cloth, 2s. 6d.; roan tuck (for the pocket), 3s. 6d.

### DR. JOHN ROWLISON PRETTY.

AIDS DURING LABOUR, including the Administration of Chloroform, the Management of Placenta and Post-partum Hæmorrhage. Fcap. 8vo. cloth, 4s. 6d.

### SIR WM. PYM, K.C.H.

OBSERVATIONS UPON YELLOW FEVER, with a Review of "A Report upon the Diseases of the African Coast, by Sir Wm. Burnett and Dr. Bryson," proving its highly Contagious Powers. Post 8vo. 6s.

### DR. RADCLIFFE.

EPILEPSY, AND OTHER AFFECTIONS OF THE NERVOUS SYSTEM which are marked by Tremor, Convulsion, or Spasm: their Pathology and Treatment. 8vo. cloth, 5s.

### DR. F. H. RAMSBOTHAM.

THE PRINCIPLES AND PRACTICE OF OBSTETRIC MEDI-CINE AND SURGERY. Illustrated with One Hundred and Twenty Plates on Steel and Wood; forming one thick handsome volume. Fourth Edition. 8vo. cloth, 22s.

#### DR. RAMSBOTHAM.

PRACTICAL OBSERVATIONS ON MIDWIFERY, with a Selection of Cases. Second Edition. 8vo. cloth, 12s.

#### DR. DU BOIS REYMOND.

ANIMAL ELECTRICITY; Edited by H. Bence Jones, M.D., F.R.S. With Fifty Engravings on Wood. Foolscap 8vo. cloth, 6s.

### DR. REYNOLDS.

THE DIAGNOSIS OF DISEASES OF THE BRAIN, SPINAL CORD, AND THEIR APPENDAGES. 8vo. cloth, 8s.

### DR. B. W. RICHARDSON.

ON THE CAUSE OF THE COAGULATION OF THE BLOOD.

Being the ASTLEY COOPER PRIZE ESSAY for 1856. With a Practical Appendix.

8vo. cloth, 16s.

THE HYGIENIC TREATMENT OF PULMONARY CONSUMP-

1-2010

Econe - DI

DR. SCHACHT.

THE MICROSCOPE, AND ITS APPLICATION TO VEGETABLE ANATOMY AND PHYSIOLOGY. Edited by FREDERICK CURREY, M.A. Fcap. 8vo. cloth, 6s.

OFF

(75 E)

OMEN

MR. SHAW, M.R.C.S.

THE MEDICAL REMEMBRANCER; OR, BOOK OF EMER-GENCIES: in which are concisely pointed out the Immediate Remedies to be adopted in the First Moments of Danger from Poisoning, Drowning, Apoplexy, Burns, and other Accidents; with the Tests for the Principal Poisons, and other useful Information. Fourth Edition. Edited, with Additions, by Jonathan Hutchinson, M.R.C.S. 32mo.

DR. SIBSON, F.R.S.

MEDICAL ANATOMY. With coloured Plates. Imperial folio. Fasciculi I. to V. 5s. each.

DR. E. H. SIEVEKING.

ON EPILEPSY AND EPILEPTIFORM SEIZURES: their Causes, Pathology, and Treatment. Post 8vo. cloth, 7s. 6d.

MR. SKEY, F.R.S.

OPERATIVE SURGERY: with Illustrations engraved on Wood. 8vo. cloth, 12s. 6d.

DR. SMELLIE.

OBSTETRIC PLATES: being a Selection from the more Important and Practical Illustrations contained in the Original Work. With Anatomical and Practical Directions. 8vo. cloth. 5s.

MR. HENRY SMITH, F.R.C.S.

ON STRICTURE OF THE URETHRA. 8vo. cloth, 7s. 6d.

DR. W. TYLER SMITH.

A MANUAL OF OBSTETRICS, THEORETICAL AND PRAC-TICAL. Illustrated with 186 Engravings. Fcap. 8vo. cloth, 12s. 6d.

THE PATHOLOGY AND TREATMENT OF LEUCORRHEA. With Engravings on Wood. 8vo. cloth, 7s.

THE PERIODOSCOPE, a new Instrument for determining the Date of Labour, and other Obstetric Calculations, with an Explanation of its Uses, and an Essay on the Periodic Phenomena attending Pregnancy and Parturition. 8vo. cloth, 4s.

> DR. SNOW. MODE OF COMMUNICATION OF CHOLERA. nuch Enlarged, and Illustrated with Maps. 8vo. cloth, 7s.



# MR. CHURCHILL'S PUBLICATIONS.

MR. STOWE, M.R.C.S.

A TOXICOLOGICAL CHART, exhibiting at one view the Symptoms, Treatment, and Mode of Detecting the various Poisons, Mineral, Vegetable, and Animal. To which are added, concise Directions for the Treatment of Suspended Animation. Eleventh Edition. On Sheet, 2s.; mounted on Roller, 5s.

MR. TAMPLIN, F.R.C.S.E.

LATERAL CURVATURE OF THE SPINE: its Causes, Nature, and Treatment. 8vo. cloth, 4s.

DR. ALFRED S. TAYLOR, F.R.S.

A MANUAL OF MEDICAL JURISPRUDENCE. Sixth Edition. Fcap. 8vo. cloth, 12s. 6d.

ON POISONS, in relation to MEDICAL JURISPRUDENCE AND MEDICINE. Fcap. 8vo. cloth, 12s. 6d.

DR. THEOPHILUS THOMPSON, F.R.S.

CLINICAL LECTURES ON PULMONARY CONSUMPTION.
With Plates. 8vo. cloth, 7s. 6d.

LETTSOMIAN LECTURES ON PULMONARY CONSUMPTION; with Remarks on Microscopical Indications, and on Cocoa-nut Oil. Post 8vo., 2s. 6d.

DR. THOMAS.

THE MODERN PRACTICE OF PHYSIC; exhibiting the Symptoms, Causes, Morbid Appearances, and Treatment of the Diseases of all Climates. Eleventh Edition. Revised by Algernon Frampton, M.D. 2 vols. 8vo. cloth, 28s.

HENRY THOMPSON, M.B. LOND., F.R.C.S.

STRICTURE OF THE URETHRA; its Pathology and Treatment.

The last Jacksonian Treatise of the Royal College of Surgeons. With Plates. 8vo. cloth, 10s.

DR. TILT.

ON DISEASES OF WOMEN AND OVARIAN INFLAM-MATION IN RELATION TO MORBID MENSTRUATION, STERILITY, PELVIC TUMOURS, AND AFFECTIONS OF THE WOMB. Second Edition. 8vo. cloth, 9s.

THE CHANGE OF LIFE IN HEALTH AND DISEASE:

Practical Treatise on the Nervous and other Affections incidental to Women at the Decl.

of Life. Second Edition. 8vo. cloth, 6s.

)

-10-3

DR. VAN OVEN.

ON THE DECLINE OF LIFE IN HEALTH AND DISEASE; being an Attempt to Investigate the Causes of LONGEVITY, and the Best Means of Attaining a Healthful Old Age. 8vo. cloth, 10s. 6d.

### MR. WADE, F.R.C.S.

STRICTURE OF THE URETHRA; its Complications and Effects. With Practical Observations on its Causes, Symptoms, and Treatment; and on a Safe and Efficient Mode of Treating its more Intractable Forms. 8vo. cloth, 5s.

### DR. WALLER.

ELEMENTS OF PRACTICAL MIDWIFERY; or, Companion to the Lying-in Room. Fourth Edition, with Plates. Fcap. cloth, 4s. 6d.

MR. HAYNES WALTON, F.R.C.S.

OPERATIVE OPHTHALMIC SURGERY. With Engravings on Wood. 8vo. cloth, 18s.

DR. WARDROP.

ON DISEASES OF THE HEART. 8vo. cloth, 12s.

DR. EBEN. WATSON, A.M.

ON THE TOPICAL MEDICATION OF THE LARYNX IN CERTAIN DISEASES OF THE RESPIRATORY AND VOCAL ORGANS. 8vo. cloth, 5s.

DR. WEBER.

A CLINICAL HAND-BOOK OF AUSCULTATION AND PER-CUSSION. Translated by JOHN COCKLE, M.D. 5s.

DR. WEGG.

OBSERVATIONS RELATING TO THE SCIENCE AND ART OF MEDICINE. 8vo. cloth, 8s.

DR. WEST.

LECTURES ON THE DISEASES OF WOMEN. 8vo. cloth, 10s. 6d-

MR. ERASMUS WILSON, F.R.S.

THE ANATOMIST'S VADE-MECUM: A SYSTEM OF HUMAN ANATOMY. With numerous Illustrations on Wood. Seventh Edition. Foolscap 8vo. cloth, 12s. 6d.

DISEASES OF THE SKIN: A Practical and Theoretical Treatise on the DIAGNOSIS, PATHOLOGY, and TREATMENT OF CUTANEOUS DIS-EASES. Fourth Edition. 8vo. cloth, 16s.

The same Work; illustrated with finely executed Engravings on Steel, accurately coloured. 8vo. cloth, 34s.

HEALTHY SKIN: A Treatise on the Management of the Skin and Hair in relation to Health. Fifth Edition. Foolscap 8vo. 2s. 6d.

PORTRAITS OF DISEASES OF THE SKIN. Folio. Fasciculi I. to XII., completing the Work. 20s. each.

ON SYPHILIS, CONSTITUTIONAL AND HEREDITARY;
AND ON SYPHILITIC ERUPTIONS. With Four Coloured Plates. 8vo. cloth,

A THREE WEEKS' SCAMPER THROUGH THE SPAS OF GERMANY AND BELGIUM, with an Appendix on the Nature and Uses of Mineral Waters. Post 8vo. cloth, 6s. 6d.

DR. FORBES WINSLOW, D.C.L. OXON.

LETTSOMIAN LECTURES ON INSANITY. 8vo. cloth, 5s.

A SYNOPSIS OF THE LAW OF LUNACY; as far as it relates to the Organization and Management of Private Asylums for the Care and Treatment of the Insane. In the form of a Chart, varnished, mounted on canvas and rollers, price 6s.

### DR. G. C. WITTSTEIN.

PRACTICAL PHARMACEUTICAL CHEMISTRY: An Explanation of Chemical and Pharmaceutical Processes, with the Methods of Testing the Purity of the Preparations, deduced from Original Experiments. Translated from the Second German Edition, by STEPHEN DARBY. 18mo. cloth, 6s.

#### MR. YEARSLEY.

DEAFNESS PRACTICALLY ILLUSTRATED; being an Exposition of Original Views as to the Causes and Treatment of Diseases of the Ear. Fifth Edition. Foolscap 8vo., 2s. 6d.

ON THE ENLARGED TONSIL AND ELONGATED UVULA, and other Morbid Conditions of the Throat, Sixth Edition. 8vo. cloth, 5s.

1

S OF 1

OFFSPRING

TION to the Plant Leucore here

ow of the brief R cloth

nding cloth

AND LEGI

tion. P. 8

\$ 6000 - or

10-->

